

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
28 December 2000 (28.12.2000)

PCT

(10) International Publication Number  
**WO 00/78953 A2**

(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
5/10, C07K 14/47, 14/705, 16/18, 16/28, C12Q 1/68,  
A61K 38/17, G01N 33/50, A01K 67/027

(21) International Application Number: PCT/US00/16668

(22) International Filing Date: 16 June 2000 (16.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/139,923 17 June 1999 (17.06.1999) US  
60/148,177 10 August 1999 (10.08.1999) US  
60/149,357 18 August 1999 (18.08.1999) US  
60/162,287 28 October 1999 (28.10.1999) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier applications:

US 60/139,923 (CIP)  
Filed on 17 June 1999 (17.06.1999)  
US 60/148,177 (CIP)  
Filed on 10 August 1999 (10.08.1999)  
US 60/149,357 (CIP)  
Filed on 18 August 1999 (18.08.1999)  
US 60/162,287 (CIP)  
Filed on 28 October 1999 (28.10.1999)

(71) Applicant (for all designated States except US): INCYTE GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LAL, Preeti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). YANG, Junming [CN/US]; 7125 Bark Lane, San Jose, CA 95129 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). HILLMAN, Jennifer, L.

[US/US]; 230 Monroe Drive, #12, Mountain View, CA 94040 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). BURFORD, Neil [GB/US]; 1308 4th Avenue, San Francisco, CA 94122 (US). BAUGHN, Mariab, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). AZIMZAI, Yalda [US/US]; 2045 Rocksprings Drive, Hayward, CA 94545 (US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US). AU-YOUNG, Janice [US/US]; 233 Golden Eagle Lane, Brisbane, CA 94005 (US). PATTERSON, Chandra [US/US]; 490 Sherwood Way, #1, Menlo Park, CA 94025 (US).

(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics, Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).

(81) Designated States (national): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HUMAN TRANSPORT PROTEINS

(57) Abstract: The invention provides human transport proteins (TPPT) and polynucleotides which identify and encode TPPT. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of TPPT.

WO 00/78953 A2

## HUMAN TRANSPORT PROTEINS

### TECHNICAL FIELD

5 This invention relates to nucleic acid and amino acid sequences of human transport proteins and to the use of these sequences in the diagnosis, treatment, and prevention of transport, metabolic, neurological, cardiovascular, reproductive, and immune disorders, and cell proliferative disorders including cancer.

### 10 BACKGROUND OF THE INVENTION

Eukaryotic cells are surrounded and subdivided into functionally distinct organelles by hydrophobic lipid bilayer membranes. These membranes act as a barrier to most molecules, and maintain the essential differences between the cytosol, the extracellular environment, and the contents of each intracellular organelle. Transport of essential nutrients, certain metal ions, metabolic waste products, cell signaling molecules, macromolecules, and proteins across lipid membranes and between organelles must be mediated by a variety of transport molecules. Transport between the cytoplasm and the extracellular environment, and between the cytoplasm and luminal spaces of cellular organelles requires specific transport proteins. Each transport protein carries a particular class of molecule, such as ions, sugars, or amino acids, and often is specific to a certain molecular species of the class.

Cells and organelles require transport molecules to import and export essential nutrients and metal ions including  $K^+$ ,  $NH_4^+$ ,  $P_i$ ,  $SO_4^{2-}$ , sugars, and vitamins, as well as various metabolic waste products. Transport proteins also play roles in antibiotic resistance, toxin secretion, ion balance, synaptic neurotransmission, kidney function, intestinal absorption, tumor growth, and other diverse cell functions (Griffith, J. and C. Sansom (1998) The Transporter Facts Book, Academic Press, San Diego CA, pp. 3-29). Transport can occur by a passive concentration-dependent mechanism, or can be linked to an energy source such as ATP hydrolysis or an ion gradient. Proteins that function in transport include carrier proteins, which bind to a specific solute and undergo a conformational change that transfers the bound solute across the membrane, and channel proteins, which form hydrophilic pores that allow specific solutes to diffuse through the membrane down an electrochemical solute gradient.

Transport proteins are often multi-pass transmembrane proteins, which either actively transport molecules across the membrane or passively allow them to cross. Active transport involves directional pumping of a solute across the membrane, usually against an electrochemical gradient. Active transport is tightly coupled to a source of metabolic energy, such as ATP hydrolysis or an electrochemically favorable ion gradient. Passive transport involves the movement of a solute down

its electrochemical gradient. Transport proteins can be further classified as either carrier proteins or channel proteins. Carrier proteins, which can function in active or passive transport, bind to a specific solute to be transported and undergo a conformational change which transfers the bound solute across the membrane. Channel proteins, which only function in passive transport, form hydrophilic pores across the membrane. When the pores open, specific solutes, such as inorganic ions, pass through the membrane and down the electrochemical gradient of the solute. Examples include facilitative transporters, the secondary active symporters and antiporters driven by ion gradients, and active ATP binding cassette transporters involved in multiple-drug resistance and targeting of antigenic peptides to MHC Class I molecules. Transported substrates range from nutrients and ions to a broad variety of drugs, peptides and proteins.

Information on the action of ARL-6 (ADP-ribosylation like factor), an endoplasmic reticulum transmembrane protein, can be found in Greenfield, J.J. and S. High (1999; *J. Cell Sci.* 112:1477-1486). Information on reduced folate carrier transporter proteins can be found in Dixon, K.H. et al. (1994; *J. Biol. Chem.* 269:17-20) and Moscow, J.A. et al. (1995; *Cancer Res.* 55:5983-5987).

Carrier proteins which transport a single solute from one side of the membrane to the other are called uniporters. In contrast, coupled transporters link the transfer of one solute with simultaneous or sequential transfer of a second solute, either in the same direction (symport) or in the opposite direction (antiport). For example, intestinal and kidney epithelia contain a variety of symporter systems driven by the sodium gradient that exists across the plasma membrane. Sodium moves into the cell down its electrochemical gradient and brings the solute into the cell with it. The sodium gradient that provides the driving force for solute uptake is maintained by the ubiquitous  $\text{Na}^+/\text{K}^+$  ATPase. Sodium-coupled transporters include the mammalian glucose transporter (SGLT1), iodide transporter (NIS), and multivitamin transporter (SMVT). All three transporters have twelve putative transmembrane segments, extracellular glycosylation sites, and cytoplasmically-oriented N- and C-termini.

Mitochondrial carrier proteins are transmembrane-spanning proteins which transport ions and charged metabolites between the cytosol and the mitochondrial matrix. Examples include the ADP, ATP carrier protein; the 2-oxoglutarate/malate carrier; the phosphate carrier protein; the brown fat uncoupling protein which transports protons from the cytosol into the matrix; the pyruvate carrier; the dicarboxylate carrier which transports malate, succinate, fumarate, and phosphate; the tricarboxylate carrier which transports citrate and malate; and the Grave's disease carrier protein, a protein recognized by IgG in patients with active Grave's disease, an autoimmune disorder resulting in hyperthyroidism (Stryer, L. (1995) Biochemistry, W.H. Freeman and Company, New York NY, p. 551; PROSITE PDOC00189 Mitochondrial energy transfer proteins signature; Online Mendelian Inheritance in Man (OMIM) \*275000 Graves Disease).

This class of transporters also includes the mitochondrial uncoupling proteins, which create

proton leaks across the inner mitochondrial membrane, thus uncoupling oxidative phosphorylation from ATP synthesis. The result is energy dissipation in the form of heat. Mitochondrial uncoupling proteins have been implicated as modulators of thermoregulation and metabolic rate, and have been proposed as potential targets for drugs against metabolic diseases such as obesity (Ricquier, D. et al. (1999) *J. Int. Med.* 245:637-642).

A number of metal ions such as iron, zinc, copper, cobalt, manganese, molybdenum, selenium, nickel, and chromium are important as cofactors for a number of enzymes. For example, zinc is required for the function of enzymes such as the extracellular matrix metalloproteinases, and zinc ions stabilize several motifs commonly found in transcription factors, including zinc fingers, zinc clusters, and LIM domains. Zinc and other metal ions must be provided in the diet, and are absorbed by transporters in the gastrointestinal tract. Plasma proteins transport the metal ions to the liver and other target organs, where specific transporters move the ions into cells and cellular organelles as needed. Imbalances in metal ion metabolism have been associated with a number of disease states (Danks, D.M. (1986) *J. Med. Genet.* 23:99-106).

The largest and most diverse family of transport proteins known are the ATP-binding cassette (ABC) transporters. As a family, ABC transporters can transport substances that differ markedly in chemical structure and size, ranging from small molecules such as ions, sugars, amino acids, peptides, and phospholipids, to lipopeptides, large proteins, and complex hydrophobic drugs. ABC proteins consist of four modules: two nucleotide-binding domains (NBD), which hydrolyze ATP to supply the energy required for transport, and two membrane-spanning domains (MSD), each containing six putative transmembrane segments. These four modules may be encoded by a single gene, as is the case for the cystic fibrosis transmembrane regulator (CFTR), or by separate genes. When encoded by separate genes, each gene product contains a single NBD and MSD. These "half-molecules" form homo- and heterodimers, such as Tap1 and Tap2, the endoplasmic reticulum-based major histocompatibility (MHC) peptide transport system. Several genetic diseases are attributed to defects in ABC transporters, such as the following diseases and their corresponding proteins: cystic fibrosis (CFTR, an ion channel), adrenoleukodystrophy (adrenoleukodystrophy protein, ALDP), Zellweger syndrome (peroxisomal membrane protein-70, PMP70), and hyperinsulinemic hypoglycemia (sulfonylurea receptor, SUR). Overexpression of the multidrug resistance (MDR) protein, another ABC transporter, in human cancer cells makes the cells resistant to a variety of cytotoxic drugs used in chemotherapy (Taglicht, D. and S. Michaelis (1998) *Methods Enzymol.* 292:131-163).

The nuclear pore complex (NPC) is a large multiprotein complex spanning the nuclear envelope which mediates the transport of proteins and RNA molecules between the nucleus and the cytoplasm, thus contributing to the regulation of gene expression. The NPC allows passive diffusion of ions, small molecules, and macromolecules under about 60kD, while larger macromolecules are transported by facilitated, energy-dependent pathways. Nuclear localization signals (NLS), consisting



of short stretches of amino acids enriched in basic residues, are found on proteins that are targeted to the nucleus, such as the glucocorticoid receptor. The NLS is recognized by the NLS receptor, importin, which then interacts with the monomeric GTP-binding protein Ran. This NLS protein/receptor/Ran complex navigates the nuclear pore with the help of the homodimeric protein nuclear transport factor 2 (NTF2) (Nakielnny, S. and G. Dreyfuss (1997) *Curr. Opin. Cell Biol.* 9:420-429; Gorlich, D. (1997) *Curr. Opin. Cell Biol.* 9:412-419). Four O-linked glycoproteins, p62, p58, p54, and p45, exist as a stable "p62 complex" that forms a ring localized on both nucleoplasmic and cytoplasmic surfaces of the NPC. The p62, p58, and p54 proteins all interact directly with the cytosolic transport factors p97 and NTF2, suggesting that the p62 complex is an important ligand binding site near the central gated channel of the NPC (Hu, T. et al. (1996) *J. Cell Biol.* 134:589-601).

Transport can also occur through intercellular bridges which connect the cytoplasms of sister cells, for example in the male and female germline of species ranging from fruit flies to humans. These bridges allow passage of cytoplasmic materials between cells during development. Intercellular bridges have also been found to connect somatic cells. The nurse cells and oocyte of a *Drosophila* egg chamber, which are derived from a single precursor cell through four rounds of mitosis, are connected to each other through intercellular bridges called ring canals. The cells do not completely separate after mitosis; the mitotic cleavage furrows are transformed into ring canals by the addition of an actin cytoskeleton lining the tunnels between the cells. The *Drosophila* kelch protein functions in organizing actin in the ring canal. Mutations in kelch cause female sterility in *Drosophila*. Kelch contains four protein domains: the NTR domain at the N-terminus, the BTB or POZ domain, the IVR or intervening region; and the kelch repeat domain, which contains six 50-amino acid kelch repeats. The BTB or POZ domain, a 120-amino acid motif that is also found in several zinc-finger containing transcription factors, may be important in dimerization of kelch. Kelch repeats are found in other proteins as well and may be important for actin binding (Robinson, D.N. and L. Cooley (1997) *J. Cell Biol.* 138:799-810; Cooley, L. (1998) *Cell* 93:913-915).

### Ion Channels

The electrical potential of a cell is generated and maintained by controlling the movement of ions across the plasma membrane. The movement of ions requires ion channels, which form an ion-selective pore within the membrane. Ion channels share common structural and mechanistic themes. The channel consists of four or five subunits or protein monomers that are arranged like a barrel in the plasma membrane. Each subunit typically consists of six potential transmembrane segments (S1, S2, S3, S4, S5, and S6). The center of the barrel forms a pore lined by  $\alpha$ -helices or  $\beta$ -strands. The side chains of the amino acid residues comprising the  $\alpha$ -helices or  $\beta$ -strands establish the charge (cation or anion) selectivity of the channel. The degree of selectivity, or what specific ions are allowed to pass through the channel, depends on the diameter of the narrowest part of the pore. There

are two basic types of ion channels, ion transporters and gated ion channels. Ion transporters utilize the energy obtained from ATP hydrolysis to actively transport an ion against the ion's concentration gradient. Gated ion channels allow passive flow of an ion down the ion's electrochemical gradient under restricted conditions. Together, these types of ion channels generate, maintain, and utilize an electrochemical gradient that is used in 1) electrical impulse conduction down the axon of a nerve cell, 2) transport of molecules into cells against concentration gradients, 3) initiation of muscle contraction, and 4) endocrine cell secretion.

Transmembrane ATPases are divided into three families. The phosphorylated (P) class ion transporters, including  $\text{Na}^+\text{-K}^+$  ATPase,  $\text{Ca}^{2+}$  ATPase,  $\text{H}^+$  ATPase, and  $\text{Cu}^{++}$  ATPase, are activated by a phosphorylation event. P-class ion transporters are responsible for maintaining resting potential distributions such that cytosolic concentrations of  $\text{Na}^+$  and  $\text{Ca}^{2+}$  are low and cytosolic concentration of  $\text{K}^+$  is high. The vacuolar (V) class of ion transporters include  $\text{H}^+$  pumps on intracellular organelles, such as lysosomes and Golgi. V-class ion transporters are responsible for generating the low pH within the lumen of these organelles that is required for function. The coupling factor (F) class consists of  $\text{H}^+$  pumps in the mitochondria. F-class ion transporters utilize a proton gradient to generate ATP from ADP and inorganic phosphate ( $\text{P}_i$ ).

$\text{Cu}^{++}$  ATPases export copper from cells (PROSITE PDOC00139 E1-E2 ATPases phosphorylation site). Mutations in one  $\text{Cu}^{++}$  ATPase cause Wilson disease, in which toxic amounts of copper accumulate in a number of organs, particularly the liver and brain (Tanzi, R.E. et al. (1993) Nat. Genet. 5:344-350). Mutations in another  $\text{Cu}^{++}$  ATPase cause Menkes disease and occipital horn syndrome. Menkes disease mutations block export of copper from the gastrointestinal tract, leading to skeletal abnormalities, severe mental retardation, neurologic degeneration, and mortality in early childhood (Harrison, M.D. and C.T. Dameron (1999) J. Biochem. Mol. Toxicol. 13:93-106). Occipital horn syndrome mutations cause connective tissue defects (Harrison, supra; Levinson, B. et al. (1996) Hum. Mol. Genet. 5:1737-1742).

The coupling factor (F) class of ion transporters consists of  $\text{H}^+$  pumps in mitochondria, chloroplasts, and bacteria. For example, the  $\text{F}_0\text{F}_1$  ATPase utilizes a proton gradient across the inner mitochondrial membrane to generate ATP from ADP and inorganic phosphate ( $\text{P}_i$ ). The  $\text{F}_0\text{F}_1$  ATPase is composed of the  $\text{F}_0$  complex, which is the transmembrane channel through which protons flow, and the  $\text{F}_1$  complex, where ATP synthesis activity resides.  $\text{F}_0$  has three subunits, A (also known as protein 6), B, and C (Lodish, H. et al. (1995) Molecular Cell Biology, Scientific American Books, New York NY, pp. 752-756; PROSITE PDOC00420 ATP synthase a subunit signature).

Voltage-gated  $\text{Ca}^{2+}$  channels are involved in presynaptic neurotransmitter release, and heart and skeletal muscle contraction. The voltage-gated  $\text{Ca}^{2+}$  channels from skeletal muscle (L-type) and brain (N-type) have been purified and, though their functions differ dramatically, they have similar subunit compositions. The channels are composed of three subunits. The  $\alpha_1$  subunit forms the

membrane pore and voltage sensor, while the  $\alpha_2\delta$  and  $\beta$  subunits modulate the voltage-dependence, gating properties, and the current amplitude of the channel. These subunits are encoded by at least six  $\alpha_1$ , one  $\alpha_2\delta$ , and four  $\beta$  genes. A fourth subunit,  $\gamma$ , has been identified in skeletal muscle (Walker, D. et al. (1998) *J. Biol. Chem.* 273:2361-2367; and Jay, S.D. et al. (1990) *Science* 248:490-492). The human  $\beta 4$  subunit is homologous to the mouse epilepsy gene lethargic, and is a candidate for involvement in neurological disorders including ataxia and absence epilepsy (Escayg, A. et al. (1998) *Genomics* 50:14-22).

Ligand-gated channels open their pores when an extracellular or intracellular mediator binds to the channel. Neurotransmitter-gated channels are channels that open when a neurotransmitter binds to their extracellular domain. These channels exist in the postsynaptic membrane of nerve or muscle cells. There are two types of neurotransmitter-gated channels. Sodium channels open in response to excitatory neurotransmitters, such as acetylcholine, glutamate, and serotonin. This opening causes an influx of  $\text{Na}^+$  and produces the initial localized depolarization that activates the voltage-gated channels and starts the action potential. Chloride channels open in response to inhibitory neurotransmitters, such as  $\gamma$ -aminobutyric acid (GABA) and glycine, leading to hyperpolarization of the membrane and the subsequent generation of an action potential.

Ion channels are expressed in a number of tissues where they are implicated in a variety of processes. CNG channels, while abundantly expressed in photoreceptor and olfactory sensory cells, are also found in kidney, lung, pineal, retinal ganglion cells, testis, aorta, and brain. Calcium-activated  $\text{K}^+$  channels may be responsible for the vasodilatory effects of bradykinin in the kidney and for shunting excess  $\text{K}^+$  from brain capillary endothelial cells into the blood. They are also implicated in repolarizing granulocytes after agonist-stimulated depolarization (Ishi, T.M. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:11651-11656). Another transmembrane protein, the leukotriene B<sub>4</sub> receptor (BLT) appears to be involved in inflammation responses and host cell defense against infection. BLT also functions as an HIV coreceptor (Izumi, T. et al. (1997) *Nature* 387:620-624; Martin, V. et al. (1999) *J. Biol. Chem.* 274:8597-8603).

Ion channels have been the target for many drug therapies. Neurotransmitter-gated channels have been targeted in therapies for treatment of insomnia, anxiety, depression, and schizophrenia. Voltage-gated channels have been targeted in therapies for arrhythmia, ischemic stroke, head trauma, and neurodegenerative disease (Taylor, C.P. and L.S. Narasimhan (1997) *Adv. Pharmacol.* 39:47-98).

$\text{K}^+$  channels are located in all cell types, and may be regulated by voltage, ATP concentration, or second messengers such as  $\text{Ca}^{++}$  and cAMP. In non-excitabile tissue,  $\text{K}^+$  channels are involved in protein synthesis, control of endocrine secretions, and the maintenance of osmotic equilibrium across membranes. In neurons and other excitable cells, in addition to regulating action potentials and repolarizing membranes,  $\text{K}^+$  channels are responsible for setting resting membrane potential. The cytosol contains non-diffusible anions and, to balance this net negative charge, the cell

contains a  $\text{Na}^+\text{-K}^+$  pump and ion channels that provide the redistribution of  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Cl}^-$ . The pump actively transports  $\text{Na}^+$  out of the cell and  $\text{K}^+$  into the cell in a 3:2 ratio. Ion channels in the plasma membrane allow  $\text{K}^+$  and  $\text{Cl}^-$  to flow by passive diffusion. Because of the high negative charge within the cytosol,  $\text{Cl}^-$  flows out of the cell. The flow of  $\text{K}^+$  is balanced by an electromotive force pulling  $\text{K}^+$  into the cell, and a  $\text{K}^+$  concentration gradient pushing  $\text{K}^+$  out of the cell. Thus, the resting membrane potential is primarily regulated by  $\text{K}^+$  flow (Salkoff, L. and T. Jegla (1995) *Neuron* 15:489-492). Information on NY-REN-45, a  $\text{K}^+$  channel integral membrane protein, can be found in Scanlan, M.J. et al. (1998; *Int. J. Cancer* 76:652-658). The emopamil-binding protein (EBP) shares structural features with both pro- and eukaryotic drug transport proteins (Hanner, M. et al. (1995) *J. Biol. Chem.* 270:7551-7557). The  $\text{Na}^+$  channel, transmembrane protein myelin protein zero (MPZ) may be responsible for some sporadic cases of Dejerine-Scott's disease (hereditary motor and sensory neuropathy type III) (Hayasaka, K. et al. (1993) *Nat. Genet.* 5:266-268).

$\text{K}^+$  pore-forming subunits generally have six transmembrane-spanning domains with a short region between the fifth and sixth transmembrane regions that senses membrane potential; and the amino and carboxy termini are located intracellularly. In mammalian heart, the duration of ventricular action potential is controlled by a  $\text{K}^+$  current. Thus, the  $\text{K}^+$  channel is central to the control of heart rate and rhythm.  $\text{K}^+$  channel dysfunctions are associated with a number of renal diseases including hypertension, hypokalemia, and the associated Bartter's syndrome and Getelman's syndrome, as well as neurological disorders including epilepsy.  $\text{K}^+$  channels have been implicated in Alzheimer's disease by observations that a significant component of senile plaques, beta amyloid or A beta, also blocks voltage-gated potassium channels in hippocampal neurons (Antes, L.M. et al. (1998) *Seminars Nephrol.* 18:31-45; Stoffel, M. and L.Y. Jan (1998) *Nat. Genet.* 18:6-8; Madeja, M. et al. (1997) *Eur. J. Neurosci.* 9:390-395; Good, T.A. et al. (1996) *Biophys. J.* 70:296-304).

Gated ion channels control ion flow by regulating the opening and closing of pores. These channels are categorized according to the manner of regulating the gating function. Mechanically-gated channels open pores in response to mechanical stress, voltage-gated channels open pores in response to changes in membrane potential, and ligand-gated channels open pores in the presence of a specific ion, nucleotide, or neurotransmitter.

Voltage-gated  $\text{Na}^+$  channels are responsible for electrical excitability of neurons, skeletal muscle, heart, and neuroendocrine tissues. For example, the sequential opening and closing of voltage-gated  $\text{Na}^+$  channels results in the propagation of action potentials down neuronal axons.  $\text{Na}^+$  channels isolated from rat brain tissue are heterotrimeric complexes composed of a 260 kDa pore forming  $\alpha$  subunit that associates with two smaller auxiliary subunits,  $\beta 1$  and  $\beta 2$ . The  $\beta 2$  subunit is an integral membrane glycoprotein that contains an extracellular Ig domain, and its association with  $\alpha$  and  $\beta 1$  subunits correlates with increased function of the channel, a change in the channel's gating properties, as well as an increase in whole cell capacitance (Isom, L.L. et al. (1995) *Cell* 83:433-442).

### Integral Membrane Proteins

The majority of known integral membrane proteins are transmembrane proteins (TM) which are characterized by an extracellular, a transmembrane, and an intracellular domain. TM domains are typically comprised of 15 to 25 hydrophobic amino acids which are predicted to adopt an  $\alpha$ -helical conformation. TM proteins are classified as bitopic (Types I and II) and polytopic (Types III and IV) (Singer, S.J. (1990) *Annu. Rev. Cell Biol.* 6:247-96). Bitopic proteins span the membrane once while polytopic proteins contain multiple membrane-spanning segments. TM proteins that act as cell-surface receptor proteins involved in signal transduction include growth and differentiation factor receptors, and receptor-interacting proteins such as *Drosophila* pecanex and frizzled proteins, LIV-1 protein, NF2 protein, and GNS1/SUR4 eukaryotic integral membrane proteins. TM proteins also act as transporters of ions or metabolites, such as gap junction channels (connexins) and ion channels, and as cell anchoring proteins, such as lectins, integrins, and fibronectins. TM proteins act as vesicle organelle-forming molecules, such as calveolins, or as cell recognition molecules, such as cluster of differentiation (CD) antigens, glycoproteins, and mucins. Information on connexin can be found in Kanter, H.L. et al. (1994; *J. Mol. Cell. Cardiol.* 26:861-868).

Many membrane proteins (MPs) contain amino acid sequence motifs that target these proteins to specific subcellular sites. Examples of these motifs include PDZ domains, KDEL, RGD, NGR, and GSL sequence motifs, von Willebrand factor A (vWFA) domains, and EGF-like domains. RGD, NGR, and GSL motif-containing peptides have been used as drug delivery agents in cancer treatments which target tumor vasculature (Arap, W. et al. (1998) *Science*, 279:377-380.) Furthermore, MPs may also contain amino acid sequence motifs, such as the carbohydrate recognition domain (CRD), also known as the C-type lectin domain, that mediate interactions with extracellular or intracellular molecules.

G-protein coupled receptors (GPCR) comprise a superfamily of integral membrane proteins which transduce extracellular signals. GPCRs include receptors for biogenic amines, lipid mediators of inflammation, peptide hormones, and sensory signal mediators. The structure of these highly-conserved receptors consists of seven hydrophobic transmembrane regions, an extracellular N-terminus, and a cytoplasmic C-terminus. Three extracellular loops alternate with three intracellular loops to link the seven transmembrane regions. The most conserved parts of these proteins are the transmembrane regions and the first two cytoplasmic loops. Cysteine disulfide bridges connect the second and third extracellular loops. A conserved, acidic-Arg-aromatic residue triplet present in the second cytoplasmic loop may interact with G proteins. A GPCR consensus pattern is characteristic of most proteins belonging to this superfamily (ExPASy PROSITE document PS00237; and Watson, S. and S. Arkinstall (1994) *The G-protein Linked Receptor Facts Book*, Academic Press, San Diego CA, pp 2-6). Mutations and changes in transcriptional activation of GPCR-encoding genes have been

associated with neurological disorders such as schizophrenia, Parkinson's disease, Alzheimer's disease, drug addiction, and feeding disorders.

Cytochromes are electron-transferring proteins that contain a heme prosthetic group, a porphyrin ring containing a tightly bound iron atom. Cytochromes act as oxidoreductases in such  
 5 diverse cellular processes as respiration, photosynthesis, fatty acid metabolism, and neurotransmitter biosynthesis. The heme iron atom serves as the actual electron carrier by changing from the ferric to the ferrous oxidation state when accepting an electron. Cytochromes accept electrons from one substrate such as NADH or ascorbate and donate them to other electron carriers such as other cytochromes, ubiquinone, or semidehydroascorbic acid (Lodish, H. et al. (1995) Molecular Cell  
 10 Biology, Scientific American Books, New York NY, pp. 759-770, 786-797; Sperling, P. et al. (1995) Eur. J. Biochem. 232:798-805; and Online Mendelian Inheritance in Man (OMIM) \*600019 Cytochrome b561, CYB561).

Cytochrome b5 is an electron donor in membrane-linked redox enzyme systems involved in lipid and drug metabolism. Cytochrome b5 has been found in Golgi, plasma, outer mitochondrial,  
 15 endoplasmic reticulum (ER), and microbody membranes. Conserved amino acids in cytochrome b5 include eight invariant amino acids at W34, H51, P52, G53, G54, G63, F70, and H74, and fifteen conserved amino acids at L24, I35, S36, V41, Y42, N43, T45, W47, A48, L58, D65, T67, L85, T87, and G88 (numbering based on the sunflower cytochrome b5/delta-6 desaturase fusion protein; GI 1040729, Sperling, supra). The invariant residues H51PGG are involved in heme-binding.  
 20 Cytochrome b5-like domains have also been found linked to other enzymes. For example, cytochrome b5-like domains are part of delta-9 fatty acid desaturases in yeast and Histoplasma capsulatum, nitrate reductase, sulfite reductase, flavocytochrome b2, Arabidopsis thaliana acyl lipid desaturase, and Borago officinalis (borage) and Helianthus annuus (sunflower) delta-6 desaturases (Sperling, supra; Sayanova, O. et al (1997) Proc. Natl. Acad. Sci. USA 94:4211-4216; and Mitchell,  
 25 A.G. and C.E. Martin (1997) J. Biol. Chem. 272:28281-28288).

Signal peptides are found on proteins that are targeted to the endoplasmic reticulum (ER). Signal peptides consist of stretches of amino acids enriched in hydrophobic residues. Signal peptides are usually found at the extreme N-terminus of the protein and are recognized by a cytosolic signal-recognition peptide (SRP). The SRP binds to the signal peptide and to an SRP receptor, an integral  
 30 membrane protein in the ER. Once bound to the SRP receptor, the newly formed protein containing the signal peptide is translocated across the ER membrane. Proteins containing signal peptides may end up inserted into the lipid bilayer, or they may end up in the lumen of an organelle or secreted from the cell.

### 35 Disease Correlation

The etiology of numerous human diseases and disorders can be attributed to defects in the

transport of molecules across membranes. Defects in the trafficking of membrane-bound transporters and ion channels are associated with several disorders, e.g. cystic fibrosis, glucose-galactose malabsorption syndrome, hypercholesterolemia, von Gierke disease, and certain forms of diabetes mellitus. Single-gene defect diseases resulting in an inability to transport small molecules across  
 5 membranes include, e.g., cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease (van't Hoff, W.G. (1996) *Exp. Nephrol.* 4:253-262; Talente, G.M. et al. (1994) *Ann. Intern. Med.* 120:218-226; and Chillon, M. et al. (1995) *New Engl. J. Med.* 332:1475-1480).

Cystinuria is an inherited disease that results from the inability to transport cystine, the disulfide-linked dimer of cysteine, from the urine into the blood. Accumulation of cystine in the  
 10 urine leads to the formation of cystine stones in the kidneys.

Transthyretin (TTR), present in human plasma, binds to and transports the thyroid hormone thyroxine. Mutations in TTR result in the conversion of TTR to amyloid, an insoluble fibrillar structure. The resulting amyloid plaques have been shown to be the causative agent in the development of familial amyloid polyneuropathy and senile systemic amyloidosis (Miroy, G.J. et al.  
 15 (1996) *Proc. Natl. Acad. Sci. USA* 93:15051-15056).

Stomatin, a 31-kDa erythrocyte integral membrane protein has been linked to the hereditary anemia stomatocytosis. This anemia is characterized by red blood cells that lack stomatin and leak Na<sup>+</sup> and K<sup>+</sup>. Thus, stomatin is presumed to play a role in the regulation of ion transport. Red blood cell ion transport defects are also linked to other disorders such as hypertension (Stewart, G.W.  
 20 (1997) *Int. J. Biochem. Cell Biol.* 29:271-274).

The discovery of new human transport proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of transport, metabolic, neurological, cardiovascular, reproductive, and  
 25 immune disorders, and cell proliferative disorders including cancer.

### SUMMARY OF THE INVENTION

The invention features purified polypeptides, human transport proteins, referred to collectively as "TPPT" and individually as "TPPT-1," "TPPT-2," "TPPT-3," "TPPT-4," "TPPT-5,"  
 30 "TPPT-6," "TPPT-7," "TPPT-8," "TPPT-9," "TPPT-10," "TPPT-11," "TPPT-12," "TPPT-13," "TPPT-14," "TPPT-15," "TPPT-16," "TPPT-17," "TPPT-18," "TPPT-19," "TPPT-20," "TPPT-21," "TPPT-22," "TPPT-23," "TPPT-24," "TPPT-25," "TPPT-26," "TPPT-27," "TPPT-28," "TPPT-29," "TPPT-30," "TPPT-31," "TPPT-32," "TPPT-33," "TPPT-34," "TPPT-35," "TPPT-36," "TPPT-37," "TPPT-38," "TPPT-39," "TPPT-40," "TPPT-41," "TPPT-42," and "TPPT-43." In one aspect, the  
 35 invention provides an isolated polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-

43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-43.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-43. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:44-86.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a



polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43.

The invention further provides an isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide comprising a polynucleotide sequence selected from the group consisting of a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:44-86, c) a polynucleotide sequence complementary to a), d) a polynucleotide sequence complementary to b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a pharmaceutical composition comprising an effective amount of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and a pharmaceutically acceptable excipient. In one embodiment, the pharmaceutical composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional TPPT, comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional TPPT, comprising administering to a patient in need of such treatment the pharmaceutical composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional TPPT, comprising administering to a patient in

need of such treatment the pharmaceutical composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide comprising an amino acid sequence selected from the group consisting of a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43, and d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-43. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:44-86, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

### BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding TPPT.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of TPPT.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases, disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

5 Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding TPPT were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

10

### DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing  
15 particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a  
20 reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be  
25 used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

### 30 DEFINITIONS

"TPPT" refers to the amino acid sequences of substantially purified TPPT obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of  
35 TPPT. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of TPPT either by directly interacting with

TPPT or by acting on components of the biological pathway in which TPPT participates.

An "allelic variant" is an alternative form of the gene encoding TPPT. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or  
5 many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding TPPT include those sequences with deletions,  
10 insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as TPPT or a polypeptide with at least one functional characteristic of TPPT. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding TPPT, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding  
15 TPPT. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent TPPT. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of TPPT is retained. For example,  
20 negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

25 The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to a sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

30 "Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of TPPT. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small  
35 molecules, or any other compound or composition which modulates the activity of TPPT either by directly interacting with TPPT or by acting on components of the biological pathway in which TPPT

participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind TPPT polypeptides can be prepared using intact polypeptides or using fragments  
5 containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

10 The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen  
15 used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified  
20 sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or  
25 translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic TPPT, or of any oligopeptide thereof,  
30 to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

35 A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or

amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding TPPT or fragments of TPPT may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be  
 5 deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (PE Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from  
 10 one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least  
 15 interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
20	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
25	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
30	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
35	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

40 Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

10 A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

A "fragment" is a unique portion of TPPT or the polynucleotide encoding TPPT which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

25 A fragment of SEQ ID NO:44-86 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:44-86, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:44-86 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:44-86 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:44-86 and the region of SEQ ID NO:44-86 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

30 A fragment of SEQ ID NO:1-43 is encoded by a fragment of SEQ ID NO:44-86. A fragment of SEQ ID NO:1-43 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-43. For example, a fragment of SEQ ID NO:1-43 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-43. The precise length of a fragment of SEQ ID NO:1-43 and the region of SEQ ID NO:1-43 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended



purpose for the fragment.

A "full-length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full-length" polynucleotide sequence encodes a "full-length" polypeptide sequence.

5 "Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps  
10 in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of  
15 molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent  
20 similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at  
25 <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The  
30 "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

35 *Reward for match: 1*

*Penalty for mismatch: -2*

*Open Gap: 5 and Extension Gap: 2 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 11*

5 *Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous  
10 nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes  
15 in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some  
20 alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e  
25 sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

30 Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (Apr-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

*Matrix: BLOSUM62*

35 *Open Gap: 11 and Extension Gap: 1 penalties*

*Gap x drop-off: 50*

*Expect: 10*

*Word Size: 3*

*Filter: on*

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating  $T_m$  and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C<sub>0</sub>t or R<sub>0</sub>t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of TPPT which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of TPPT which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of TPPT. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of TPPT.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide,

polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an TPPT may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of TPPT.

"Probe" refers to nucleic acid sequences encoding TPPT, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

"Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, 2<sup>nd</sup> ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs

can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence. This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is

expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription,  
5 translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent, chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

10 An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic  
15 acids encoding TPPT, or fragments thereof, or TPPT itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or  
20 synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

25 The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides  
30 by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

35 A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

“Transformation” describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term “transformed” cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

10 A “transgenic organism,” as used herein, is any organism, including but not limited to animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants, and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

25 A “variant” of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the “BLAST 2 Sequences” tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an “allelic” (as defined above), “splice,” “species,” or “polymorphic” variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass “single nucleotide



polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having  
 5 at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

## 10 THE INVENTION

The invention is based on the discovery of new human transport proteins (TPPT), the polynucleotides encoding TPPT, and the use of these compositions for the diagnosis, treatment, or prevention of transport, metabolic, neurological, cardiovascular, reproductive, and immune disorders, and cell proliferative disorders including cancer.

15 Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding TPPT. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each TPPT were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their corresponding cDNA  
 20 libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. In some cases, GenBank sequence identifiers are also shown in column 5. The Incyte clones and GenBank cDNA sequences, where indicated, in column 5 were used to assemble the consensus nucleotide sequence of each TPPT and are useful as fragments in hybridization technologies.

25 The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows homologous sequences as identified by BLAST analysis; and column 7 shows analytical  
 30 methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding TPPT. The first column of Table 3 lists the nucleotide  
 35 SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID NO:44-86

and to distinguish between SEQ ID NO:44-86 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express TPPT as a fraction of total tissues expressing TPPT. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing TPPT as a fraction of total tissues expressing TPPT. Column 5 lists the vectors used to subclone each cDNA library.

Of particular interest is the expression of SEQ ID NO:50 exclusively in cardiovascular tissue, the expression of SEQ ID NO:56 in nervous and gastrointestinal tissues, the expression of SEQ ID NO:57 in gastrointestinal tissues, and the expression of SEQ ID NO:66 in nervous system tissues. Of particular note is the tissue-specific expression of SEQ ID NO:75. Over 71% of the cDNA libraries expressing SEQ ID NO:75 are derived from lung tissue.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries from which cDNA clones encoding TPPT were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

SEQ ID NO:44 maps to chromosome 7 within the interval from 38.80 to 42.10 centiMorgans. SEQ ID NO:48 maps to chromosome X within the interval from 107.90 to 122.80 centiMorgans. SEQ ID NO:60 maps to chromosome 2 within the interval from 157.0 to 167.0 centiMorgans. SEQ ID NO:65 maps to chromosome 2 within the interval from 17.4 to 40.7 centiMorgans and to chromosome 5 within the interval from 61.1 to 69.6 centiMorgans. The interval on chromosome 5 from 61.1 to 69.6 centiMorgans also contains genes associated with Cockayne syndrome. SEQ ID NO:69 maps to chromosome 3 within the interval from 157.40 to 162.00 centiMorgans. SEQ ID NO:70 maps to chromosome 3 within the interval from 176.40 to 179.80 centiMorgans. SEQ ID NO:71 maps to chromosome 18 within the interval from the p-terminus to 52.30 centiMorgans. SEQ ID NO:73 maps to chromosome 17 within the interval from 75.70 to 84.20 centiMorgans, and to chromosome 2 within the interval from 204.70 to 209.30 centiMorgans. SEQ ID NO:76 maps to chromosome 20 within the interval from 79.00 to 94.40 centiMorgans. SEQ ID NO:80 maps to chromosome 18 within the interval from 1.60 to 6.20 centiMorgans, and to chromosome 11 within the interval from 117.90 to 126.00 centiMorgans. SEQ ID NO:83 maps to chromosome 17 within the interval from 67.60 to 69.30 centiMorgans, and from 83.8 centiMorgans to the q-terminus, and to chromosome 7 within the interval from 105.20 to 114.50 centiMorgans.

The invention also encompasses TPPT variants. A preferred TPPT variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the TPPT amino acid sequence, and which contains at least one functional or structural characteristic of TPPT.

The invention also encompasses polynucleotides which encode TPPT. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected

from the group consisting of SEQ ID NO:44-86, which encodes TPPT. The polynucleotide sequences of SEQ ID NO:44-86, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

5       The invention also encompasses a variant of a polynucleotide sequence encoding TPPT. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding TPPT. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:44-  
10 86 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:44-86. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of TPPT.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the  
15 genetic code, a multitude of polynucleotide sequences encoding TPPT, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the  
20 polynucleotide sequence of naturally occurring TPPT, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode TPPT and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring TPPT under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding TPPT or  
25 its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding TPPT and its derivatives without altering the encoded amino acid sequences  
30 include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode TPPT and TPPT derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems  
35 using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding TPPT or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:44-86 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (PE Biosystems, Foster City CA), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (PE Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (PE Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding TPPT may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060.) Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo

Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, PE Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode TPPT may be cloned in recombinant DNA molecules that direct expression of TPPT, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express TPPT.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter TPPT-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Cramer, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or

improve the biological properties of TPPT, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding TPPT may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.) Alternatively, TPPT itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) *Proteins, Structures and Molecular Properties*, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (PE Biosystems). Additionally, the amino acid sequence of TPPT, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, *supra*, pp. 28-53.)

In order to express a biologically active TPPT, the nucleotide sequences encoding TPPT or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding TPPT. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding TPPT. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding TPPT and its initiation codon and upstream regulatory sequences are inserted into

the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both  
 5 natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding TPPT and appropriate transcriptional and translational control  
 10 elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences  
 15 encoding TPPT. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or  
 20 animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984)  
 25 *Science* 224:838-843; Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences  
 30 to the targeted organ, tissue, or cell population. (See, e.g., Di Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al., (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.) The invention is not limited by the host cell employed.

35 In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding TPPT. For example, routine cloning,

subcloning, and propagation of polynucleotide sequences encoding TPPT can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding TPPT into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of TPPT are needed, e.g. for the production of antibodies, vectors which direct high level expression of TPPT may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of TPPT. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, *supra*; and Scorer, *supra*.)

Plant systems may also be used for expression of TPPT. Transcription of sequences encoding TPPT may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, *supra*; Broglie, *supra*; and Winter, *supra*.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding TPPT may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses TPPT in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet.



15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of TPPT in cell lines is preferred. For example, sequences encoding TPPT can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous  
5 expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue  
10 culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk<sup>-</sup>* and *apr<sup>-</sup>* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or  
15 herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which  
20 alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech),  $\beta$  glucuronidase and its substrate  $\beta$ -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system.  
25 (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding TPPT is inserted within a marker gene sequence, transformed cells containing sequences encoding TPPT can be identified by the absence of marker gene function. Alternatively, a  
30 marker gene can be placed in tandem with a sequence encoding TPPT under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding TPPT and that express TPPT may be identified by a variety of procedures known to those of skill in the art. These  
35 procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or

chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of TPPT using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and  
5 fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on TPPT is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and  
10 Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding TPPT  
15 include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding TPPT, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety  
20 of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding TPPT may be cultured under  
25 conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode TPPT may be designed to contain signal sequences which direct secretion of TPPT through a prokaryotic or eukaryotic cell membrane.

30 In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity.  
35 Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the

American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding TPPT may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric TPPT protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of TPPT activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the TPPT encoding sequence and the heterologous protein sequence, so that TPPT may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled TPPT may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, <sup>35</sup>S-methionine.

TPPT of the present invention or fragments thereof may be used to screen for compounds that specifically bind to TPPT. At least one and up to a plurality of test compounds may be screened for specific binding to TPPT. Examples of test compounds include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of TPPT, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which TPPT binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express TPPT, either as a secreted

protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing TPPT or cell membrane fractions which contain TPPT are then contacted with a test compound and binding, stimulation, or inhibition of activity of either TPPT or the compound is analyzed.

5 An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with TPPT, either in solution or affixed to a solid support, and detecting the binding of TPPT to the compound.

Alternatively, the assay may detect or measure binding of a test compound in the presence of a  
10 labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

TPPT of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of TPPT. Such compounds may include agonists, antagonists, or partial or  
15 inverse agonists. In one embodiment, an assay is performed under conditions permissive for TPPT activity, wherein TPPT is combined with at least one test compound, and the activity of TPPT in the presence of a test compound is compared with the activity of TPPT in the absence of the test compound. A change in the activity of TPPT in the presence of the test compound is indicative of a compound that modulates the activity of TPPT. Alternatively, a test compound is combined with an  
20 in vitro or cell-free system comprising TPPT under conditions suitable for TPPT activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of TPPT may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding TPPT or their mammalian homologs may  
25 be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of human disease. (See, e.g., U.S. Patent No. 5,175,383 and U.S. Patent No. 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of  
30 interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids  
35 Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred

to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding TPPT may also be manipulated *in vitro* in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding TPPT can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding TPPT is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress TPPT, e.g., by secreting TPPT in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

#### THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of TPPT and human transport proteins. In addition, the expression of TPPT is closely associated with neurological, cardiovascular, reproductive, gastrointestinal, and hematopoietic/immune tissues, and inflammation, cell proliferation, and cancer. Therefore, TPPT appears to play a role in transport, metabolic, neurological, cardiovascular, reproductive, and immune disorders, and cell proliferative disorders including cancer. In the treatment of disorders associated with increased TPPT expression or activity, it is desirable to decrease the expression or activity of TPPT. In the treatment of disorders associated with decreased TPPT expression or activity, it is desirable to increase the expression or activity of TPPT.

Therefore, in one embodiment, TPPT or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TPPT. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie Tooth disease, diabetes mellitus, diabetes insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis, normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, multidrug resistance, myasthenia gravis, myotonic dystrophy, catatonia, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer, cardiac disorders associated with transport, e.g., angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis, cardiomyopathy, nemaline

myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, thyrotoxic myopathy, ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis; neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychoses, and schizophrenia; and other disorders associated with transport, e.g., neurofibromatosis, postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease, cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Menkes disease, occipital horn syndrome, von Gierke disease, cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease; a metabolic disorder such as Addison's disease, cerebrotendinous xanthomatosis, congenital adrenal hyperplasia, coumarin resistance, cystic fibrosis, diabetes, fatty hepatocirrhosis, fructose-1,6-diphosphatase deficiency, galactosemia, goiter, glucagonoma, glycogen storage diseases, hereditary fructose intolerance, hyperadrenalism, hypoadrenalism, hyperparathyroidism, hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypoglycemia, hypothyroidism, hyperlipidemia, hyperlipemia, lipid myopathies, lipodystrophies, lysosomal storage diseases, mannosidosis, neuraminidase deficiency, obesity, pentosuria phenylketonuria, and pseudovitamin D-deficiency rickets; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a cardiovascular disorder such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins,

thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve,

5 mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation, congenital lung anomalies, atelectasis, pulmonary congestion and edema, pulmonary embolism, pulmonary

10 hemorrhage, pulmonary infarction, pulmonary hypertension, vascular sclerosis, obstructive pulmonary disease, restrictive pulmonary disease, chronic obstructive pulmonary disease, emphysema, chronic bronchitis, bronchial asthma, bronchiectasis, bacterial pneumonia, viral and mycoplasmal pneumonia, lung abscess, pulmonary tuberculosis, diffuse interstitial diseases, pneumoconioses, sarcoidosis, idiopathic pulmonary fibrosis, desquamative interstitial pneumonitis, hypersensitivity pneumonitis,

15 pulmonary eosinophilia bronchiolitis obliterans-organizing pneumonia, diffuse pulmonary hemorrhage syndromes, Goodpasture's syndromes, idiopathic pulmonary hemosiderosis, pulmonary involvement in collagen-vascular disorders, pulmonary alveolar proteinosis, lung tumors, inflammatory and noninflammatory pleural effusions, pneumothorax, pleural tumors, drug-induced lung disease, radiation-induced lung disease, and complications of lung transplantation; a

20 reproductive disorder such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm

25 physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis,

30 autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia,

35 irritable bowel syndrome, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis,

pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, 5 fungal, parasitic, protozoal, and helminthic infections, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, 10 myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

In another embodiment, a vector capable of expressing TPPT or a fragment or derivative 15 thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TPPT including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified TPPT in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TPPT including, but not 20 limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of TPPT may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of TPPT including, but not limited to, those listed above.

In a further embodiment, an antagonist of TPPT may be administered to a subject to treat or 25 prevent a disorder associated with increased expression or activity of TPPT. Examples of such disorders include, but are not limited to, those transport, metabolic, neurological, cardiovascular, reproductive, and immune disorders, and cell proliferative disorders including cancer described above. In one aspect, an antibody which specifically binds TPPT may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to 30 cells or tissues which express TPPT.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding TPPT may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of TPPT including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary 35 sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made



by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

5 An antagonist of TPPT may be produced using methods which are generally known in the art. In particular, purified TPPT may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind TPPT. Antibodies to TPPT may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and  
10 fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with TPPT or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to  
15 increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to TPPT  
20 have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of TPPT amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

25 Monoclonal antibodies to TPPT may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and  
30 Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda,  
35 S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce TPPT-specific single

chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte  
5 population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for TPPT may also be generated. For example, such fragments include, but are not limited to,  $F(ab')_2$  fragments produced by pepsin  
10 digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the  $F(ab)_2$  fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired  
15 specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between TPPT and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering TPPT epitopes is generally used, but a competitive binding assay may  
20 also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for TPPT. Affinity is expressed as an association constant,  $K_a$ , which is defined as the molar concentration of TPPT-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions.  
25 The  $K_a$  determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple TPPT epitopes, represents the average affinity, or avidity, of the antibodies for TPPT. The  $K_a$  determined for a preparation of monoclonal antibodies, which are monospecific for a particular TPPT epitope, represents a true measure of affinity. High-affinity antibody preparations with  $K_a$  ranging from about  $10^9$  to  $10^{12}$  L/mole are preferred for use in immunoassays in which the  
30 TPPT-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with  $K_a$  ranging from about  $10^6$  to  $10^7$  L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of TPPT, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons,  
35 New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to

determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of TPPT-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines  
 5 for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al., supra.)

In another embodiment of the invention, the polynucleotides encoding TPPT, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA,  
 10 RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding TPPT. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding TPPT. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense  
 15 sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral  
 20 vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.*  
 25 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding TPPT may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined  
 30 immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal,  
 35 R.G. (1995) *Science* 270:404-410; Verma, I.M. and Somia, N. (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated

cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) Nature 335:395-396; Poeschla, E. et al. (1996) Proc. Natl. Acad. Sci. USA. 93:11395-11399), hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in TPPT expression or regulation causes disease, the expression of TPPT from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in TPPT are treated by constructing mammalian expression vectors encoding TPPT and introducing these vectors by mechanical means into TPPT-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) Annu. Rev. Biochem. 62:191-217; Ivics, Z. (1997) Cell 91:501-510; Boulay, J-L. and H. Récipon (1998) Curr. Opin. Biotechnol. 9:445-450).

Expression vectors that may be effective for the expression of TPPT include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). TPPT may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or  $\beta$ -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) Proc. Natl. Acad. Sci. USA 89:5547-5551; Gossen, M. et al. (1995) Science 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) Curr. Opin. Biotechnol. 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and H.M. Blau, supra), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding TPPT from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) Virology 52:456-467), or by electroporation (Neumann, E. et al. (1982) EMBO J. 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with respect to TPPT expression are treated by constructing a retrovirus vector consisting of (i) the p lynucle tide encoding TPPT under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) Proc. Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al. (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference. Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4<sup>+</sup> T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).

In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding TPPT to cells which have one or more genetic abnormalities with respect to the expression of TPPT. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544; and Verma, I.M. and N. Somia (1997) Nature 18:389:239-242, both incorporated by reference herein.

In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding TPPT to target cells which have one or more genetic abnormalities with respect to the expression of TPPT. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing TPPT to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has

been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) *Exp. Eye Res.* 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) *J. Virol.* 73:519-532 and Xu, H. et al. (1994) *Dev. Biol.* 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding TPPT to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) *Curr. Opin. Biotech.* 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full-length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for TPPT into the alphavirus genome in place of the capsid-coding region results in the production of a large number of TPPT-coding RNAs and the synthesis of high levels of TPPT in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) *Virology* 228:74-83). The wide host range of alphaviruses will allow the introduction of TPPT into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have

been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

5 Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding TPPT.

10 Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of  
15 candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis.

20 Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding TPPT. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

25 RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2'-O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine,  
30 and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding TPPT. Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not  
35 limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular

chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased TPPT expression or activity, a compound which specifically inhibits expression of the polynucleotide  
5 encoding TPPT may be therapeutically useful, and in the treatment of disorders associated with decreased TPPT expression or activity, a compound which specifically promotes expression of the polynucleotide encoding TPPT may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method  
10 commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a  
15 polynucleotide encoding TPPT is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an in vitro cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding TPPT are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence of the  
20 polynucleotide encoding TPPT. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific polynucleotide  
25 can be carried out, for example, using a Schizosaccharomyces pombe gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide  
30 nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide sequence (Bruce, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruce, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells  
35 taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved



using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and  
5 monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest  
10 edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such pharmaceutical compositions may consist of TPPT, antibodies to TPPT, and mimetics, agonists, antagonists, or inhibitors of TPPT.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial,  
15 intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Pharmaceutical compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol  
20 delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration  
25 enhancers.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of pharmaceutical compositions may be prepared for direct intracellular  
30 delivery of macromolecules comprising TPPT or fragments thereof. For example, liposome preparations containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, TPPT or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system  
35 (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell

culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

5 A therapeutically effective dose refers to that amount of active ingredient, for example TPPT or fragments thereof, antibodies of TPPT, and agonists, antagonists or inhibitors of TPPT, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose  
10 lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the  $LD_{50}/ED_{50}$  ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with  
15 little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the  
20 severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

25 Normal dosage amounts may vary from about 0.1  $\mu\text{g}$  to 100,000  $\mu\text{g}$ , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells,  
30 conditions, locations, etc.

## DIAGNOSTICS

In another embodiment, antibodies which specifically bind TPPT may be used for the diagnosis of disorders characterized by expression of TPPT, or in assays to monitor patients being treated with TPPT or agonists, antagonists, or inhibitors of TPPT. Antibodies useful for diagnostic  
35 purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for TPPT include methods which utilize the antibody and a label to detect TPPT in human body fluids

r in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

5 A variety of protocols for measuring TPPT, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of TPPT expression. Normal or standard values for TPPT expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibody to TPPT under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of TPPT expressed in subject, 10 control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding TPPT may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and 15 quantify gene expression in biopsied tissues in which expression of TPPT may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of TPPT, and to monitor regulation of TPPT levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding TPPT or closely related molecules may be used to 20 identify nucleic acid sequences which encode TPPT. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding TPPT, allelic variants, or related sequences.

25 Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the TPPT encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:44-86 or from genomic sequences including promoters, enhancers, and introns of the TPPT gene.

Means for producing specific hybridization probes for DNAs encoding TPPT include the 30 cloning of polynucleotide sequences encoding TPPT or TPPT derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as  $^{32}\text{P}$  or  $^{35}\text{S}$ , or by enzymatic labels, 35 such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding TPPT may be used for the diagnosis of disorders

associated with expression of TPPT. Examples of such disorders include, but are not limited to, a transport disorder such as akinesia, amyotrophic lateral sclerosis, ataxia telangiectasia, cystic fibrosis, Becker's muscular dystrophy, Bell's palsy, Charcot-Marie Tooth disease, diabetes mellitus, diabetes insipidus, diabetic neuropathy, Duchenne muscular dystrophy, hyperkalemic periodic paralysis, normokalemic periodic paralysis, Parkinson's disease, malignant hyperthermia, multidrug resistance, myasthenia gravis, myotonic dystrophy, catatonia, tardive dyskinesia, dystonias, peripheral neuropathy, cerebral neoplasms, prostate cancer; cardiac disorders associated with transport, e.g., angina, bradyarrhythmia, tachyarrhythmia, hypertension, Long QT syndrome, myocarditis, cardiomyopathy, nemaline myopathy, centronuclear myopathy, lipid myopathy, mitochondrial myopathy, thyrotoxic myopathy, ethanol myopathy, dermatomyositis, inclusion body myositis, infectious myositis, polymyositis; neurological disorders associated with transport, e.g., Alzheimer's disease, amnesia, bipolar disorder, dementia, depression, epilepsy, Tourette's disorder, paranoid psychoses, and schizophrenia; and other disorders associated with transport, e.g., neurofibromatosis, postherpetic neuralgia, trigeminal neuropathy, sarcoidosis, sickle cell anemia, Wilson's disease, cataracts, infertility, pulmonary artery stenosis, sensorineural autosomal deafness, hyperglycemia, hypoglycemia, Grave's disease, goiter, Cushing's disease, Addison's disease, glucose-galactose malabsorption syndrome, hypercholesterolemia, adrenoleukodystrophy, Zellweger syndrome, Menkes disease, occipital horn syndrome, von Gierke disease, cystinuria, iminoglycinuria, Hartup disease, and Fanconi disease; a metabolic disorder such as Addison's disease, cerebrotendinous xanthomatosis, congenital adrenal hyperplasia, coumarin resistance, cystic fibrosis, diabetes, fatty hepatocirrhosis, fructose-1,6-diphosphatase deficiency, galactosemia, goiter, glucagonoma, glycogen storage diseases, hereditary fructose intolerance, hyperadrenalism, hypoadrenalism, hyperparathyroidism, hypoparathyroidism, hypercholesterolemia, hyperthyroidism, hypoglycemia, hypothyroidism, hyperlipidemia, hyperlipemia, lipid myopathies, lipodystrophies, lysosomal storage diseases, mannosidosis, neuraminidase deficiency, obesity, pentosuria phenylketonuria, and pseudovitamin D-deficiency rickets; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial

nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; a cardiovascular disorder such as arteriovenous fistula, atherosclerosis, hypertension, vasculitis, Raynaud's disease, aneurysms, arterial dissections, varicose veins, thrombophlebitis and phlebothrombosis, vascular tumors, and complications of thrombolysis, balloon angioplasty, vascular replacement, and coronary artery bypass graft surgery, congestive heart failure, ischemic heart disease, angina pectoris, myocardial infarction, hypertensive heart disease, degenerative valvular heart disease, calcific aortic valve stenosis, congenitally bicuspid aortic valve, mitral annular calcification, mitral valve prolapse, rheumatic fever and rheumatic heart disease, infective endocarditis, nonbacterial thrombotic endocarditis, endocarditis of systemic lupus erythematosus, carcinoid heart disease, cardiomyopathy, myocarditis, pericarditis, neoplastic heart disease, congenital heart disease, and complications of cardiac transplantation, congenital lung anomalies, atelectasis, pulmonary congestion and edema, pulmonary embolism, pulmonary hemorrhage, pulmonary infarction, pulmonary hypertension, vascular sclerosis, obstructive pulmonary disease, restrictive pulmonary disease, chronic obstructive pulmonary disease, emphysema, chronic bronchitis, bronchial asthma, bronchiectasis, bacterial pneumonia, viral and mycoplasmal pneumonia, lung abscess, pulmonary tuberculosis, diffuse interstitial diseases, pneumoconioses, sarcoidosis, idiopathic pulmonary fibrosis, desquamative interstitial pneumonitis, hypersensitivity pneumonitis, pulmonary eosinophilia bronchiolitis obliterans-organizing pneumonia, diffuse pulmonary hemorrhage syndromes, Goodpasture's syndromes, idiopathic pulmonary hemosiderosis, pulmonary involvement in collagen-vascular disorders, pulmonary alveolar proteinosis, lung tumors, inflammatory and noninflammatory pleural effusions, pneumothorax, pleural tumors, drug-induced lung disease, radiation-induced lung disease, and complications of lung transplantation; a reproductive disorder such as a disorder of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, a disruption of the estrous cycle, a disruption of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, an endometrial or ovarian tumor, a uterine fibroid, autoimmune disorders, an ectopic pregnancy, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; a disruption of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome,

allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, 5 episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, 10 polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, trauma, and hematopoietic cancer including 15 lymphoma, leukemia, and myeloma; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, 20 bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding TPPT may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from 25 patients to detect altered TPPT expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding TPPT may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding TPPT may be labeled by standard methods and added to a fluid or tissue sample 30 from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding TPPT in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate 35 the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of TPPT, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding TPPT, under conditions suitable for hybridization or amplification.

- 5 Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

- 10 Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

- 15 With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

- 20 Additional diagnostic uses for oligonucleotides designed from the sequences encoding TPPT may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced *in vitro*. Oligomers will preferably contain a fragment of a polynucleotide encoding TPPT, or a fragment of a polynucleotide complementary to the polynucleotide encoding TPPT, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

- 25 In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding TPPT may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding TPPT are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are

detectable using gel electrophoresis in non-denaturing gels. In fSCCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of TPPT include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described in Seilhamer, J.J. et al., "Comparative Gene Transcript Analysis," U.S. Patent No. 5,840,484, incorporated herein by reference. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, antibodies specific for TPPT, or TPPT or fragments thereof may be used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-



2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding TPPT may be used  
5 to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a  
10 chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop  
15 genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, e.g., Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic  
20 map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding TPPT on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as  
25 linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely  
30 localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

35 In another embodiment of the invention, TPPT, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug

screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between TPPT and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with TPPT, or fragments thereof, and washed. Bound TPPT is then detected by methods well known in the art. Purified TPPT can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding TPPT specifically compete with a test compound for binding TPPT. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with TPPT.

In additional embodiments, the nucleotide sequences which encode TPPT may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications and publications, mentioned above and below, in particular U.S. Ser. No. 60/139,923, U.S. Ser. No. 60/148,177, U.S. Ser. No. 60/149,357, and U.S. Ser. No. 60/162,287, are hereby expressly incorporated by reference.

## EXAMPLES

### I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A+) RNA was isolated

using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

5 In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic  
10 oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g.,  
15 PBLUEScript plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), pcDNA2.1 plasmid (Invitrogen, Carlsbad CA), or pINCY plasmid (Incyte Genomics, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 $\alpha$ , DH10B, or ElectroMAX DH10B from Life Technologies.

## II. Isolation of cDNA Clones

20 Plasmids obtained as described in Example I were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96  
25 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in  
30 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

## III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows.  
35 Sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (PE Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ

Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (PE Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VI.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such

as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) *Curr. Opin. Struct. Biol.* 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:44-86. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

#### IV. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, *supra*, ch. 7; Ausubel, 1995, *supra*, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum \{length(Seq. 1), length(Seq. 2)\}}}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and 70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding TPPT occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous,

reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in

5 Table 3.

#### V. Chromosomal Mapping of TPPT Encoding Polynucleotides

The cDNA sequences which were used to assemble SEQ ID NO:44-49 and SEQ ID NO:54-86 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these  
10 databases that matched SEQ ID NO:44-49 and SEQ ID NO:54-86 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 5). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a  
15 mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

The genetic map locations of SEQ ID NO:44, SEQ ID NO:48, SEQ ID NO:60, SEQ ID NO:65, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:73, SEQ ID NO:76, SEQ ID NO:80, and SEQ ID NO:83 are described in The Invention as ranges, or intervals, of human  
20 chromosomes. More than one map location is reported for SEQ ID NO:65, SEQ ID NO:73, SEQ ID NO:80, and SEQ ID NO:83, indicating that previously mapped sequences having similarity, but not complete identity, to SEQ ID NO:65, SEQ ID NO:73, SEQ ID NO:80, and SEQ ID NO:83 were assembled into their respective clusters. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-arm. (The centiMorgan (cM) is a unit of  
25 measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation hybrid markers whose sequences were included in each of the clusters. Diseases associated with the public and Incyte sequences located within the  
30 indicated intervals are also reported in the Invention section where applicable. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

#### VI. Extension of TPPT Encoding Polynucleotides

35 The full length nucleic acid sequences of SEQ ID NO:44-86 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this

fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing  $Mg^{2+}$ ,  $(NH_4)_2SO_4$ , and  $\beta$ -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100  $\mu$ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5  $\mu$ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was  
5 quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (PE Biosystems).

10 In like manner, the polynucleotide sequences of SEQ ID NO:44-86 are used to obtain 5' regulatory sequences using the procedure above, along with oligonucleotides designed for such extension, and an appropriate genomic library.

#### VII. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:44-86 are employed to screen cDNAs,  
15 genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ -<sup>32</sup>P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase  
20 (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10<sup>7</sup> counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

25 The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and  
30 compared.

#### VIII. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, supra),  
mechanical microspotting technologies, and derivatives thereof. The substrate in each of the  
35 aforementioned technologies should be uniform and solid with a non-porous surface (Schna (1999), supra). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers.



Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g.,

5 Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645; Marshall, A. and J. Hodgson (1998) *Nat. Biotechnol.* 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The

10 array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection. After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of

15 complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

#### Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and

20 poly(A)<sup>+</sup> RNA is purified using the oligo-(dT) cellulose method. Each poly(A)<sup>+</sup> RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/μl oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/μl RNase inhibitor, 500 μM dATP, 500 μM dGTP, 500 μM dTTP, 40 μM dCTP, 40 μM dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)<sup>+</sup> RNA with

25 GEMBRIGHT kits (Incyte). Specific control poly(A)<sup>+</sup> RNAs are synthesized by *in vitro* transcription from non-coding yeast genomic DNA. After incubation at 37 °C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85 °C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc.

30 (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μl 5X SSC/0.2% SDS.

#### Microarray Preparation

35 Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification

uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 µg. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

5 Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C  
10 oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 µl of the array element DNA, at an average concentration of 100 ng/µl, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

15 Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60 °C followed by washes in 0.2% SDS and distilled water as before.

#### 20 Hybridization

Hybridization reactions contain 9 µl of sample mixture consisting of 0.2 µg each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65 °C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm<sup>2</sup> coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly  
25 larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 µl of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60 °C. The arrays are washed for 10 min at 45 °C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45 °C in a second wash buffer (0.1X SSC), and dried.

#### Detection

30 Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-  
35 scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate filters positioned between the array and the photomultiplier tubes are used to filter the signals. The  
5 emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

The sensitivity of the scans is typically calibrated using the signal intensity generated by a cDNA control species added to the sample mixture at a known concentration. A specific location on  
10 the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two  
15 fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high  
20 signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated  
25 to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

#### **IX. Complementary Polynucleotides**

Sequences complementary to the TPPT-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring TPPT. Although use of oligonucleotides  
30 comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of TPPT. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is  
35 designed to prevent ribosomal binding to the TPPT-encoding transcript.

#### **X. Expression of TPPT**

Expression and purification of TPPT is achieved using bacterial or virus-based expression systems. For expression of TPPT in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express TPPT upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of TPPT in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding TPPT by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, TPPT is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from TPPT at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified TPPT obtained by these methods can be used directly in the assays shown in Examples XI and XV.

#### 30 XI. Demonstration of TPPT Activity

TPPT transport activity is assayed by measuring uptake of labeled substrates into Xenopus laevis oocytes. Oocytes at stages V and VI are injected with TPPT mRNA (10 ng per oocyte) and incubated for 3 days at 18°C in OR2 medium (82.5mM NaCl, 2.5 mM KCl, 1mM CaCl<sub>2</sub>, 1mM MgCl<sub>2</sub>, 1mM Na<sub>2</sub>HPO<sub>4</sub>, 5 mM Hepes, 3.8 mM NaOH, 50µg/ml gentamycin, pH 7.8) to allow expression of TPPT. Oocytes are then transferred to standard uptake medium (100mM NaCl, 2 mM KCl, 1mM CaCl<sub>2</sub>, 1mM MgCl<sub>2</sub>, 10 mM Hepes/Tris pH 7.5). Uptake of various substrates (e.g., amino acids,

sugars, drugs, ions, and neurotransmitters) is initiated by adding labeled substrate (e.g. radiolabeled with  $^3\text{H}$ , fluorescently labeled with rhodamine, etc.) to the oocytes. After incubating for 30 minutes, uptake is terminated by washing the oocytes three times in  $\text{Na}^+$ -free medium, measuring the incorporated label, and comparing with controls. TPPT activity is proportional to the level of internalized labeled substrate.

## XII. Functional Assays

TPPT function is assessed by expressing the sequences encoding TPPT at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen), both of which contain the cytomegalovirus promoter. 5-10  $\mu\text{g}$  of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2  $\mu\text{g}$  of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of TPPT on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding TPPT and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding TPPT and other genes of interest can be analyzed by northern analysis or microarray techniques.

### **XIII. Production of TPPT Specific Antibodies**

TPPT substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

5 Alternatively, the TPPT amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, *supra*, ch. 11.)

10 Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (PE Biosystems) using FMOC chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, *supra*.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-TPPT activity by, for example, binding the peptide or TPPT to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

### **XIV. Purification of Naturally Occurring TPPT Using Specific Antibodies**

Naturally occurring or recombinant TPPT is substantially purified by immunoaffinity chromatography using antibodies specific for TPPT. An immunoaffinity column is constructed by covalently coupling anti-TPPT antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing TPPT are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of TPPT (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/TPPT binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and TPPT is collected.

### **XV. Identification of Molecules Which Interact with TPPT**

TPPT, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled TPPT, washed, and any wells with labeled TPPT complex are assayed. Data obtained using different concentrations of TPPT are used to calculate values for the number, affinity, and association of TPPT with the candidate molecules.

35 Alternatively, molecules interacting with TPPT are analyzed using the yeast two-hybrid system as described in Fields, S. and O. Song (1989, *Nature* 340:245-246), or using commercially

available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

TPPT may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent  
5 No. 6,057,101).

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with certain embodiments, it should be  
10 understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	44	264114	HNT2AGT01	028972R6 (SPLNFET01), 028972T6 (SPLNFET01), 264114H1 (HNT2AGT01), 452387R6 (TLYMNOT02), 736580R1 (TONSNOT01), 747955R6 (BRAITUT01), 936731R1 (CERVNOT01), 3206282H1 (PENCNOT03), 3344943H1 (SPLNNOT09), 3742964H1 (THYMNOT08), 4028320H1 (BRAINOT23), 4726757H1 (GBLADIT01), 5473562H1 (MCLRUNT01)
2	45	1455669	COLNFET02	1455669H1 (COLNFET02), 2877376F6 (THYRNOT10), 3536452F6 (KIDNNOT25)
3	46	2084989	PANCNOT04	1281527H1 (COLNNOT16), 1412985H1 (BRAINOT12), 2084989H1 (PANCNOT04), 2084989R6 (PANCNOT04), 2084989T6 (PANCNOT04), 2470481F6 (THP1NOT03), 2539015F7 (BONRTUT01), 3109754F6 (BRSTTUT15), 3694831H1 (PANCNOT19), 3700647H1 (SININOT05)
4	47	2501034	ADRETUT05	111466F1 (PITUNOT01), 111466R1 (PITUNOT01), 414042R6 (BRSTNOT01), 687891H1 (UTRSNOT02), 2501034H1 (ADRETUT05)
5	48	2745212	LUNGTUT11	000802H1 (U937NOT01), 008963H1 (HMC1NOT01), 009314H1 (HMC1NOT01), 135428F1 (BARNOT02), 723168X19 (SYNOOAT01), 1000842R1 (BRSTNOT03), 1370189H1 (BSTMNON02), 13743329H1 (BSTMNON02), 2745212H1 (LUNGTUT11), 4920466H1 (TESTNOT11), SAIA02182F1
6	49	4833111	BRAVXT03	864776T1 (BRAITUT03), 1911267F6 (CONNTUT01), 4833111H1 (BRAVXT03), SARA02608F1, SARA02002F1
7	50	876677	LUNGAST01	876677H1 (LUNGAST01), 876677R6 (LUNGAST01), SCDA08642V1
8	51	2326143	OVARNOT02	867305R1 (BRAITUT03), 963058R2 (BRSTTUT03), 1715155F6 (UCMCNOT02), 1727927T6 (PROSNOT14), 2326143H1 (OVARNOT02), 2326143R6 (OVARNOT02), 3360563H1 (PROSTUT16)
9	52	2786302	BRSTNOT13	2786302H1 (BRSTNOT13), 2958321X303D1 (ADRENOT09), 2958321X305D1 (ADRENOT09), 2958321X308D1 (ADRENOT09)
10	53	3735780	SMCCNOS01	551126H1 (BEPINOT01), 2808373H1 (BLADTUT08), 3735780F6 (SMCCNOS01), 3735780H1 (SMCCNOS01), 3735780T6 (SMCCNOS01), 4760604T6 (BRAMNOT01)
11	54	039026	HUVENOB01	039026H1 (HUVENOB01), 159164F1 (ADENINB01), 159164R1 (ADENINB01)
12	55	260607	HNT2RAT01	063159R6 (PLACNOB01), 260607R6 (HNT2RAT01), 1272850T1 (TESTTUT02), 1273069H1 (TESTTUT02), 2867453F6 (KIDNNOT20), 3082466H1 (BRAIUNT01), 4796739H1 (LIVRTUT09), 4799318F6 (MYEPUNT01), 91424405



Table 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragmente
13	56	1429651	SINTBST01	1429651F1 (SINTBST01), 1429651H1 (SINTBST01), 1501096F6 (SINTBST01), 1989621T6 (CORPNOT02), SXLA01343V1, SXLA01183V1, SXLA01559V1, SXLA00812V1
14	57	2069971	ISLTNOT01	2069606F6 (ISLTNOT01), 2069971H1 (ISLTNOT01), 2374634F6 (ISLTNOT01), 2383754F6 (ISLTNOT01), 4171186T6 (SINTNOT21), SXLA01128V1, SXLA01348V1, SXLA01219V1, SXLA00260V1, SXLA00074V1
15	58	2329339	COLNNOT11	658662H1 (BRAINT03), 1544110R1 (PROSTUT04), 1657742F6 (URETTUT01), 1750523F6 (STOMTUT02), 2329339H1 (COLNNOT11), 2329339R6 (COLNNOT11), 3858671H1 (LNODNOT03), g1494061, g1891451
16	59	2540219	BONRTUT01	2540219H1 (BONRTUT01), 2540219T6 (BONRTUT01), 2554869F6 (THYMNOT03), g869197
17	60	2722462	LUNGUT10	883601R1 (PANCNOT05), 1525902F6 (UCMCL5T01), 1525902X18C1 (UCMCL5T01), 1525902X311D1 (UCMCL5T01), 1527325T6 (UCMCL5T01), 1554770X311D1 (BLADTUT04), 2417265H1 (HNT3AZT01), 2444786F6 (THPLNOT03), 2722462H1 (LUNGUT10), 4293114H1 (BRABDIR01), 5070268T6 (PANCNOT23), SANA01850F1, SAJA01078R1, SANA02081F1, SAJA01813F1
18	61	2739264	OVARNOT09	000573H1 (U937NOT01), 494409F1 (HNT2NOT01), 494409R1 (HNT2NOT01), 2506506F6 (CONUTUT01), 2681059H1 (SINIUCT01), 2744648F6 (BRSTTUT14), 2805590F6 (BLADTUT08), 3770643H1 (BRSTNOT25), 4204278H1 (BRAITUT29), SAEA02093F1
19	62	2758310	THP1AZS08	487309R7 (HNT2AGT01), 1361439F1 (LUNGNOT12), 2758310H1 (THP1AZS08), SCFA05584V1, SCFA05940V1, SCFA05166V1, SCFA05135V1
20	63	2762348	BRSTNOT12	632097R6 (KIDNNOT05), 632097T6 (KIDNNOT05), 2762348H1 (BRSTNOT12), SCCA02837V1, SCCA05356V1, SCCA01377V1, SCCA05963V1, SCCA05364V1, SCCA02307V1, SCCA04327V1, SCCA02009V1
21	64	3715961	PENCNOT09	961523H1 (BRSTTUT03), 1863723F6 (PROSNOT19), 2265329H1 (UTRSNOT02), 2360619R6 (LUNGFET05), 2360619T6 (LUNGFET05), 2821718H1 (ADRETUT06), 3715961H1 (PENCNOT09), 5016160H1 (BRAXNOT03), 5499583H1 (BRABDIR01)
22	65	5108194	PROSTUS19	1322651X35 (BLADNOT04), 1322651X36 (BLADNOT04), 3494841H1 (ADRETUT07), 4958978F6 (TLYMNOT05), 5108194H1 (PROSTUS19), g1379009, g1527417
23	66	5503122	BRABDIR01	5503122F6 (BRABDIR01), 5503122H1 (BRABDIR01), 5503122R6 (BRABDIR01)

Table 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
24	67	5517972	LIVDIR01	805957R1 (BSTNOT01), 953622R1 (SCORNON01), 1501080F1 (SINTBST01), 1547381R6 (PROSTUT04), 2081843T6 (UTRSNOT08), 2811524F6 (OVARNOT10), 3212921H1 (BLADNOT08), 3250443H1 (SEMVNOT03), 3269479H1 (BRAINOT20), 3699955F6 (SININOT05), 3700568H1 (SININOT05), 4944050H1 (BRAIFEN05), 5517972H1 (LIVDIR01)
25	68	5593114	COLCDIT03	2859465F6 (SININOT03), 2859465T6 (SININOT03), 3555656F6 (LUNGNOT31), 3555656T6 (LUNGNOT31), 4345952H1 (TLYMTXT01), 5593114H1 (COLCDIT03), 5874544H1 (COLTDIT04)
26	69	044775	TBLYN01	044775H1 (TBLYN01), 044775X3 (TBLYN01), 455640R1 (KERANOT01), 950702R1 (PANCNOT05), 2418550H1 (HNT3AZT01), 2798917H1 (NPOLNOT01), 2844686H1 (DRGLNOT01), 91718929
27	70	116588	KIDNNOT01	699714R6 (SYNORAT03), 831423R1 (PROSTUT04), 978875R1 (BRSTNOT02), 1350569F1 (LATRTUT02), 1447681R1 (PLACNOT02), 3177382F6 (UTRSTUT04), 3688796H1 (HEAANOT01), 3929008H1 (KIDNNOT19), 92106455, 92163092
28	71	875369	LUNGAST01	571573F1 (OVARNON01), 571573R1 (OVARNON01), 875369H1 (LUNGAST01), 875369R1 (LUNGAST01), 3569021H1 (HEAPNOT01)
29	72	1325518	LPARNOT02	1325518H1 (LPARNOT02), 1325518T6 (LPARNOT02), 1825553F6 (LSUBNOT03), SBAA02035F1
30	73	2060987	OVARNOT03	1378947T1 (LUNGNOT10), 1453290F1 (PENITUT01), 1459818R1 (COLNFET02), 1967477H1 (BRSTNOT04), 2060987H1 (OVARNOT03), 2455371F6 (ENDANOT01), 2499967F7 (ADRETUT05), 3093056T6 (BRSTNOT19), 3213366H1 (BLADNOT08), 4934158H1 (BRSTTUT20), SBYA01942U1
31	74	2172064	ENDCNOT03	2172064CT1 (ENDCNOT03), 2172064H1 (ENDCNOT03), SBLA01269F1
32	75	2219267	LUNGNOT18	2219267F6 (LUNGNOT18), 2219267H1 (LUNGNOT18), 3117478T6 (LUNGNOT13), 3126288T6 (LUNGNOT12), 3558495H1 (LUNGNOT31)
33	76	2308629	NGANNOT01	469862F1 (MMLRIDT01), 469862R1 (MMLRIDT01), 1594203X11C1 (BRAINOT14), 2191933H1 (THYRTUT03)
34	77	2660038	LUNGUT09	1326594F1 (LPARNOT02), 2256143H1 (OVARUT01), 2278689R6 (PROSNON01), 2528425H1 (GBLANOT02), 2660038H1 (LUNGUT09), 2660038T6 (LUNGUT09), 3449964H1 (UTRSNON03), 5099879H1 (PROSTUS20), 91886680, 9783969

Table 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
35	78	2670745	ESOGTUT02	259200X12 (HNT2RAT01), 1266477F1 (BRAINT09), 2383364F6 (ISLTNOT01), 2670745H1 (ESOGTUT02), 3181526H1 (TLYJNOT01)
36	79	2676443	KIDNNOT19	607375R6 (BRSTTUT01), 1728626X15C1 (PROSNOT14), 1751773F6 (LIVRTUT01), 1751994T6 (LIVRTUT01), 1796032X14C1 (PROSTUT05), 2010172H1 (TESTNOT03), 2676443H1 (KIDNNOT19)
37	80	3295764	TLYJINT01	063264H1 (PLACNOB01), 434468T6 (THYRNOT01), 487721H1 (HNT2AGT01), 907796R2 (COLNNOT09), 1212556R7 (BRSTTUT01), 1251889H1 (LUNGFET03), 1653370F6 (PROSTUT08), 1653370X309D1 (PROSTUT08), 2192762F6 (THYRTUT03), 2226786F6 (SEMVN01), 3295764H1 (TLYJINT01), 3384471H1 (ESOGNOT04), SASA01137F1
38	81	3438320	PENCNOT06	3438320H1 (PENCNOT06), 3501438H1 (PROSTUT13), 3745542H1 (THYMN08), 3751060H1 (UTRSNOT18), 4979750F6 (HELATXT04), SADA00043F1, SADA00087F1
39	82	3986488	UTRSTUT05	1634141F6 (COLNNOT19), 1692115X12C1 (PROSTUT10), 1731310F6 (BRSTTUT08), 2046232H1 (THP1T7T01), 3557951H1 (LUNGNOT31), 4726788H1 (GBLADIT01)
40	83	4378816	LUNGNOT37	1318962H1 (BLADNOT04), 1520864F1 (BLADTUT04), 1684381F6 (PROSNOT15), 2055747R6 (BEPINOT01), 4378816H1 (LUNGNOT37)
41	84	4797137	LIVRTUT09	4797137F6 (LIVRTUT09), 4797137H1 (LIVRTUT09), 4797137T6 (LIVRTUT09)
42	85	5470806	MCLRUNT01	5470806H1 (MCLRUNT01), 5470806T6 (MCLRUNT01)
43	86	5473242	MCLRUNT01	5473242F6 (MCLRUNT01), 5473242T6 (MCLRUNT01)

Table 2

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
1	623	S521 S2 T3 S16 S99 S138 S144 T193 T264 T404 S448 S589 S151 T229 T337 T457 S562 S568	N97 N333	BTB domain: C44-F56 POZ domain: N10-Q211 Kelch repeat signature: E379-G392, T398-V412, L438-M452, T498-A512 Ring canal protein repeat: E122-P254	Ring canal protein [Drosophila melanogaster] g577276	MOTIFS BLIMPS-PFAM BLIMPS-PRINTS BLAST-GenBank BLAST-PRODOM BLAST-DOMO
2	99	T17	N15	Signal peptide: M1-G36 Transmembrane region: S25-W45 MRP(2) MRP(1) repeat: C30-V74	Multi-drug resistance- associated protein (MRP)-like protein- 1 (MLP-1) [Rattus norvegicus] g3242458	MOTIFS BLAST-GenBank BLAST-PRODOM SPScan HMMER
3	374	T334 T33 S137 T146 S291 S311 T346	N103 N127 N135 N138	Signal peptide: M1-N52	Tricarboxylate carrier [Rattus sp.] g545998	MOTIFS BLAST-GenBank SPScan
4	271	S234 T126 T169 Y141		Signal peptide: M1-C30 Transmembrane region: L233-F252	Weak similarity with honeybee ATP synthase A chain [Caenorhabditis elegans] g3878801	MOTIFS BLAST-GenBank SPScan HMMER
5	323	S99 S125 S192 T277 S307 S309 T110 Y212		Leucine zipper: L284-L305	Cu <sup>2+</sup> -transporting ATPase homolog [Arabidopsis thaliana] g2464854	BLAST-GenBank MOTIFS
6	274	S96 T198 S215 T29 S121 S164 S170		Mitochondrial energy transfer proteins: G5-L266 Signal peptide: M1-G17	Pet8p [Saccharomyces cerevisiae] g495307	BLAST-GenBank HMMER-PFAM MOTIFS ProfileScan BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO SPScan

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
7	291	S6 T113 T173 T147 S230 T258	N226 N261	Signal peptide: M1-T42 Transmembrane domain: W29-I54 Band 7 protein family: C50-V62, K90-E224 Membrane stomatin: E14-N283	Stomatin [Homo sapiens] g1161562	MOTIFS BLAST-GenBank SPScan HMMER BLIMPS-BLOCKS BLAST-DOMO BLAST-PRODOM
8	381	S2 S35 T57 S92 T104 S191 S302 S334 S335 S336 T43 T250 T255 T304 S311 S370 Y65	N218 N253 N259		K <sup>+</sup> channel modulatory factor DEBT-91 [Mus musculus] g4838557	MOTIFS BLAST-GenBank
9	190	T160 S17 T71 S77 T78 S111 S134 S142	N87	ABC transporter family: R79-K177 ATP/GTP-binding site motif A (P-loop): G102-S109	ABC2 transporter [Mus musculus] g495259	MOTIFS BLAST-GenBank BLAST-DOMO
10	297	S17 S114 T136 S16	N287	Mitochondrial carrier protein signature: E117-I297 Graves Disease carrier protein: P137-T157, L259-S279	Similar to human ADP/ATP carrier protein [C. elegans] g3879938	MOTIFS BLAST-GenBank HMMER-PFAM BLIMPS-PRINTS
11	89	T37 T47 T60 S64			Mitochondrial import protein Tim9p [Saccharomyces cerevisiae] g3747026	BLAST-GenBank MOTIFS
12	115	T108 T84		Signal peptide: M1-G24 Transmembrane domain: G35-F57 Sodium neurotransmitter symporter signature: R7-S61		MOTIFS SPScan HMMER ProfileScan

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
13	675	T54 T50 S99 T127 S413 T558 S645 T654 T47 S242 T602 T611 Y501	N243 N247 N301 N601	Transmembrane domains: I29-V48, L103-I121, L177-G196, I210-M229, L417-W435, F481-Y501, Y521-W541 Sodium symporter family domain: Y58-G487 Sodium:solute symporter signature: Y35-G89, M111-R140, L173-G227, P460-G469	Sodium-glucose cotransporter [Oryctolagus cuniculus] g473969	BLAST-GenBank MOTIFS HMMER HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLAST-PRODOM BLAST-DOMO
14	320	T84 S304 T11 S75 S80 S164 Y20	N162 N234	Transmembrane domains: I92-L112, I201-K219 Zinc transporter signature: A28-V142, D199-E303 Cation transporter domain: S48-L74	Zinc transporter Znf-2 [Rattus norvegicus] g1256378	BLAST-GenBank MOTIFS HMMER BLIMPS-PRODOM BLAST-PRODOM BLAST-DOMO
15	462	S111 S145 S183 S233 T26 T185 S202 T243	N24 N279	Kelch repeat motifs: C299-N349; F350-R399 Y400-G446 BTB domain: F50-L117 POZ domain: Y27-E215	Ring canal protein [Drosophila melanogaster] g577276	BLAST-GenBank MOTIFS HMMER-PFAM BLIMPS-PRINTS BLAST-DOMO
16	98	T22 Y37		Signal peptide: M1-S17 Mitochondrial carrier proteins domain: C4-I89 Mitochondrial carrier proteins signature sequence: V6-G19, G19-A33, G63-E83	Carrier protein (cl) [Caenorhabditis elegans] g472902	BLAST-GenBank MOTIFS SPScan HMMER-PFAM ProfileScan BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-DOMO

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
17	748	S55 S196 T254 S307 S327 T491 T534 T550 T571 S635 S648 S677 T696 S283 S291 T314 S629 S701 Y556	N531 N543 N548 N627	Signal peptide: M1-A61 Transmembrane domains: L39-L56, I167-F186, C229-F252, G438-L455, M492-F509, L598-I618 Ion transport proteins signature: F85-V251, L369-I618	Voltage-gated calcium channel [Rattus norvegicus] g4586963	BLAST-GenBank MOTIFS SPScan HMMER HMMER-PFAM BLIMPS-PRINTS
18	507	T200 S183 T232 T284 T349 T150 T252 S253 S319 S383 Y454	N220 N250 N364 N496	Signal peptide: M1-G26	Nucleoporin p54 [Rattus norvegicus] g1537070	BLAST-GenBank MOTIFS SPScan
19	592	S460 S104 T178 S320 S321 T498 T531 Y365		ABC precursor signature: N153-Q162, F210-A229, G234-I254, V312-G332, T366-V378	ABC transporter [Methanobacterium thermo.] g2622773	BLAST-GenBank MOTIFS BLIMPS-PRODOM BLAST-PRODOM
20	841	T98 S120 S203 T214 T276 S388 T438 T700 T838 T167 T179 S280 T370 S435 S531 S539 S666 S693 S830	N368 N490 N624	Transmembrane domains: Y451-D469, M544-F562, F577-F597, G775-W797 Vacuolar ion transport subunit signature: M10-F831	Vacuolar H <sup>+</sup> /ATPase subunit [Rattus norvegicus] g206430	BLAST-GenBank MOTIFS HMMER BLIMPS-PRODOM BLAST-PRODOM BLAST-DOMO
21	253	S50 T139 T152 T177 S202 T143 Y55		Mitochondrial carrier proteins domain: Y31-S248 Mitochondrial energy transfer proteins signature sequence: I62-Q86, I110-G122	Mitochondrial uncoupling protein UCP-4 [Homo sapiens] g4324701	BLAST-GenBank MOTIFS HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLAST-PRODOM BLAST-DOMO

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
22	229	S69 S26 S109 T162 S178 S25 S64 S65 T210 S219		Signal peptide: M1-A47 Mitochondrial carrier proteins domain: Q32-G220 Mitochondrial carrier proteins signature sequence: S36-T49, T49-V63, G92-E112, T144-T162, Y187-F205	Grave's disease carrier protein [Bos taurus] g387	BLAST-GenBank MOTIFS SPScan HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO
23	170	S26 S31 S149 S164 T22 T157	N66 N145	Dihydroxypyridine-sensitive L-type calcium channel signature: Y2-A47, I49-V77, A83-N100, R106-E131 SH3 domain: V59-R122	Voltage-dependent calcium channel beta-4 subunit [Homo sapiens] g2058727	BLAST-GenBank MOTIFS HMMER-PFAM BLIMPS-BLOCKS BLIMPS-PRINTS BLIMPS-PFAM BLAST-PRODOM BLAST-DOMO
24	655	T194 S195 S232 T362 S655 S4 S88 T135 T153 S187 T214 S322 T345 S353 S443 T609 S261 S381 S384	N338 N418 N557 N596	Transmembrane domains: I396-K417, Y494-S522, T538-V556 ABC transporters domain: P73-G262 ABC transporter family signature sequence: I78-L89, V186-D217	Breast cancer resistance protein (multidrug transporter) [Homo sapiens] g4038352	BLAST-GenBank MOTIFS HMMER HMMER-PFAM BLIMPS-BLOCKS ProfileScan BLAST-PRODOM BLAST-DOMO
25	184	T51 S29 T100 S138 S151 Y78	N27		Cation transport protein [E. coli] g495778	BLAST-GenBank MOTIFS
26	154	S54 S42 S62 T78 Y104		Mitochondrial energy transfer proteins signatures: P89-L97, M1-E41, M73-L152 Mitochondrial carrier protein domain: G2-L152	Similar to carrier protein C2 [C. elegans] g3879669	MOTIFS HMMER-PFAM BLAST-PRODOM BLAST-DOMO BLAST-GenBank



Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
27	438	S170 T5 T51 T265 T300 S425	N50 N423	Transmembrane domains: C91-L111, L237-I257, I305-M332, M332-L354, L391-V408, I186-A204	Multidrug efflux transporter [Bacillus subtilis] g2635104	MOTIFS HMMER BLAST-GenBank
28	237	S10 S47 T72 S28 S96 S148 T173 T222 S6 S21 T32 T61 T192	N35	Nucleic acid-binding protein E5.1 domain: S6-K128	ARL-6 interacting protein-4 [Mus musculus] g4927204	MOTIFS BLAST-DOMO BLAST-GenBank
29	219	T66 S194 T200		Signal peptide: M1-R19 or M1-K15 Caseins alpha/beta signature: M1-N39	Surface antigen [Trypanosoma cruzi] g161956	MOTIFS HMMER SPScan ProfileScan BLAST-GenBank
30	707	S31 T6 T55 T263 T328 T546 T580 T594 S662 S673 T32 S50 S231 T244 T306 T385 S439 S476 S533 S553 S624	N343 N570 N638 N703	Potassium channel signature: A62-T81 Potassium channel integral membrane protein domain: S13-D117	NY-REN-45 antigen (similar to potassium channel protein) [Homo sapiens] g5360115	MOTIFS BLIMPS-PRINTS BLAST-DOMO BLAST-GenBank
31	279	T18 T245 T206	N181	Signal cleavage: M1-G45 Connexin domains: M1-V99, V20-Y44 Connexin signatures: L33-V86, L152-F205, F51-P73, S76-L96, L133-Y159, C169-T189, I190-L218 Gap junction protein connexin transmembrane regions: F5-Y97, L133- K223, M1-S130	Gap junction protein (similar to connexin) [Homo sapiens] g3006230	MOTIFS SPScan HMMER BLIMPS-BLOCKS BLIMPS-PRINTS BLAST-PFAM ProfileScan BLAST-PRODOM BLAST-DOMO BLAST-GenBank

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
32	154	S114		Signal peptide: M1-A35 or M1-A14 Transmembrane domain: F83-L102	mBOCT (potent organic cation transporter) [Mus musculus] g4589468	MOTIFS HMHER SPScan BLAST-GenBank
33	289	T83 T205 S269 T279	N60	Mitochondrial energy transfer proteins signatures: M1-G147, P17, P115, N185-K280, A101-Q181, Y184-I278 Mitochondrial carrier protein domains: M1-E176, N185-K280 Mitochondrial transmembrane transport protein regions: P17-R182, P180-I278	Mitochondrial solute carrier [Onchocerca volvulus] g1518458	MOTIFS HMHER-PFAM BLAST-DOMO BLAST-PRODOM ProfileScan BLAST-GenBank
34	300	S189 S195 S204 T257		Mitochondrial energy transfer proteins signatures: P19-M27, D2-I53, L209-L295 Mitochondrial carrier protein domain: D2-Y295 Transport protein domain: P122-Y295	YKL522=mitochondria 1 ADP/ATP carrier protein homolog [Saccharomyces cerevisiae] g254449	MOTIFS HMHER-PFAM ProfileScan BLAST-PRODOM BLAST-DOMO BLAST-GenBank
35	382	S34 S207 T221 S312 T40 S53 T112 T117 T277 S337	N96 N372	Kelch motifs: H191-G249, E250-D301	Similarity to Human host cell factor C1 [Homo sapiens] g3875291	MOTIFS HMHER BLAST-PFAM BLAST-GenBank

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
36	287	T36 T118 S180 S230 T84 S168 T244		Mitochondrial energy transfer proteins signatures: P26-L34, P219-L227, L97-G193, W10-V89, D197-F281, P96-Y194 Mitochondrial carrier protein domain: A5-F281 Mitochondrial brown fat region: Y82-Q94, V151-S168, Y194-C212	Mitochondrial dicarboxylate carrier [Rattus norvegicus] g3646426	MOTIFS HMMER-PFAM BLIMPS-PRINTS BLAST-DOMO BLAST-GenBank
37	497	T65 T135 S147 T360 S8 T22 S45 S291	N63 N314 N414	Transmembrane domains: M114-T137, M364-M380, Y390-A413, A421-D444, F456-V478 Folate transporter domains: W30-H218, I253-K484	Reduced folate carrier [Homo sapiens] g1041934	MOTIFS HMMER BLAST-PRODOM BLAST-DOMO SPScan BLAST-GenBank
38	228	T21 S124 T145 S158 T190 T95 S132 S137 T177		Heme-binding domain in cytochrome b5: Y19-G98 Cytochrome b5 family domain: H28-P75	cytochrome b5 containing fusion protein [Helianthus annuus] g1040729 P=1.2e-07	MOTIFS HMMER-PFAM BLAST-GenBank ProfileScan
39	273	T63 S158 T48	N214	Transmembrane domains: L85-N105, F180-Y200 Intermembrane space domain: L30-L251	Sqv-7-like protein (similar to nucleotide-sugar transporters) [Homo sapiens] g4008517	MOTIFS HMMER BLAST-DOMO BLAST-GenBank

Table 2 (cont.)

SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Motifs, and Domains	Homologous Sequences	Analytical Methods & Databases
40	206	S187 S201	N158	Signal peptide: M1-G29 or M1-A27 Emopamil binding protein: G37-S187, L15-K203 Transmembrane domain: Y164-L183	C-8,7 sterol isomerase, aSI1 [Arabidopsis thaliana] g2772934	MOTIFS HMMER ProfileScan BLAST-DOMO BLAST-GenBank
41	235	S192 S200 S56 T95 T146 S199 T207 S229 T53 T61 T69 T119 T148 Y70	N123	Transmembrane domain: F15-I34, M155-V174 Channel myelin protein: L18-M181 Sodium channel beta-2 subunit precursor: F15-E210 Immunoglobulin domain: I34-V136	Myelin protein zero (MPZ) [Homo sapiens] g2160399	MOTIFS HMMER BLIMPS-PRINTS BLAST-PRODOM BLAST-DOMO BLAST-GenBank
42	147	T79 T116 S3 S66 Y89 Y98	N118	Signal peptide: M1-G23 or M1-A20 Transthyretin signature: S28-S132 Transthyretin domain: G21-Q146	Transthyretin precursor [Sus scrofa] g1009702	MOTIFS HMMER ProfileScan BLAST-PRODOM BLAST-DOMO BLAST-GenBank BLIMPS-BLOCKS BLIMPS-PRINTS
43	147	T5 S88 T39		Globin domain: V2-H147 Heme oxygen transport protein domain: L32-H147	III beta-3 globin [Rattus norvegicus] g395943	MOTIFS HMMER-PFAM BLAST-PRODOM BLAST-DOMO BLIMPS-BLOCKS BLIMPS-PRINTS

Table 3

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
44	1567-1611 2107-2151	Gastrointestinal (0.203) Hematopoietic/Immune (0.188) Nervous (0.156)	Cell Proliferation and Cancer (0.547) Inflammation (0.422)	PBLUESCRIPT
45	1-92 351-434	Endocrine (0.333) Developmental (0.167) Gastrointestinal (0.167) Musculoskeletal (0.167) Reproductive (0.167)	Cell Proliferation and Cancer (0.833) Inflammation (0.167)	pINCY
46	920-964 1352-1396	Reproductive (0.304) Gastrointestinal (0.174) Cardiovascular (0.130) Hematopoietic/Immune (0.130) Nervous (0.130)	Cell Proliferation and Cancer (0.478) Inflammation (0.391)	PSPORT1
47	1-80 768-848	Nervous (0.273) Reproductive (0.273) Gastrointestinal (0.127) Hematopoietic/Immune (0.127)	Cell Proliferation and Cancer (0.564) Inflammation (0.400)	pINCY
48	111-194 687-758	Reproductive (0.221) Nervous (0.185) Gastrointestinal (0.124)	Cell Proliferation and Cancer (0.552) Inflammation (0.343)	pINCY
49	1-97	Nervous (0.234) Hematopoietic/Immune (0.191) Gastrointestinal (0.149)	Cell Proliferation and Cancer (0.617) Inflammation (0.340)	pINCY
50	218-262	Cardiovascular (1.000)	Cancer (0.333) Inflammation/Trauma (0.333) Cell Proliferation (0.333)	PSPORT1
51	811-855	Hematopoietic/Immune (0.180) Gastrointestinal (0.146) Reproductive (0.281)	Cancer (0.393) Inflammation/Trauma (0.515) Cell Proliferation (0.146)	PSPORT1
52	595-639	Gastrointestinal (0.286) Reproductive (0.714)	Cancer (0.429) Inflammation/Trauma (0.429)	pINCY
53	96-140	Cardiovascular (0.167) Hematopoietic/Immune (0.167) Nervous (0.250) Reproductive (0.167)	Cancer (0.250) Inflammation/Trauma (0.167) Cell Proliferation (0.167)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
54	507-551	Reproductive (0.323) Gastrointestinal (0.154) Nervous (0.123)	Cancer (0.446) Inflammation/Trauma (0.308) Cell Proliferation (0.185)	PBLUESCRIPT
55	455-499	Urologic (0.333) Nervous (0.222) Reproductive (0.222)	Cancer (0.667) Cell Proliferation (0.333)	PBLUESCRIPT
56	1835-1879	Nervous (0.625) Gastrointestinal (0.375)	Inflammation/Trauma (0.375) Cancer (0.250) Neurological (0.250)	pINCY
57	811-855	Gastrointestinal (1.000)	Inflammation/Trauma (0.667)	pINCY
58	390-434	Reproductive (0.320) Nervous (0.240) Urologic (0.120)	Cancer (0.520) Inflammation/Trauma (0.240) Cell Proliferation (0.160)	PSPORT1
59	413-457	Gastrointestinal (0.333) Musculoskeletal (0.333) Nervous (0.333)	Cancer (0.333) Neurological (0.333)	pINCY
60	2021-2084	Nervous (0.197) Gastrointestinal (0.184) Reproductive (0.184)	Cancer (0.461) Inflammation/Trauma (0.316) Cell Proliferation (0.118)	pINCY
61	65-109	Nervous (0.226) Reproductive (0.208) Cardiovascular (0.113) Gastrointestinal (0.113)	Cancer (0.528) Inflammation/Trauma (0.301) Cell Proliferation (0.208)	pINCY
62	379-423 1867-1911	Reproductive (0.282) Gastrointestinal (0.205) Nervous (0.154)	Cancer (0.538) Inflammation/Trauma (0.282) Cell Proliferation (0.103)	PSPORT1
63	362-406 1193-1237	Urologic (0.500) Reproductive (0.333) Cardiovascular (0.167)	Cancer (0.667) Inflammation/Trauma (0.333)	pINCY
64	394-438	Nervous (0.294) Reproductive (0.265) Cardiovascular (0.118)	Cancer (0.382) Inflammation/Trauma (0.235) Cell Proliferation (0.118)	pINCY

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
65	768-812	Reproductive (0.300) Endocrine (0.200) Gastrointestinal (0.200) Hematopoietic/Immune (0.200)	Inflammation/Trauma (0.500) Cancer (0.400)	pINCY
66	77-121	Nervous (1.000)	Neurological (1.000)	pINCY
67	1999-2043	Reproductive (0.324) Nervous (0.265) Gastrointestinal (0.235)	Cancer (0.500) Inflammation/Trauma (0.294) Cell Proliferation (0.118)	pINCY
68	561-605	Hematopoietic/Immune (0.455) Gastrointestinal (0.182) Nervous (0.182)	Inflammation/Trauma (0.546) Cell Proliferation (0.182)	pINCY
69	679-729	Nervous (0.292) Gastrointestinal (0.208) Hematopoietic/Immune (0.125)	Cancer (0.250) Cell Proliferation (0.375) Inflammation/Trauma (0.416)	PBLUESCRIPT
70	95-366 1078-1185	Reproductive (0.206) Hematopoietic/Immune (0.186) Cardiovascular (0.127)	Cancer (0.373) Inflammation/Trauma (0.382) Cell Proliferation (0.176)	PBLUESCRIPT
71	33-152	Reproductive (0.275) Nervous (0.163) Gastrointestinal (0.137)	Cancer (0.438) Inflammation/Trauma (0.314) Cell Proliferation (0.176)	PSPORT1
72	81-779	Gastrointestinal (1.000)	Cancer (1.000)	pINCY
73	719-817 1202-1414	Reproductive (0.311) Hematopoietic/Immune (0.203) Gastrointestinal (0.122)	Cancer (0.459) Inflammation/Trauma (0.379) Cell Proliferation (0.203)	PSPORT1
74	1-848	Nervous (0.750) Dermatologic (0.250)	Cancer (0.250) Cell Proliferation (0.250) Inflammation/Trauma (0.500)	pINCY
75	1-478	Cardiovascular (0.714) Developmental (0.143) Hematopoietic/Immune (0.143)	Cancer (0.571) Cell Proliferation (0.286) Inflammation (0.143)	pINCY
76	1-134	Reproductive (0.253) Nervous (0.241) Gastrointestinal (0.127) Hematopoietic (0.127)	Cancer (0.494) Inflammation (0.215) Cell Proliferation (0.127)	PSPORT1

Table 3 (cont.)

Nucleotide SEQ ID NO:	Selected Fragments	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
77	510-719 960-1100	Reproductive (0.467) Cardiovascular (0.133) Gastrointestinal (0.133)	Cancer (0.467) Inflammation/Trauma (0.467)	pINCY
78	180-293	Reproductive (0.230) Nervous (0.225) Gastrointestinal (0.124)	Cancer (0.478) Inflammation/Trauma (0.292) Cell Proliferation (0.191)	pINCY
79	192-653 795-935	Reproductive (0.417) Gastrointestinal (0.292) Urologic (0.125)	Cancer (0.750) Cell Proliferation (0.125) Inflammation/Trauma (0.167)	pINCY
80	139-1044	Reproductive (0.245) Nervous (0.143) Developmental (0.122)	Cancer (0.490) Inflammation/Trauma (0.286) Cell Proliferation (0.224)	pINCY
81	233-916	Reproductive (0.667) Cardiovascular (0.167) Nervous (0.167)	Cancer (0.500) Cell Proliferation (0.333) Inflammation (0.167)	pINCY
82	1-153 760-816	Gastrointestinal (0.282) Hematopoietic/Immune (0.205) Reproductive (0.205)	Inflammation/Trauma (0.461) Cancer (0.308) Cell Proliferation (0.205)	pINCY
83	57-299	Nervous (0.179) Reproductive (0.179) Gastrointestinal (0.128)	Cancer (0.564) Cell Proliferation (0.256) Inflammation/Trauma (0.180)	pINCY
84	1-707	Gastrointestinal (0.500) Hematopoietic/Immune (0.500)	Cancer (0.500) Inflammation (0.500)	pINCY
85	451-594	Hematopoietic/Immune (1.000)	Cell Proliferation (1.000)	pINCY
86	8-124 161-187 407-472	Developmental (1.000)	Cell Proliferation (1.000)	pINCY



Table 4

SEQ ID NO.	Library	Library Comment
44	HNT2AGT01	Library was constructed at Stratagene (STR937233), using RNA isolated from the HNT2 cell line derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor. Cells were treated with retinoic acid for 5 weeks and with mitotic inhibitors for two weeks and allowed to mature for an additional 4 weeks in conditioned medium.
45	COLNFET02	Library was constructed using RNA isolated from the colon tissue of a Caucasian female fetus, who died at 20 weeks' gestation.
46	PANCN0T04	Library was constructed using RNA isolated from the pancreatic tissue of a 5-year-old Caucasian male, who died in a motor vehicle accident. Serology was positive for cytomegalovirus (CMV).
47	ADRETUT05	Library was constructed from adrenal tumor tissue removed from a 52-year-old Caucasian female during a unilateral adrenalectomy. Pathology indicated a pheochromocytoma.
48	LUNGTTUT11	Library was constructed using RNA isolated from lung tumor tissue removed from the right lower lobe of a 57-year-old Caucasian male during a segmental lung resection. Pathology indicated an infiltrating grade 4 squamous cell carcinoma. Multiple intrapulmonary peribronchial lymph nodes showed metastatic squamous cell carcinoma. Patient history included a benign brain neoplasm and tobacco abuse. Family history included spinal cord cancer, type II diabetes, cerebrovascular disease, and malignant prostate neoplasm.
49	BRAVTXT03	Library was constructed using RNA isolated from treated astrocytes removed from the brain of a female fetus who died after 22 weeks' gestation. The cells were treated with tumor necrosis factor (TNF) alpha and interleukin 1 (IL-1), 10ng/ml each for 24 hours.
50	LUNGAST01	Library was constructed using RNA isolated from the lung tissue of a 17-year-old Caucasian male, who died from head trauma. Patient history included asthma.
51	OVARNOT02	Library was constructed using RNA isolated from ovarian tissue removed from a 59-year-old Caucasian female who died of a myocardial infarction. Patient history included cardiomyopathy, coronary artery disease, previous myocardial infarctions, hypercholesterolemia, hypotension, and arthritis.
52	BRSTNOT13	Library was constructed using RNA isolated from breast tissue removed from the left medial lateral breast of a 36-year-old Caucasian female during bilateral simple mastectomy and total breast reconstruction. Pathology indicated benign breast tissue. Patient history included a breast neoplasm, depressive disorder, hyperlipidemia, chronic stomach ulcer, and an ectopic pregnancy. Family history included myocardial infarction, cerebrovascular disease, atherosclerotic coronary artery disease, hyperlipidemia, skin cancer, breast cancer, depressive disorder, esophageal cancer, bone cancer, Hodgkin's lymphoma, bladder cancer, and a heart condition.

Table 4 (cont.)

SEQ ID NO:	Library	Library Comment
53	SMCCNOS01	Library was constructed using 7.56 X 10e6 clones from a coronary artery smooth muscle cell library and was subjected to two rounds of subtraction hybridization for 48 hours with 6.12 X 10e6 clones from a control coronary artery smooth muscle cell library. The starting library for subtraction was constructed using RNA isolated from coronary artery smooth muscle cells removed from a 3-year-old Caucasian male. The cells were treated with TNF alpha & IL-1 beta 10ng/ml each for 20 hours. The hybridization probe for subtraction was derived from a similarly constructed library from RNA isolated from untreated coronary artery smooth muscle cells from the same donor.
54	HUVENOB01	Library was constructed using RNA isolated from HUV-EC-C (ATCC CRL 1730) cells.
55	HNT2RAT01	Library was constructed at Stratagene (STR937231), using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated with retinoic acid for 24 hours.
56	SINTBST01	Library was constructed using RNA isolated from ileum tissue obtained from an 18-year-old Caucasian female during bowel anastomosis. Pathology indicated Crohn's disease of the ileum, involving 15 cm of the small bowel. Family history included cerebrovascular disease and atherosclerotic coronary artery disease.
57	ISLTNOT01	Library was constructed using RNA isolated from a pooled collection of pancreatic islet cells.
58	COLNNOT11	Library was constructed using RNA isolated from colon tissue removed from a 60-year-old Caucasian male during a left hemicolectomy.
59	BONRTUT01	Library was constructed using RNA isolated from rib tumor tissue removed from a 16-year-old Caucasian male during a rib osteotomy and a wedge resection of the lung. Pathology indicated metastatic grade 3 (of 4) osteosarcoma, forming a mass involving the chest wall.
60	LUNGUT10	Library was constructed using RNA isolated from lung tumor tissue removed from the left upper lobe of a 65-year-old Caucasian female during a segmental lung resection. Pathology indicated metastatic grade 2 myxoid liposarcoma and metastatic grade 4 liposarcoma. Patient history included soft tissue cancer, breast cancer, and secondary lung cancer.
61	OVARNOT09	Library was constructed using RNA isolated from ovarian tissue removed from a 28-year-old Caucasian female during a vaginal hysterectomy and removal of the fallopian tubes and ovaries. Pathology indicated multiple follicular cysts ranging in size from 0.4 to 1.5 cm in the right and left ovaries, chronic cervicitis and squamous metaplasia of the cervix, and endometrium in weakly proliferative phase. Family history included benign hypertension, hyperlipidemia, and atherosclerotic coronary artery disease.

Table 4 (cont.)

SEQ ID NO:	Library	Library Comment
62	THPLAZS08	Library was constructed using 5.76 million clones from a 5-aza-2'-deoxycytidine (AZ) treated THP-1 promonocyte cell line library. Starting RNA was made from THP-1 promonocyte cells treated for three days with 0.8 micromolar AZ. 5.76 million clones from the AZ-treated THP-1 cell library were then subjected to two rounds of subtractive hybridization with 5 million clones from the untreated THP-1 cell library. Subtractive hybridization conditions were based on the methodologies of Swaroop et al. (1991) Nucleic Acids Res. 19:1954, and Bonaldo et al. (1996) Genome Research 6:791. THP-1 (ATCC TIB 202) is a human promonocyte cell line derived from peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia (ref: Int. J. Cancer (1980) 26:171).
63	BRSTNOT12	Library was constructed using RNA isolated from diseased breast tissue removed from a 32-year-old Caucasian female during a bilateral reduction mammoplasty. Pathology indicated nonproliferative fibrocystic disease. Family history included benign hypertension and atherosclerotic coronary artery disease.
64	PENCNOT09	Library was constructed using RNA isolated from penis right corpora cavernosa tissue.
65	PROSTUS19	Library was constructed using 2.36 million clones from a prostate tumor library which was subjected to two rounds of subtraction hybridization with 2.36 million clones from a normal prostate library. The starting library for subtraction was constructed using RNA isolated from prostate tumor tissue removed from a 59-year-old Caucasian male during a radical prostatectomy with regional lymph node excision. Pathology indicated adenocarcinoma (Gleason grade 3+3) involving the prostate peripherally with invasion of the capsule. Adenofibromatous hyperplasia was present. The patient presented with elevated prostate-specific antigen (PSA). Patient history included diverticulitis of the colon, asbestosis, and thrombophlebitis. Family history included benign hypertension, multiple myeloma, hyperlipidemia, and rheumatoid arthritis. The hybridization probe for subtraction was derived from a similarly constructed library, except that NotI-anchored oligo(dT) primer was used. Subtractive hybridization conditions were based on the methodologies of Swaroop et al. (1991) Nucleic Acids Res. 19:1954 and Bonaldo, et al. (1996) Genome Research 6:791.
66	BRABDIR01	Library was constructed using RNA isolated from diseased cerebellum tissue removed from the brain of a 57-year-old Caucasian male, who died from a cerebrovascular accident. Patient history included Huntington's disease, emphysema, and tobacco abuse.
67	LIVRDIR01	Library was constructed using RNA isolated from diseased liver tissue removed from a 63-year-old Caucasian female during a liver transplant. Patient history included primary biliary cirrhosis. Serology was positive for anti-mitochondrial antibody.

Table 4 (cont.)

SEQ ID NO:	Library	Library Comment
68	COLCDIT03	Library was constructed using RNA isolated from diseased colon polyp tissue removed from the cecum of a 67-year-old female. Pathology indicated a benign cecum polyp. Pathology for the associated tumor tissue indicated invasive grade 3 adenocarcinoma that arose in tubulovillous adenoma forming a fungating mass in the cecum.
69	TBLYNOT01	Library was constructed at Stratagene (STR937214) using RNA isolated from a hybrid of T-B lymphoblasts from an untreated leukemic cell line.
70	KIDNNOT01	Library was constructed using RNA isolated from the kidney tissue of a 64-year-old Caucasian female, who died from an intracranial bleed. Patient history included rheumatoid arthritis and tobacco use.
71	LUNGAST01	Library was constructed using RNA isolated from the lung tissue of a 17-year-old Caucasian male, who died from head trauma. Patient history included asthma.
72	LPARNOT02	Library was constructed using RNA isolated from tissue obtained from the left parotid (salivary) gland of a 70-year-old male with parotid cancer.
73	OVARNOT03	Library was constructed using RNA isolated from ovarian tissue removed from a 43-year-old Caucasian female during a bilateral salpingo-oophorectomy. Pathology for the associated tumor tissue indicated grade 2 mucinous cyst adenocarcinoma. The patient presented with stress incontinence. Patient history included mitral valve disorder, pneumonia, and viral hepatitis. Family history included atherosclerotic coronary artery disease, cerebrovascular disease, breast cancer, and uterine cancer.
74	ENDCNOT03	Library was constructed using RNA isolated from dermal microvascular endothelial cells removed from a neonatal Caucasian male.
75	LUNGNOT18	Library was constructed using RNA isolated from left upper lobe lung tissue removed from a 66-year-old Caucasian female. Pathology for the associated tumor tissue indicated a grade 2 adenocarcinoma. Patient history included cerebrovascular disease, atherosclerotic coronary artery disease, and pulmonary insufficiency. Family history included a myocardial infarction and atherosclerotic coronary artery disease.
76	NGANNOT01	Library was constructed using RNA isolated from tumorous neuroganglion tissue removed from a 9-year-old Caucasian male during a soft tissue excision of the chest wall. Pathology indicated a ganglioneuroma. Family history included asthma.
77	LUNGTUT09	Library was constructed using RNA isolated from lung tumor tissue removed from a 68-year-old Caucasian male during segmental lung resection. Pathology indicated invasive grade 3 squamous cell carcinoma and a metastatic tumor. Patient history included type II diabetes, thyroid disorder, depressive disorder, hyperlipidemia, esophageal ulcer, and tobacco use.

Table 4 (cont.)

SEQ ID NO:	Library	Library Comment
78	ESOGTUT02	Library was constructed using RNA isolated from esophageal tumor tissue obtained from a 61-year-old Caucasian male during a partial esophagectomy, proximal gastrectomy, pyloromyotomy, and regional lymph node excision. Pathology indicated an invasive grade 3 adenocarcinoma in the esophagus. Family history included atherosclerotic coronary artery disease, type II diabetes, chronic liver disease, primary cardiomyopathy, benign hypertension, and cerebrovascular disease.
79	KIDNOT19	Library was constructed using RNA isolated from kidney tissue removed a 65-year-old Caucasian male during an exploratory laparotomy and nephroureterectomy. Pathology for the associated tumor tissue indicated a grade 1 renal cell carcinoma within the upper pole of the left kidney. Patient history included malignant melanoma of the abdominal skin, benign neoplasm of colon, cerebrovascular disease, and umbilical hernia. Family history included myocardial infarction, atherosclerotic coronary artery disease, cerebrovascular disease, and prostate cancer.
80	TLYJINT01	Library was constructed using RNA isolated from a Jurkat cell line derived from the T cells of a male. Patient history included acute T-cell leukemia. This is an uninduced Jurkat cell line library from the same donor.
81	PENCNOT06	Library was constructed using RNA isolated from penis corpora cavernosa tissue removed from a 3-year-old Black male. Pathology for the associated tumor tissue indicated invasive grade 4 urothelial carcinoma forming a soft tissue scrotal mass that invaded the cavernous body of the penis and encased both testicles. Right inguinal lymph node showed metastatic grade 4 urothelial carcinoma, with extranodal invasion.
82	UTRSTUT05	Library was constructed using RNA isolated from uterine tumor tissue removed from a 41-year-old Caucasian female during a vaginal hysterectomy with dilation and curettage. Pathology indicated uterine leiomyoma. The endometrium was secretory and contained fragments of endometrial polyps. Benign endo- and ectocervical mucosa were identified in the endocervix. Patient history included a ventral hernia and a benign ovarian neoplasm.
83	LUNGNOT37	Library was constructed using polyA RNA isolated from lung tissue removed from a 15-year-old Caucasian female who died from a closed head injury. Serology was positive for cytomegalovirus.
84	LIVRTUT09	Library was constructed using RNA isolated from an untreated C3A hepatocyte cell line which is a derivative of Hep G2, a cell line derived from a hepatoblastoma removed from a 15-year-old Caucasian male.
85	MCLRUNT01	Library was constructed using RNA isolated from untreated peripheral blood mononuclear cell tissue obtained from buffy coat, removed from a 60-year-old male.
86	MCLRUNT01	Library was constructed using RNA isolated from untreated peripheral blood mononuclear cell tissue obtained from buffy coat, removed from a 60-year-old male.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	PE Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	PE Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	PE Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Atwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score $\geq$ GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising an amino acid sequence selected from the group consisting of:
  - 5 a) an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28,
    - 10 SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43,
  - b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID
    - 15 NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35,
      - 20 SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43,
  - c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID
    - 25 NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43, and
  - 30 d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID
    - 35 NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID



NO:41, SEQ ID NO:42, and SEQ ID NO:43.

2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43.

3. An isolated polynucleotide encoding a polypeptide of claim 1.

4. An isolated polynucleotide encoding a polypeptide of claim 2.

5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86.

6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.

7. A cell transformed with a recombinant polynucleotide of claim 6.

8. A transgenic organism comprising a recombinant polynucleotide of claim 6.

9. A method for producing a polypeptide of claim 1, the method comprising:

a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

b) recovering the polypeptide so expressed.

10. An isolated antibody which specifically binds to a polypeptide of claim 1.

5 11. An isolated polynucleotide comprising a polynucleotide sequence selected from the group consisting of:

a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,

15 b) a naturally occurring polynucleotide sequence having at least 70% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:44, SEQ ID NO:45, SEQ ID NO:46, SEQ ID NO:47, SEQ ID NO:49, SEQ ID NO:50, SEQ ID NO:51, SEQ ID NO:52, SEQ ID NO:53, SEQ ID NO:55, SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58, SEQ ID NO:59, SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62, SEQ ID NO:63, SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66, SEQ ID NO:67, SEQ ID NO:68, SEQ ID NO:69, SEQ ID NO:70, SEQ ID NO:71, SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74, SEQ ID NO:75, SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:85, SEQ ID NO:86,

20 c) a polynucleotide sequence complementary to a),  
 25 d) a polynucleotide sequence complementary to b), and  
 e) an RNA equivalent of a)-d).

12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.

30

13. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe  
 35 specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and

b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

5 14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.

15 15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:

a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and

10 b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

16. A pharmaceutical composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

15

17. A pharmaceutical composition of claim 16, wherein the polypeptide comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:41, SEQ ID NO:42, and SEQ ID NO:43.

25 18. A method for treating a disease or condition associated with decreased expression of functional TPPT, comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

30

a) exposing a sample comprising a polypeptide of claim 1 to a compound, and

b) detecting agonist activity in the sample.

20. A pharmaceutical composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

35

21. A method for treating a disease or condition associated with decreased expression of functional TPPT, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 20.
- 5        22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:
- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
  - b) detecting antagonist activity in the sample.
- 10       23. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.
- 15       24. A method for treating a disease or condition associated with overexpression of functional TPPT, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 23.
25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:
- a) combining the polypeptide of claim 1 with at least one test compound under suitable
  - 20 conditions, and
  - b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.
26. A method of screening for a compound that modulates the activity of the polypeptide of
- 25 claim 1, said method comprising:
- a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
  - b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound,
  - and
  - 30 c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.
- 35       27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method

comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

5           28. An isolated polynucleotide comprising a polynucleotide sequence of SEQ ID NO:83.

          29. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 28.

10          30. A cell transformed with a recombinant polynucleotide of claim 29.

          31. A transgenic organism comprising a recombinant polynucleotide of claim 29.

          32. A method for producing a polypeptide comprising an amino acid sequence of SEQ ID  
15 NO:40, the method comprising:

- a) culturing the cell of claim 30 under conditions suitable for expression of the polypeptide,  
and
- b) recovering the polypeptide so expressed.

20          33. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 28, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

## SEQUENCE LISTING

<110> INCYTE GENOMICS, INC.  
 LAL, Preeti  
 YANG, Junming  
 YUE, Henry  
 HILLMAN, Jennifer L.  
 TANG, Y. Tom  
 BANDMAN, Olga  
 BURFORD, Neil  
 BAUGHN, Mariah R.  
 AZIMZAI, Yalda  
 LU, Dyung Aina M.  
 AU-YOUNG, Janice  
 PATTERSON, Chandra

<120> HUMAN TRANSPORT PROTEINS

<130> PF-0709 PCT

<140> To Be Assigned  
 <141> Herewith

<150> 60/139,923; 60/148,177; 60/149,357; 60/162,287  
 <151> 1999-06-17; 1999-08-10; 1999-08-18; 1999-10-28

<160> 86  
 <170> PERL Program

<210> 1  
 <211> 623  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 264114CD1

<400> 1  
 Met Ser Thr Gln Asp Glu Arg Gln Ile Asn Thr Glu Tyr Ala Val  
 1 5 10 15  
 Ser Leu Leu Glu Gln Leu Lys Leu Phe Tyr Glu Gln Gln Leu Phe  
 20 25 30  
 Thr Asp Ile Val Leu Ile Val Glu Gly Thr Glu Phe Pro Cys His  
 35 40 45  
 Lys Met Val Leu Ala Thr Cys Ser Ser Tyr Phe Arg Ala Met Phe  
 50 55 60  
 Met Ser Gly Leu Ser Glu Ser Lys Gln Thr His Val His Leu Arg  
 65 70 75  
 Asn Val Asp Ala Ala Thr Leu Gln Ile Ile Ile Thr Tyr Ala Tyr  
 80 85 90  
 Thr Gly Asn Leu Ala Met Asn Asp Ser Thr Val Glu Gln Leu Tyr  
 95 100 105  
 Glu Thr Ala Cys Phe Leu Gln Val Glu Asp Val Leu Gln Arg Cys  
 110 115 120  
 Arg Glu Tyr Leu Ile Lys Lys Ile Asn Ala Glu Asn Cys Val Arg  
 125 130 135  
 Leu Leu Ser Phe Ala Asp Leu Phe Ser Cys Glu Glu Leu Lys Gln  
 140 145 150  
 Ser Ala Lys Arg Met Val Glu His Lys Phe Thr Ala Val Tyr His  
 155 160 165  
 Gln Asp Ala Phe Met Gln Leu Ser His Asp Leu Leu Ile Asp Ile  
 170 175 180  
 Leu Ser Ser Asp Asn Leu Asn Val Glu Lys Glu Glu Thr Val Arg  
 185 190 195  
 Glu Ala Ala Met Leu Trp Leu Glu Tyr Asn Thr Glu Ser Arg Ser

Gln Tyr Leu Ser	200	Ser Val Leu Ser Gln	205	Ile Arg Ile Asp Ala	210
Ser Glu Val Thr	215	Gln Arg Ala Trp Phe	220	Gln Gly Leu Pro Pro	225
Asp Lys Ser Val	230	Val Val Gln Gly Leu	235	Tyr Lys Ser Met Pro	240
Phe Phe Lys Pro	245	Arg Leu Gly Met Thr	250	Lys Glu Glu Met Met	255
Phe Ile Glu Ala	260	Ser Ser Glu Asn Pro	265	Cys Ser Leu Tyr Ser	270
Val Cys Tyr Ser	275	Pro Gln Ala Glu Lys	280	Val Tyr Lys Leu Cys	285
Pro Pro Ala Asp	290	Leu His Lys Val Gly	295	Thr Val Val Thr Pro	300
Asn Asp Ile Tyr	305	Ile Ala Gly Gly Gln	310	Val Pro Leu Lys Asn	315
Lys Thr Asn His	320	Ser Lys Thr Ser Lys	325	Leu Gln Thr Ala Phe	330
Thr Val Asn Cys	335	Phe Tyr Trp Phe Asp	340	Ala Gln Gln Asn Thr	345
Phe Pro Lys Thr	350	Pro Met Leu Phe Val	355	Arg Ile Lys Pro Ser	360
Val Cys Cys Glu	365	Gly Tyr Ile Tyr Ala	370	Ile Gly Gly Asp Ser	375
Gly Gly Glu Leu	380	Asn Arg Arg Thr Val	385	Glu Arg Tyr Asp Thr	390
Lys Asp Glu Trp	395	Thr Met Val Ser Pro	400	Leu Pro Cys Ala Trp	405
Trp Ser Ala Ala	410	Val Val His Asp Cys	415	Ile Tyr Val Met Thr	420
Leu Asn Leu Met	425	Tyr Cys Tyr Phe Pro	430	Arg Ser Asp Ser Trp	435
Glu Met Ala Met	440	Arg Gln Thr Ser Arg	445	Ser Phe Ala Ser Ala	450
Ala Phe Gly Asp	455	Lys Ile Phe Tyr Ile	460	Gly Gly Leu His Ile	465
Thr Asn Ser Gly	470	Ile Arg Leu Pro Ser	475	Gly Thr Val Asp Gly	480
Ser Val Thr Val	485	Glu Ile Tyr Asp Val	490	Asn Lys Asn Glu Trp	495
Met Ala Ala Asn	500	Ile Pro Ala Lys Arg	505	Tyr Ser Asp Pro Cys	510
Arg Ala Val Val	515	Ile Ser Asn Ser Leu	520	Cys Val Phe Met Arg	525
Thr His Leu Asn	530	Glu Arg Ala Lys Tyr	535	Val Thr Tyr Gln Tyr	540
Leu Glu Leu Asp	545	Arg Trp Ser Leu Arg	550	Gln His Ile Ser Glu	555
Val Leu Trp Asp	560	Leu Gly Arg Asp Phe	565	Arg Cys Thr Val Gly	570
Leu Tyr Pro Ser	575	Cys Leu Glu Glu Ser	580	Pro Trp Lys Pro Pro	585
Tyr Leu Phe Ser	590	Thr Asp Gly Thr Glu	595	Glu Phe Glu Leu Asp	600
Glu Met Val Ala	605	Leu Pro Pro Val	610		615
	620				

&lt;210&gt; 2

&lt;211&gt; 99

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1455669CD1

&lt;400&gt; 2

Met Ala Ala Pro Ala Glu Pro Cys Ala Gly Gln Gly Val Trp Asn

```

1      5      10      15
Gln Thr Glu Pro Glu Pro Ala Ala Thr Ser Leu Leu Ser Leu Cys
20      25      30
Phe Leu Arg Thr Ala Gly Val Trp Val Pro Pro Met Tyr Leu Trp
35      40      45
Val Leu Gly Pro Ile Tyr Leu Leu Phe Ile His His His Gly Arg
50      55      60
Gly Tyr Leu Arg Met Ser Pro Leu Phe Lys Ala Lys Met Val Ala
65      70      75
Ala Ile Pro Gly Ser Leu Glu Pro Gly Asn Val Arg Gly Arg Gln
80      85      90
Gly Thr Gly Trp Asn Leu Val Lys Ser
95

<210> 3
<211> 374
<212> PRT
<213> Homo sapiens

<220>
<221> misc_feature
<223> Incyte ID No: 2084989CD1

<400> 3
Met Glu Ser Lys Met Gly Glu Leu Pro Leu Asp Ile Asn Ile Gln
1      5      10      15
Glu Pro Arg Trp Asp Gln Ser Thr Phe Leu Gly Arg Ala Arg His
20      25      30
Phe Phe Thr Val Thr Asp Pro Arg Asn Leu Leu Leu Ser Gly Ala
35      40      45
Gln Leu Glu Ala Ser Arg Asn Ile Val Gln Asn Tyr Arg Ala Gly
50      55      60
Val Val Thr Pro Gly Ile Thr Glu Asp Gln Leu Trp Arg Ala Lys
65      70      75
Tyr Val Tyr Asp Ser Ala Phe His Pro Asp Thr Gly Glu Lys Val
80      85      90
Val Leu Ile Gly Arg Met Ser Ala Gln Val Pro Met Asn Met Thr
95      100      105
Ile Thr Gly Cys Met Leu Thr Phe Tyr Arg Lys Thr Pro Thr Val
110      115      120
Val Phe Trp Gln Trp Val Asn Gln Ser Phe Asn Ala Ile Val Asn
125      130      135
Tyr Ser Asn Arg Ser Gly Asp Thr Pro Ile Thr Val Arg Gln Leu
140      145      150
Gly Thr Ala Tyr Val Ser Ala Thr Thr Gly Ala Val Ala Thr Ala
155      160      165
Leu Gly Leu Lys Ser Leu Thr Lys His Leu Pro Pro Leu Val Gly
170      175      180
Arg Phe Val Pro Phe Ala Ala Val Ala Ala Ala Asn Cys Ile Asn
185      190      195
Ile Pro Leu Met Arg Gln Arg Glu Leu Gln Val Gly Ile Pro Val
200      205      210
Ala Asp Glu Ala Gly Gln Arg Leu Gly Tyr Ser Val Thr Ala Ala
215      220      225
Lys Gln Gly Ile Phe Gln Val Val Ile Ser Arg Ile Cys Met Ala
230      235      240
Ile Pro Ala Met Ala Ile Pro Pro Leu Ile Met Asp Thr Leu Glu
245      250      255
Lys Lys Asp Phe Leu Lys Val Gly Asp Cys Thr Ser Leu Val Leu
260      265      270
Glu Trp Ala Met Ala Gly Arg Ser Asp Gln Ala Pro Thr Leu Ser
275      280      285
Pro Ala Ser Pro Asp Ser Leu Arg Leu Ala Ser Pro Ser Pro Asp
290      295      300
Pro Cys Thr Ala Ser Ser Thr Phe Val His Ser Ala Arg Met Asn
305      310      315
Trp Ala Gly Val Lys Glu Leu Cys Arg Gly Arg Arg Arg Gly Gln
320      325      330
Arg Lys Glu Thr Asn Phe Ile Ser Val Thr Pro Val Ala Ser Asp

```



Thr Gln Lys Gly	335	Thr Val Ile Val Met	340	Leu Asp Leu Met Leu	345
	350		355		360
Leu Leu Pro Pro	365	Ser Ala Ser Ile Leu	Arg Gly Thr His Gly		

&lt;210&gt; 4

&lt;211&gt; 271

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2501034CD1

&lt;400&gt; 4

Met Gly Asn Gly Gly	Arg Ser Gly Leu Gln	Gln Gly Lys Gly Asn
1	5	10
Val Asp Gly Val	Ala Ala Thr Pro Thr	Ala Ala Ser Ala Ser Cys
	20	25
Gln Tyr Arg Cys	Ile Glu Cys Asn Gln	Glu Ala Lys Glu Leu Tyr
	35	40
Arg Asp Tyr Asn	His Gly Val Leu Lys	Ile Thr Ile Cys Lys Ser
	50	55
Cys Gln Lys Pro	Val Asp Lys Tyr Ile	Glu Tyr Asp Pro Val Ile
	65	70
Ile Leu Ile Asn	Ala Ile Leu Cys Lys	Ala Gln Ala Tyr Arg His
	80	85
Ile Leu Phe Asn	Thr Gln Ile Asn Ile	His Gly Lys Leu Cys Ile
	95	100
Phe Cys Leu Leu	Cys Glu Ala Tyr Leu	Arg Trp Trp Gln Leu Gln
	110	115
Asp Ser Asn Gln	Asn Thr Ala Pro Asp	Asp Leu Ile Arg Tyr Ala
	125	130
Lys Glu Trp Asp	Phe Tyr Arg Met Phe	Ala Ile Ala Ala Leu Glu
	140	145
Gln Thr Ala Tyr	Phe Ile Gly Ile Phe	Thr Phe Leu Trp Val Glu
	155	160
Arg Pro Met Thr	Ala Lys Lys Lys Pro	Asn Phe Ile Leu Leu Leu
	170	175
Lys Ala Leu Leu	Leu Ser Ser Tyr Gly	Lys Leu Leu Leu Ile Pro
	185	190
Ala Val Ile Trp	Glu His Asp Tyr Thr	Ser Val Cys Leu Lys Leu
	200	205
Ile Lys Val Phe	Val Leu Thr Ser Asn	Phe Gln Ala Ile Arg Val
	215	220
Thr Leu Asn Ile	Asn Arg Lys Leu Ser	Phe Leu Ala Val Leu Ser
	230	235
Gly Leu Leu Leu	Glu Ser Ile Met Val	Tyr Phe Phe Gln Ser Met
	245	250
Glu Trp Asp Val	Gly Ser Asp Tyr Ala	Ile Phe Lys Ser Gln Asp
	260	265
Phe		270

&lt;210&gt; 5

&lt;211&gt; 323

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2745212CD1

&lt;400&gt; 5

Met Ala Pro Lys	Gln Asp Pro Lys	Pro Lys Phe Gln Glu Gly Glu
1	5	10
Arg Val Leu Cys	Phe His Gly Pro Leu	Leu Tyr Glu Ala Lys Cys
	20	25
Val Lys Val Ala	Ile Lys Asp Lys Gln	Val Lys Tyr Phe Ile His

```

35      40      45
Tyr Ser Gly Trp Asn Lys Asn Trp Asp Glu Trp Val Pro Glu Ser
50      55      60
Arg Val Leu Lys Tyr Val Asp Thr Asn Leu Gln Lys Gln Arg Glu
65      70      75
Leu Gln Lys Ala Asn Gln Glu Gln Tyr Ala Glu Gly Lys Met Arg
80      85      90
Gly Ala Ala Pro Gly Lys Lys Thr Ser Gly Leu Gln Gln Lys Asn
95      100     105
Val Glu Val Lys Thr Lys Lys Asn Lys Gln Lys Thr Pro Gly Asn
110     115     120
Gly Asp Gly Gly Ser Thr Ser Glu Thr Pro Gln Pro Pro Arg Lys
125     130     135
Lys Arg Ala Arg Val Asp Pro Thr Val Glu Asn Glu Glu Thr Phe
140     145     150
Met Asn Arg Val Glu Val Lys Val Lys Ile Pro Glu Glu Leu Lys
155     160     165
Pro Trp Leu Val Asp Asp Trp Asp Leu Ile Thr Arg Gln Lys Gln
170     175     180
Leu Phe Tyr Leu Pro Ala Lys Lys Asn Val Asp Ser Ile Leu Glu
185     190     195
Asp Tyr Ala Asn Tyr Lys Lys Ser Arg Gly Asn Thr Asp Asn Lys
200     205     210
Glu Tyr Ala Val Asn Glu Val Val Ala Gly Ile Lys Glu Tyr Phe
215     220     225
Asn Val Met Leu Gly Thr Gln Leu Leu Tyr Lys Phe Glu Arg Pro
230     235     240
Gln Tyr Ala Glu Ile Leu Ala Asp His Pro Asp Ala Pro Met Ser
245     250     255
Gln Val Tyr Gly Ala Pro His Leu Leu Arg Leu Phe Val Arg Ile
260     265     270
Gly Ala Met Leu Ala Tyr Thr Pro Leu Asp Glu Lys Ser Leu Ala
275     280     285
Leu Leu Leu Asn Tyr Leu His Asp Phe Leu Lys Tyr Leu Ala Lys
290     295     300
Asn Ser Ala Thr Leu Phe Ser Ala Ser Asp Tyr Glu Val Ala Pro
305     310     315
Pro Glu Tyr His Arg Lys Ala Val
320

```

&lt;210&gt; 6

&lt;211&gt; 274

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4833111CD1

&lt;400&gt; 6

```

Met Asp Arg Pro Gly Phe Val Ala Ala Leu Val Ala Gly Gly Val
1      5      10      15
Ala Gly Val Ser Val Asp Leu Ile Leu Phe Pro Leu Asp Thr Ile
20     25     30
Lys Thr Arg Leu Gln Ser Pro Gln Gly Phe Ser Lys Ala Gly Gly
35     40     45
Phe His Gly Ile Tyr Ala Gly Val Pro Ser Ala Ala Ile Gly Ser
50     55     60
Phe Pro Asn Ala Ala Ala Phe Phe Ile Thr Tyr Glu Tyr Val Lys
65     70     75
Trp Phe Leu His Ala Asp Ser Ser Ser Tyr Leu Thr Pro Met Lys
80     85     90
His Met Leu Ala Ala Ser Ala Gly Glu Val Val Ala Cys Leu Ile
95     100    105
Arg Val Pro Ser Glu Val Val Lys Gln Arg Ala Gln Val Ser Ala
110    115    120
Ser Thr Arg Thr Phe Gln Ile Phe Ser Asn Ile Leu Tyr Glu Glu
125    130    135
Gly Ile Gln Gly Leu Tyr Arg Gly Tyr Lys Ser Thr Val Leu Arg

```

Glu Ile Pro Phe	140	Ser Leu Val Gln Phe	145	Pro Leu Trp Glu Ser	150
	155		160		165
Lys Ala Leu Trp	170	Ser Trp Arg Gln Asp	175	His Val Val Asp Ser	180
	185		190		195
Gln Ser Ala Val	200	Cys Gly Ala Phe Ala	205	Gly Gly Phe Ala Ala	210
	215		220		225
Val Thr Thr Pro	230	Leu Asp Val Ala Lys	235	Thr Arg Ile Thr Leu	240
	245		250		255
Lys Ala Gly Ser	260	Ser Thr Ala Asp Gly	265	Asn Val Leu Ser Val	270
His Gly Val Trp		Arg Ser Gln Gly Leu		Ala Gly Leu Phe Ala	
Val Phe Pro Arg		Met Ala Ala Ile Ser		Leu Gly Gly Phe Ile	
Leu Gly Ala Tyr		Asp Arg Thr His Ser		Leu Leu Leu Glu Val	
Arg Lys Ser Pro					

&lt;210&gt; 7

&lt;211&gt; 291

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 876677CD1

&lt;400&gt; 7

Met Asp Ser Arg Val	1	Ser Ser Pro Glu Lys	10	Gln Asp Lys Glu Asn	15
	5		25		30
Phe Val Gly Val Asn	20	Asn Lys Arg Leu Gly	25	Val Cys Gly Trp Ile	30
	35		40		45
Leu Phe Ser Leu Ser	50	Phe Leu Leu Val Ile	55	Ile Thr Phe Pro Ile	60
	65		70		75
Ser Ile Trp Met Cys	80	Leu Lys Ile Ile Lys	85	Glu Tyr Glu Arg Ala	90
	95		100		105
Val Val Phe Arg Leu	110	Gly Arg Ile Gln Ala	115	Asp Lys Ala Lys Gly	120
	125		130		135
Pro Gly Leu Ile Leu	140	Val Leu Pro Cys Ile	145	Asp Val Phe Val Lys	150
	155		160		165
Val Asp Leu Arg Thr	170	Val Thr Cys Asn Ile	175	Pro Pro Gln Glu Ile	180
	185		190		195
Leu Thr Arg Asp Ser	200	Val Thr Thr Gln Val	205	Asp Gly Val Val Tyr	210
	215		220		225
Tyr Arg Ile Tyr Ser	230	Ala Val Ser Ala Val	235	Ala Asn Val Asn Asp	240
	245		250		255
Val His Gln Ala Thr	260	Phe Leu Leu Ala Gln	265	Thr Thr Leu Arg Asn	270
	275		280		285
Val Leu Gly Thr Gln		Thr Leu Ser Gln Ile		Leu Ala Gly Arg Glu	
Glu Ile Ala His Ser		Ile Gln Thr Leu Leu		Asp Asp Ala Thr Glu	
Leu Trp Gly Ile Arg		Val Ala Arg Val Glu		Ile Lys Asp Val Arg	
Ile Pro Val Gln Leu		Gln Arg Ser Met Ala		Ala Glu Ala Glu Ala	
Thr Arg Glu Ala Arg		Ala Lys Val Leu Ala		Ala Glu Gly Glu Met	
Asn Ala Ser Lys Ser		Leu Lys Ser Ala Ser		Met Val Leu Ala Glu	
Ser Pro Ile Ala Leu		Gln Leu Arg Tyr Leu		Gln Thr Leu Ser Thr	
Val Ala Thr Glu Lys		Asn Ser Thr Ile Val		Phe Pro Leu Pro Met	
Asn Ile Leu Glu Gly		Ile Gly Gly Val Ser		Tyr Asp Asn His Lys	
Lys Leu Pro Asn Lys					

290

<210> 8  
 <211> 381  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2326143CD1

<400> 8  
 Met Ser Arg His Glu Gly Val Ser Cys Asp Ala Cys Leu Lys Gly  
 1 5 10 15  
 Asn Phe Arg Gly Arg Tyr Lys Cys Leu Ile Cys Tyr Asp Tyr  
 20 25 30  
 Asp Leu Cys Ala Ser Cys Tyr Glu Ser Gly Ala Thr Thr Thr Arg  
 35 40 45  
 His Thr Thr Asp His Pro Met Gln Cys Ile Leu Thr Arg Val Asp  
 50 55 60  
 Phe Asp Leu Tyr Tyr Gly Gly Glu Ala Phe Ser Val Glu Gln Pro  
 65 70 75  
 Gln Ser Phe Thr Cys Pro Tyr Cys Gly Lys Met Gly Tyr Thr Glu  
 80 85 90  
 Thr Ser Leu Gln Glu His Val Thr Ser Glu His Ala Glu Thr Ser  
 95 100 105  
 Thr Glu Val Ile Cys Pro Ile Cys Ala Ala Leu Pro Gly Gly Asp  
 110 115 120  
 Pro Asn His Val Thr Asp Asp Phe Ala Ala His Leu Thr Leu Glu  
 125 130 135  
 His Arg Ala Pro Arg Asp Leu Asp Glu Ser Ser Gly Val Arg His  
 140 145 150  
 Val Arg Arg Met Phe His Pro Gly Arg Gly Leu Gly Gly Pro Arg  
 155 160 165  
 Ala Arg Arg Ser Asn Met His Phe Thr Ser Ser Ser Thr Gly Gly  
 170 175 180  
 Leu Ser Ser Ser Gln Ser Ser Tyr Ser Pro Ser Asn Arg Glu Ala  
 185 190 195  
 Met Asp Pro Ile Ala Glu Leu Leu Ser Gln Leu Ser Gly Val Arg  
 200 205 210  
 Arg Ser Ala Gly Gly Gln Leu Asn Ser Ser Gly Pro Ser Ala Ser  
 215 220 225  
 Gln Leu Gln Gln Leu Gln Met Gln Leu Gln Leu Glu Arg Gln His  
 230 235 240  
 Ala Gln Ala Ala Arg Gln Gln Leu Glu Thr Ala Arg Asn Ala Thr  
 245 250 255  
 Arg Arg Thr Asn Thr Ser Ser Val Thr Thr Thr Ile Thr Gln Ser  
 260 265 270  
 Thr Ala Thr Thr Asn Ile Ala Asn Thr Glu Ser Ser Gln Gln Thr  
 275 280 285  
 Leu Gln Asn Ser Gln Phe Leu Leu Thr Arg Leu Asn Asp Pro Lys  
 290 295 300  
 Met Ser Glu Thr Glu Arg Gln Ser Met Glu Ser Glu Arg Ala Asp  
 305 310 315  
 Arg Ser Leu Phe Val Gln Glu Leu Leu Leu Ser Thr Leu Val Arg  
 320 325 330  
 Glu Glu Ser Ser Ser Ser Asp Glu Asp Asp Arg Gly Glu Met Ala  
 335 340 345  
 Asp Phe Gly Ala Met Gly Cys Val Asp Ile Met Pro Leu Asp Val  
 350 355 360  
 Ala Leu Glu Asn Leu Asn Leu Lys Glu Ser Asn Lys Gly Asn Glu  
 365 370 375  
 Pro Pro Pro Pro Pro Leu  
 380

<210> 9  
 <211> 190  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2786302CD1

<400> 9

Met	Lys	Tyr	Gly	Asn	Glu	Ile	Met	Asn	Lys	Asp	Pro	Val	Phe	Arg	
1				5					10					15	
Ile	Ser	Pro	Arg	Ser	Arg	Glu	Thr	His	Pro	Asn	Pro	Glu	Glu	Pro	
				20					25					30	
Glu	Glu	Glu	Asp	Glu	Asp	Val	Gln	Ala	Glu	Arg	Val	Gln	Ala	Ala	
				35					40					45	
Asn	Ala	Leu	Thr	Ala	Pro	Asn	Leu	Glu	Glu	Glu	Pro	Val	Ile	Thr	
				50					55					60	
Ala	Ser	Cys	Leu	His	Lys	Glu	Tyr	Tyr	Glu	Thr	Lys	Lys	Ser	Cys	
				65					70					75	
Phe	Ser	Thr	Arg	Lys	Lys	Lys	Ile	Ala	Ile	Arg	Asn	Val	Ser	Phe	
				80					85					90	
Cys	Val	Lys	Lys	Gly	Glu	Val	Leu	Gly	Leu	Leu	Gly	His	Asn	Gly	
				95					100					105	
Ala	Gly	Lys	Ser	Thr	Ser	Ile	Lys	Met	Ile	Thr	Gly	Cys	Thr	Lys	
				110					115					120	
Pro	Thr	Ala	Gly	Val	Val	Val	Leu	Gln	Gly	Ser	Arg	Ala	Ser	Val	
				125					130					135	
Arg	Gln	Gln	His	Asp	Asn	Ser	Leu	Lys	Phe	Leu	Gly	Tyr	Cys	Pro	
				140					145					150	
Gln	Glu	Asn	Ser	Leu	Trp	Pro	Lys	Leu	Thr	Met	Lys	Glu	His	Leu	
				155					160					165	
Glu	Leu	Tyr	Ala	Ala	Val	Glu	Arg	Leu	Gly	Gln	Lys	Arg	Cys	Cys	
				170					175					180	
Ser	Gln	Tyr	Phe	Thr	Ile	Gly	Gly	Arg	Ser						
				185					190						

<210> 10  
 <211> 297  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3735780CD1

<400> 10

Met	Met	Asp	Ser	Glu	Ala	His	Glu	Lys	Arg	Pro	Pro	Ile	Leu	Thr	
1				5					10					15	
Ser	Ser	Lys	Gln	Asp	Ile	Ser	Pro	His	Ile	Thr	Asn	Val	Gly	Glu	
				20					25					30	
Met	Lys	His	Tyr	Leu	Cys	Gly	Cys	Cys	Ala	Ala	Phe	Asn	Asn	Val	
				35					40					45	
Ala	Ile	Thr	Phe	Pro	Ile	Gln	Lys	Val	Leu	Phe	Arg	Gln	Gln	Leu	
				50					55					60	
Tyr	Gly	Ile	Lys	Thr	Arg	Asp	Ala	Ile	Leu	Gln	Leu	Arg	Arg	Asp	
				65					70					75	
Gly	Phe	Arg	Asn	Leu	Tyr	Arg	Gly	Ile	Leu	Pro	Pro	Leu	Met	Gln	
				80					85					90	
Lys	Thr	Thr	Thr	Leu	Ala	Leu	Met	Phe	Gly	Leu	Tyr	Glu	Asp	Leu	
				95					100					105	
Ser	Cys	Leu	Leu	His	Lys	His	Val	Ser	Ala	Pro	Glu	Phe	Ala	Thr	
				110					115					120	
Ser	Gly	Val	Ala	Ala	Val	Leu	Ala	Gly	Thr	Thr	Glu	Ala	Ile	Phe	
				125					130					135	
Thr	Pro	Leu	Glu	Arg	Val	Gln	Thr	Leu	Leu	Gln	Asp	His	Lys	His	
				140					145					150	
His	Asp	Lys	Phe	Thr	Asn	Thr	Tyr	Gln	Ala	Phe	Lys	Ala	Leu	Lys	
				155					160					165	
Cys	His	Gly	Ile	Gly	Glu	Tyr	Tyr	Arg	Gly	Leu	Val	Pro	Ile	Leu	
				170					175					180	
Phe	Arg	Asn	Gly	Leu	Ser	Asn	Val	Leu	Phe	Phe	Gly	Leu	Arg	Gly	
				185					190					195	
Pro	Ile	Lys	Glu	His	Leu	Pro	Thr	Ala	Thr	Thr	His	Ser	Ala	His	

200	205	210
Leu Val Asn Asp Phe Ile Cys Gly Gly	Leu Leu Gly Ala Met Leu	
215	220	225
Gly Phe Leu Phe Phe Pro Ile Asn Val	Val Lys Thr Arg Ile Gln	
230	235	240
Ser Gln Ile Gly Gly Glu Phe Gln Ser	Phe Pro Lys Val Phe Gln	
245	250	255
Lys Ile Trp Leu Glu Arg Asp Arg Lys	Leu Ile Asn Leu Phe Arg	
260	265	270
Gly Ala His Leu Asn Tyr His Arg Ser	Leu Ile Ser Trp Gly Ile	
275	280	285
Ile Asn Ala Thr Tyr Glu Phe Leu Leu	Lys Val Ile	
290	295	

<210> 11  
 <211> 89  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 039026CD1

<400> 11	
Met Ala Ala Gln Ile Pro Glu Ser Asp Gln Ile Lys Gln Phe Lys	
1 5 10 15	
Glu Phe Leu Gly Thr Tyr Asn Lys Leu Thr Glu Thr Cys Phe Leu	
20 25 30	
Asp Cys Val Lys Asp Phe Thr Thr Arg Glu Val Lys Pro Glu Glu	
35 40 45	
Thr Thr Cys Ser Glu His Cys Leu Gln Lys Tyr Leu Lys Met Thr	
50 55 60	
Gln Arg Ile Ser Met Arg Phe Gln Glu Tyr His Ile Gln Gln Asn	
65 70 75	
Glu Ala Leu Ala Ala Lys Ala Gly Leu Leu Gly Gln Pro Arg	
80 85	

<210> 12  
 <211> 115  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 260607CD1

<400> 12	
Met Ala Leu Ile Pro Ser Arg Val Trp Leu Pro Phe Ala Val Trp	
1 5 10 15	
Val Val Asp Ser Ala Pro Val Arg Gly Leu Val Arg Arg Glu Pro	
20 25 30	
Phe Leu Arg Thr Gly Ser Phe Ile Ala Leu Phe Tyr Phe Pro Pro	
35 40 45	
Leu Leu Pro Val Leu Ile Asn Leu Phe Ser Phe Phe Leu Thr Pro	
50 55 60	
Ser Phe Trp Arg Gln Leu Gly Ala Ile Leu Val Tyr Ala Ser Leu	
65 70 75	
Leu Ala Glu Lys Thr Pro Phe Lys Thr Gln Arg Thr Leu Glu Gly	
80 85 90	
Asp Ala Leu Val Gly Ser Val Ser Ile Phe Leu Cys Ala Lys Asp	
95 100 105	
Arg Gln Thr Glu Ala Glu Arg Gly Cys Ser	
110 115	

<210> 13  
 <211> 675  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature

&lt;223&gt; Incyte ID No: 1429651CD1

&lt;400&gt; 13

Met	Glu	Ser	Gly	Thr	Ser	Ser	Pro	Gln	Pro	Pro	Gln	Leu	Asp	Pro
1				5					10					15
Leu	Asp	Ala	Phe	Pro	Gln	Lys	Gly	Leu	Glu	Pro	Gly	Asp	Ile	Ala
				20					25					30
Val	Leu	Val	Leu	Tyr	Phe	Leu	Phe	Val	Leu	Ala	Val	Gly	Leu	Trp
				35					40					45
Ser	Thr	Val	Lys	Thr	Lys	Arg	Asp	Thr	Val	Lys	Gly	Tyr	Phe	Leu
				50					55					60
Ala	Gly	Gly	Asp	Met	Val	Trp	Trp	Pro	Val	Gly	Ala	Ser	Leu	Phe
				65					70					75
Ala	Ser	Asn	Val	Gly	Ser	Gly	His	Phe	Ile	Gly	Leu	Ala	Gly	Ser
				80					85					90
Gly	Ala	Ala	Thr	Gly	Ile	Ser	Val	Ser	Ala	Tyr	Glu	Leu	Asn	Gly
				95					100					105
Leu	Phe	Ser	Val	Leu	Met	Leu	Ala	Trp	Ile	Phe	Leu	Pro	Ile	Tyr
				110					115					120
Ile	Ala	Gly	Gln	Val	Thr	Thr	Met	Pro	Glu	Tyr	Leu	Arg	Lys	Arg
				125					130					135
Phe	Gly	Gly	Ile	Arg	Ile	Pro	Ile	Ile	Leu	Ala	Val	Leu	Tyr	Leu
				140					145					150
Phe	Ile	Tyr	Ile	Phe	Thr	Lys	Ile	Ser	Val	Asp	Met	Tyr	Ala	Gly
				155					160					165
Ala	Ile	Phe	Ile	Gln	Ser	Leu	His	Leu	Asp	Leu	Tyr	Leu	Ala	
				170					175					180
Ile	Val	Gly	Leu	Leu	Ala	Ile	Thr	Ala	Val	Tyr	Thr	Val	Ala	Gly
				185					190					195
Gly	Leu	Ala	Ala	Val	Ile	Tyr	Thr	Asp	Ala	Leu	Gln	Thr	Leu	Ile
				200					205					210
Met	Leu	Ile	Gly	Ala	Leu	Thr	Leu	Met	Gly	Tyr	Ser	Phe	Ala	Ala
				215					220					225
Val	Gly	Gly	Met	Glu	Gly	Leu	Lys	Glu	Lys	Tyr	Phe	Leu	Ala	Leu
				230					235					240
Ala	Ser	Asn	Arg	Ser	Glu	Asn	Ser	Ser	Cys	Gly	Leu	Pro	Arg	Glu
				245					250					255
Asp	Ala	Phe	His	Ile	Phe	Arg	Asp	Pro	Leu	Thr	Ser	Asp	Leu	Pro
				260					265					270
Trp	Pro	Gly	Val	Leu	Phe	Gly	Met	Ser	Ile	Pro	Ser	Leu	Trp	Tyr
				275					280					285
Trp	Cys	Thr	Asp	Gln	Val	Ile	Val	Gln	Arg	Thr	Leu	Ala	Ala	Lys
				290					295					300
Asn	Leu	Ser	His	Ala	Lys	Gly	Gly	Ala	Leu	Met	Ala	Ala	Tyr	Leu
				305					310					315
Lys	Val	Leu	Pro	Leu	Phe	Ile	Met	Val	Phe	Pro	Gly	Met	Val	Ser
				320					325					330
Arg	Ile	Leu	Phe	Pro	Asp	Gln	Val	Ala	Cys	Ala	Asp	Pro	Glu	Ile
				335					340					345
Cys	Gln	Lys	Ile	Cys	Ser	Asn	Pro	Ser	Gly	Cys	Ser	Asp	Ile	Ala
				350					355					360
Tyr	Pro	Lys	Leu	Val	Leu	Glu	Leu	Leu	Pro	Thr	Gly	Leu	Arg	Gly
				365					370					375
Leu	Met	Met	Ala	Val	Met	Val	Ala	Ala	Leu	Met	Ser	Ser	Leu	Thr
				380					385					390
Ser	Ile	Phe	Asn	Ser	Ala	Ser	Thr	Ile	Phe	Thr	Met	Asp	Leu	Trp
				395					400					405
Asn	His	Leu	Arg	Pro	Arg	Ala	Ser	Glu	Lys	Glu	Leu	Met	Ile	Val
				410					415					420
Gly	Arg	Val	Phe	Val	Leu	Leu	Leu	Val	Leu	Val	Ser	Ile	Leu	Trp
				425					430					435
Ile	Pro	Val	Val	Gln	Ala	Ser	Gln	Gly	Gly	Gln	Leu	Phe	Ile	Tyr
				440					445					450
Ile	Gln	Ser	Ile	Ser	Ser	Tyr	Leu	Gln	Pro	Pro	Val	Ala	Val	Val
				455					460					465
Phe	Ile	Met	Gly	Cys	Phe	Trp	Lys	Arg	Thr	Asn	Glu	Lys	Gly	Ala
				470					475					480
Phe	Trp	Gly	Leu	Ile	Ser	Gly	Leu	Leu	Leu	Gly	Leu	Val	Arg	Leu

	485		490		495
Val Leu Asp Phe	Ile Tyr Val Gln Pro	Arg Cys Asp Gln Pro	Asp		
	500		505		510
Glu Arg Pro Val	Leu Val Lys Ser Ile	His Tyr Leu Tyr Phe	Ser		
	515		520		525
Met Ile Leu Ser	Thr Val Thr Leu Ile	Thr Val Ser Thr Val	Ser		
	530		535		540
Trp Phe Thr Glu	Pro Pro Ser Lys Glu	Met Val Ser His Leu	Thr		
	545		550		555
Trp Phe Thr Arg	His Asp Pro Val Val	Gln Lys Glu Gln Ala	Pro		
	560		565		570
Pro Ala Ala Pro	Leu Ser Leu Thr Leu	Ser Gln Asn Gly Met	Pro		
	575		580		585
Glu Ala Ser Ser	Ser Ser Ser Val Gln	Phe Glu Met Val Gln	Glu		
	590		595		600
Asn Thr Ser Lys	Thr His Ser Cys Asp	Met Thr Pro Lys Gln	Ser		
	605		610		615
Lys Val Val Lys	Ala Ile Leu Trp Leu	Cys Gly Ile Gln Glu	Lys		
	620		625		630
Gly Lys Glu Glu	Leu Pro Ala Arg Ala	Glu Ala Ile Ile Val	Ser		
	635		640		645
Leu Glu Glu Asn	Pro Leu Val Lys Thr	Leu Leu Asp Val Asn	Leu		
	650		655		660
Ile Phe Cys Val	Ser Cys Ala Ile Phe	Ile Trp Gly Tyr Phe	Ala		
	665		670		675

&lt;210&gt; 14

&lt;211&gt; 320

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2069971CD1

<400> 14	
Met Tyr His Cys His	Ser Gly Ser Lys Pro Thr Glu Lys Gly Ala
1	5
Asn Glu Tyr Ala Tyr	Ala Lys Trp Lys Leu Cys Ser Ala Ser Ala
	20
Ile Cys Phe Ile Phe	Met Ile Ala Glu Val Val Gly Gly His Ile
	35
Ala Gly Ser Leu Ala	Val Val Thr Asp Ala Ala His Leu Leu Ile
	50
Asp Leu Thr Ser Phe	Leu Leu Ser Leu Phe Ser Leu Trp Leu Ser
	65
Ser Lys Pro Pro Ser	Lys Arg Leu Thr Phe Gly Trp His Arg Ala
	80
Glu Ile Leu Gly Ala	Leu Leu Ser Ile Leu Cys Ile Trp Val Val
	95
Thr Gly Val Leu Val	Tyr Leu Ala Cys Glu Arg Leu Leu Tyr Pro
	110
Asp Tyr Gln Ile Gln	Ala Thr Val Met Ile Ile Val Ser Ser Cys
	125
Ala Val Ala Ala Asn	Ile Val Leu Thr Val Val Leu His Gln Arg
	140
Cys Leu Gly His Asn	His Lys Glu Val Gln Ala Asn Ala Ser Val
	155
Arg Ala Ala Phe Val	His Ala Leu Gly Asp Leu Phe Gln Ser Ile
	170
Ser Val Leu Ile Ser	Ala Leu Ile Ile Tyr Phe Lys Pro Glu Tyr
	185
Lys Ile Ala Asp Pro	Ile Cys Thr Phe Ile Phe Ser Ile Leu Val
	200
Leu Ala Ser Thr Ile	Thr Ile Leu Lys Asp Phe Ser Ile Leu Leu
	215
Met Glu Gly Val Pro	Lys Ser Leu Asn Tyr Ser Gly Val Lys Glu
	230
	235
	240



Leu	Ile	Leu	Ala	Val	Asp	Gly	Val	Leu	Ser	Val	His	Ser	Leu	His
				245					250					255
Ile	Trp	Ser	Leu	Thr	Met	Asn	Gln	Val	Ile	Leu	Ser	Ala	His	Val
				260					265					270
Ala	Thr	Ala	Ala	Ser	Arg	Asp	Ser	Gln	Val	Val	Arg	Arg	Glu	Ile
				275					280					285
Ala	Lys	Ala	Leu	Ser	Lys	Ser	Phe	Thr	Met	His	Ser	Leu	Thr	Ile
				290					295					300
Gln	Met	Glu	Ser	Pro	Val	Asp	Gln	Asp	Pro	Asp	Cys	Leu	Phe	Cys
				305					310					315
Glu	Asp	Pro	Cys	Asp										
				320										

&lt;210&gt; 15

&lt;211&gt; 462

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2329339CD1

&lt;400&gt; 15

Met	Ala	Glu	Glu	Gln	Glu	Phe	Thr	Gln	Leu	Cys	Lys	Leu	Pro	Ala
1				5					10					15
Gln	Pro	Ser	His	Pro	His	Cys	Val	Asn	Asn	Thr	Tyr	Arg	Ser	Ala
				20					25					30
Gln	His	Ser	Gln	Ala	Leu	Leu	Arg	Gly	Leu	Leu	Ala	Leu	Arg	Asp
				35					40					45
Ser	Gly	Ile	Leu	Phe	Asp	Val	Val	Leu	Val	Val	Glu	Gly	Arg	His
				50					55					60
Ile	Glu	Ala	His	Arg	Ile	Leu	Leu	Ala	Ala	Ser	Cys	Asp	Tyr	Phe
				65					70					75
Arg	Gly	Met	Phe	Ala	Gly	Gly	Leu	Lys	Glu	Met	Glu	Gln	Glu	Glu
				80					85					90
Val	Leu	Ile	His	Gly	Val	Ser	Tyr	Asn	Ala	Met	Cys	Gln	Ile	Leu
				95					100					105
His	Phe	Ile	Tyr	Thr	Ser	Glu	Leu	Glu	Leu	Ser	Leu	Ser	Asn	Val
				110					115					120
Gln	Glu	Thr	Leu	Val	Ala	Ala	Cys	Gln	Leu	Gln	Ile	Pro	Glu	Ile
				125					130					135
Ile	His	Phe	Cys	Cys	Asp	Phe	Leu	Met	Ser	Trp	Val	Asp	Glu	Glu
				140					145					150
Asn	Ile	Leu	Asp	Val	Tyr	Arg	Leu	Ala	Glu	Leu	Phe	Asp	Leu	Ser
				155					160					165
Arg	Leu	Thr	Glu	Gln	Leu	Asp	Thr	Tyr	Ile	Leu	Lys	Asn	Phe	Val
				170					175					180
Ala	Phe	Ser	Arg	Thr	Asp	Lys	Tyr	Arg	Gln	Leu	Pro	Leu	Glu	Lys
				185					190					195
Val	Tyr	Ser	Leu	Leu	Ser	Ser	Asn	Arg	Leu	Glu	Val	Ser	Cys	Glu
				200					205					210
Thr	Glu	Val	Tyr	Glu	Gly	Ala	Leu	Leu	Tyr	His	Tyr	Ser	Leu	Glu
				215					220					225
Gln	Val	Gln	Ala	Asp	Gln	Ile	Ser	Leu	His	Glu	Pro	Pro	Lys	Leu
				230					235					240
Leu	Glu	Thr	Val	Arg	Phe	Pro	Leu	Met	Glu	Ala	Glu	Val	Leu	Gln
				245					250					255
Arg	Leu	His	Asp	Lys	Leu	Asp	Pro	Ser	Pro	Leu	Arg	Asp	Thr	Val
				260					265					270
Ala	Ser	Gly	Leu	Met	Tyr	His	Arg	Asn	Glu	Ser	Leu	Gln	Pro	Ser
				275					280					285
Leu	Gln	Ser	Pro	Gln	Thr	Glu	Leu	Arg	Ser	Asp	Phe	Gln	Cys	Val
				290					295					300
Val	Gly	Phe	Gly	Gly	Ile	His	Ser	Thr	Pro	Ser	Thr	Val	Leu	Ser
				305					310					315
Asp	Gln	Ala	Lys	Tyr	Leu	Asn	Pro	Leu	Leu	Gly	Glu	Trp	Lys	His
				320					325					330
Phe	Thr	Ala	Ser	Leu	Ala	Pro	Arg	Met	Ser	Asn	Gln	Gly	Ile	Ala
				335					340					345

```

Val Leu Asn Asn Phe Val Tyr Leu Ile Gly Gly Asp Asn Asn Val
350 355 360
Gln Gly Phe Arg Ala Glu Ser Arg Cys Trp Arg Tyr Asp Pro Arg
365 370 375
His Asn Arg Trp Phe Gln Ile Gln Ser Leu Gln Gln Glu His Ala
380 385 390
Asp Leu Ser Val Cys Val Val Gly Arg Tyr Ile Tyr Ala Val Ala
395 400 405
Gly Arg Asp Tyr His Asn Asp Leu Asn Ala Val Glu Arg Tyr Asp
410 415 420
Pro Ala Thr Asn Ser Trp Ala Tyr Val Ala Pro Leu Lys Arg Glu
425 430 435
Val Tyr Ala His Ala Gly Ala Thr Leu Glu Gly Lys Met Tyr Ile
440 445 450
Thr Cys Gly Arg Lys Leu Ile Pro Phe Ser Glu Gly
455 460

```

<210> 16  
 <211> 98  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2540219CD1

```

<400> 16
Met Arg Ala Cys Ala Val Trp Leu Ala Gly Gly Met Ala Gly Ala
1 5 10 15
Ile Ser Trp Gly Thr Ala Thr Pro Met Asp Val Val Lys Ser Arg
20 25 30
Leu Gln Ala Asp Gly Val Tyr Leu Asn Lys Tyr Lys Gly Val Leu
35 40 45
Asp Cys Ile Ser Gln Ser Tyr Gln Lys Glu Gly Leu Lys Val Phe
50 55 60
Phe Arg Gly Ile Thr Val Asn Ala Val Arg Gly Phe Pro Met Ser
65 70 75
Ala Ala Met Phe Leu Gly Tyr Glu Leu Ser Leu Gln Ala Ile Arg
80 85 90
Gly Asp His Ala Val Thr Ser Pro
95

```

<210> 17  
 <211> 748  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2722462CD1

```

<400> 17
Met Asn Tyr Gln Glu Ala Ala Ile Tyr Leu Gln Glu Gly Glu Asn
1 5 10 15
Asn Asp Lys Phe Phe Thr His Pro Lys Asp Ala Lys Ala Leu Ala
20 25 30
Ala Tyr Leu Phe Ala His Asn His Leu Phe Tyr Leu Met Glu Leu
35 40 45
Ala Thr Ala Leu Leu Leu Leu Leu Ser Leu Cys Glu Ala Pro
50 55 60
Ala Val Pro Ala Leu Arg Leu Gly Ile Tyr Val His Ala Thr Leu
65 70 75
Glu Leu Phe Ala Leu Met Val Val Val Phe Glu Leu Cys Met Lys
80 85 90
Leu Arg Trp Leu Gly Leu His Thr Phe Ile Arg His Lys Arg Thr
95 100 105
Met Val Lys Thr Ser Val Leu Val Val Gln Phe Val Glu Ala Ile
110 115 120
Val Val Leu Val Arg Gln Met Ser His Val Arg Val Thr Arg Ala
125 130 135

```

Leu	Arg	Cys	Ile	Phe	Leu	Val	Asp	Cys	Arg	Tyr	Cys	Gly	Gly	Val
				140					145					150
Arg	Arg	Asn	Leu	Arg	Gln	Ile	Phe	Gln	Ser	Leu	Pro	Pro	Phe	Met
				155					160					165
Asp	Ile	Leu	Leu	Leu	Leu	Phe	Phe	Phe	Met	Ile	Ile	Phe	Ala	Ile
				170					175					180
Leu	Gly	Phe	Tyr	Leu	Phe	Ser	Pro	Asn	Pro	Ser	Asp	Pro	Tyr	Phe
				185					190					195
Ser	Thr	Leu	Glu	Asn	Ser	Ile	Val	Ser	Leu	Phe	Val	Leu	Leu	Thr
				200					205					210
Thr	Ala	Asn	Phe	Pro	Asp	Val	Met	Met	Pro	Ser	Tyr	Ser	Arg	Asn
				215					220					225
Pro	Trp	Ser	Cys	Val	Phe	Phe	Ile	Val	Tyr	Leu	Ser	Ile	Glu	Leu
				230					235					240
Tyr	Phe	Ile	Met	Asn	Leu	Leu	Leu	Ala	Val	Val	Phe	Asp	Thr	Phe
				245					250					255
Asn	Asp	Ile	Glu	Lys	Arg	Lys	Phe	Lys	Ser	Leu	Leu	Leu	His	Lys
				260					265					270
Arg	Thr	Ala	Ile	Gln	His	Ala	Tyr	Arg	Leu	Leu	Ile	Ser	Gln	Arg
				275					280					285
Arg	Pro	Ala	Gly	Ile	Ser	Tyr	Arg	Gln	Phe	Glu	Gly	Leu	Met	Arg
				290					295					300
Phe	Tyr	Lys	Pro	Arg	Met	Ser	Ala	Arg	Glu	Arg	Tyr	Leu	Thr	Phe
				305					310					315
Lys	Ala	Leu	Asn	Gln	Asn	Asn	Thr	Pro	Leu	Leu	Ser	Leu	Lys	Asp
				320					325					330
Phe	Tyr	Asp	Ile	Tyr	Glu	Val	Ala	Ala	Leu	Lys	Trp	Lys	Ala	Lys
				335					340					345
Lys	Asn	Arg	Glu	His	Trp	Phe	Asp	Glu	Leu	Pro	Arg	Thr	Ala	Leu
				350					355					360
Leu	Ile	Phe	Lys	Gly	Ile	Asn	Ile	Leu	Val	Lys	Ser	Lys	Ala	Phe
				365					370					375
Gln	Tyr	Phe	Met	Tyr	Leu	Val	Val	Ala	Val	Asn	Gly	Val	Trp	Ile
				380					385					390
Leu	Val	Glu	Thr	Phe	Met	Leu	Lys	Gly	Gly	Asn	Phe	Phe	Ser	Lys
				395					400					405
His	Val	Pro	Trp	Ser	Tyr	Leu	Val	Phe	Leu	Thr	Ile	Tyr	Gly	Val
				410					415					420
Glu	Leu	Phe	Leu	Lys	Val	Ala	Gly	Leu	Gly	Pro	Val	Glu	Tyr	Leu
				425					430					435
Ser	Ser	Gly	Trp	Asn	Leu	Phe	Asp	Phe	Ser	Val	Thr	Val	Phe	Ala
				440					445					450
Phe	Leu	Gly	Leu	Leu	Ala	Leu	Ala	Leu	Asn	Met	Glu	Pro	Phe	Tyr
				455					460					465
Phe	Ile	Val	Val	Leu	Arg	Pro	Leu	Gln	Leu	Leu	Arg	Leu	Phe	Lys
				470					475					480
Leu	Lys	Glu	Arg	Tyr	Arg	Asn	Val	Leu	Asp	Thr	Met	Phe	Glu	Leu
				485					490					495
Leu	Pro	Arg	Met	Ala	Ser	Leu	Gly	Leu	Thr	Leu	Leu	Ile	Phe	Tyr
				500					505					510
Tyr	Ser	Phe	Ala	Ile	Val	Gly	Met	Glu	Phe	Phe	Cys	Gly	Ile	Val
				515					520					525
Phe	Pro	Asn	Cys	Cys	Asn	Thr	Ser	Thr	Val	Ala	Asp	Ala	Tyr	Arg
				530					535					540
Trp	Arg	Asn	His	Thr	Val	Gly	Asn	Arg	Thr	Val	Val	Glu	Glu	Gly
				545					550					555
Tyr	Tyr	Tyr	Leu	Asn	Asn	Phe	Asp	Asn	Ile	Leu	Asn	Ser	Phe	Val
				560					565					570
Thr	Leu	Phe	Glu	Leu	Thr	Val	Val	Asn	Asn	Trp	Tyr	Ile	Ile	Met
				575					580					585
Glu	Gly	Val	Thr	Ser	Gln	Thr	Ser	His	Trp	Ser	Arg	Leu	Tyr	Phe
				590					595					600
Met	Thr	Phe	Tyr	Ile	Val	Thr	Met	Val	Val	Met	Thr	Ile	Ile	Val
				605					610					615
Ala	Phe	Ile	Leu	Glu	Ala	Phe	Val	Phe	Arg	Met	Asn	Tyr	Ser	Arg
				620					625					630
Lys	Asn	Gln	Asp	Ser	Glu	Val	Asp	Gly	Gly	Ile	Thr	Leu	Glu	Lys
				635					640					645

Glu Ile Ser Lys Glu Glu Leu Val Ala Val Leu Glu Leu Tyr Arg  
 650 655 660  
 Glu Ala Arg Gly Ala Ser Ser Asp Val Thr Arg Leu Leu Glu Thr  
 665 670 675  
 Leu Ser Gln Met Glu Arg Tyr Gln Gln His Ser Met Val Phe Leu  
 680 685 690  
 Gly Arg Arg Ser Arg Thr Lys Ser Asp Leu Ser Leu Lys Met Tyr  
 695 700 705  
 Gln Glu Glu Ile Gln Glu Trp Tyr Glu Glu His Ala Arg Glu Gln  
 710 715 720  
 Glu Gln Gln Arg Gln Leu Ser Ser Ser Ala Ala Pro Ala Ala Gln  
 725 730 735  
 Gln Pro Pro Gly Ser Arg Gln Arg Ser Gln Thr Val Thr  
 740 745

&lt;210&gt; 18

&lt;211&gt; 507

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2739264CD1

&lt;400&gt; 18

Met Ala Phe Asn Phe Gly Ala Pro Ser Gly Thr Ser Gly Thr Ala  
 1 5 10 15  
 Ala Ala Thr Ala Ala Pro Ala Gly Gly Phe Gly Gly Phe Gly Thr  
 20 25 30  
 Thr Ser Thr Thr Thr Gly Ser Ala Phe Ser Phe Ser Ala Pro Thr  
 35 40 45  
 Asn Thr Gly Thr Thr Gly Leu Phe Gly Gly Thr Gln Asn Lys Gly  
 50 55 60  
 Phe Gly Phe Gly Thr Gly Phe Gly Thr Thr Thr Gly Thr Ser Thr  
 65 70 75  
 Gly Leu Gly Thr Gly Leu Gly Thr Gly Leu Gly Phe Gly Gly Phe  
 80 85 90  
 Asn Thr Gln Gln Gln Gln Gln Thr Thr Leu Gly Gly Leu Phe Ser  
 95 100 105  
 Gln Pro Thr Gln Ala Pro Thr Gln Ser Asn Gln Leu Ile Asn Thr  
 110 115 120  
 Ala Ser Ala Leu Ser Ala Pro Thr Leu Leu Gly Asp Glu Arg Asp  
 125 130 135  
 Ala Ile Leu Ala Lys Trp Asn Gln Leu Gln Ala Phe Trp Gly Thr  
 140 145 150  
 Gly Lys Gly Tyr Phe Asn Asn Asn Ile Pro Pro Val Glu Phe Thr  
 155 160 165  
 Gln Glu Asn Pro Phe Cys Arg Phe Lys Ala Val Gly Tyr Ser Cys  
 170 175 180  
 Met Pro Ser Asn Lys Asp Glu Asp Gly Leu Val Val Leu Val Phe  
 185 190 195  
 Asn Lys Lys Glu Thr Glu Ile Arg Ser Gln Gln Gln Gln Leu Val  
 200 205 210  
 Glu Ser Leu His Lys Val Leu Gly Gly Asn Gln Thr Leu Thr Val  
 215 220 225  
 Asn Val Glu Gly Thr Lys Thr Leu Pro Asp Asp Gln Thr Glu Val  
 230 235 240  
 Val Ile Tyr Val Val Glu Arg Ser Pro Asn Gly Thr Ser Arg Arg  
 245 250 255  
 Val Pro Ala Thr Thr Leu Tyr Ala His Phe Glu Gln Ala Asn Ile  
 260 265 270  
 Lys Thr Gln Leu Gln Gln Leu Gly Val Thr Leu Ser Met Thr Arg  
 275 280 285  
 Thr Glu Leu Ser Pro Ala Gln Ile Lys Gln Leu Leu Gln Asn Pro  
 290 295 300  
 Pro Ala Gly Val Asp Pro Ile Ile Trp Glu Gln Ala Lys Val Asp  
 305 310 315  
 Asn Pro Asp Ser Glu Lys Leu Ile Pro Val Pro Met Val Gly Phe  
 320 325 330

Lys Glu Leu Leu Arg Arg Leu Lys Val Gln Asp Gln Met Thr Lys  
 335 340 345  
 Gln His Gln Thr Arg Leu Asp Ile Ile Ser Glu Asp Ile Ser Glu  
 350 355 360  
 Leu Gln Lys Asn Gln Thr Thr Ser Val Ala Lys Ile Ala Gln Tyr  
 365 370 375  
 Lys Arg Lys Leu Met Asp Leu Ser His Arg Thr Leu Gln Val Leu  
 380 385 390  
 Ile Lys Gln Glu Ile Gln Arg Lys Ser Gly Tyr Ala Ile Gln Ala  
 395 400 405  
 Asp Glu Glu Gln Leu Arg Val Gln Leu Asp Thr Ile Gln Gly Glu  
 410 415 420  
 Leu Asn Ala Pro Thr Gln Phe Lys Gly Arg Leu Asn Glu Leu Met  
 425 430 435  
 Ser Gln Ile Arg Met Gln Asn His Phe Gly Ala Val Arg Ser Glu  
 440 445 450  
 Glu Arg Tyr Tyr Ile Asp Ala Asp Leu Leu Arg Glu Ile Lys Gln  
 455 460 465  
 His Leu Lys Gln Gln Gln Glu Gly Leu Ser His Leu Ile Ser Ile  
 470 475 480  
 Ile Lys Asp Asp Leu Glu Asp Ile Lys Leu Val Glu His Gly Leu  
 485 490 495  
 Asn Glu Thr Ile His Ile Arg Gly Gly Val Phe Ser  
 500 505

<210> 19  
 <211> 592  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2758310CD1

<400> 19  
 Met Trp Phe Cys Gly Gln Ser Thr Pro Phe Gly Cys Glu Leu His  
 1 5 10 15  
 Asp Thr Cys Val Gln Leu Cys His Phe His Ser Ala Leu Leu His  
 20 25 30  
 Arg Arg Gln Lys Pro Trp Pro Ser Pro Ala Val Phe Phe Arg Arg  
 35 40 45  
 Asn Val Arg Gly Leu Pro Pro Arg Phe Ser Ser Pro Thr Pro Leu  
 50 55 60  
 Trp Arg Lys Val Leu Ser Thr Ala Val Val Gly Ala Pro Leu Leu  
 65 70 75  
 Leu Gly Ala Arg Tyr Val Met Ala Glu Ala Arg Glu Lys Arg Arg  
 80 85 90  
 Met Arg Leu Val Val Asp Gly Met Gly Arg Phe Gly Arg Ser Leu  
 95 100 105  
 Lys Val Gly Leu Gln Ile Ser Leu Asp Tyr Trp Trp Cys Thr Asn  
 110 115 120  
 Val Val Leu Arg Gly Trp Lys Ser Pro Gly Tyr Leu Glu Val Met  
 125 130 135  
 Ser Ala Cys His Gln Arg Ala Ala Asp Ala Leu Val Ala Gly Ala  
 140 145 150  
 Ile Ser Asn Gly Gly Leu Tyr Val Lys Leu Gly Gln Gly Leu Cys  
 155 160 165  
 Ser Phe Asn His Leu Leu Pro Pro Glu Tyr Thr Arg Thr Leu Arg  
 170 175 180  
 Val Leu Glu Asp Arg Ala Leu Lys Arg Gly Phe Gln Glu Val Asp  
 185 190 195  
 Glu Leu Phe Leu Glu Asp Phe Gln Ala Leu Pro His Glu Leu Phe  
 200 205 210  
 Gln Glu Phe Asp Tyr Gln Pro Ile Ala Ala Ala Ser Leu Ala Gln  
 215 220 225  
 Val His Arg Ala Lys Leu His Asp Gly Thr Ser Val Ala Val Lys  
 230 235 240  
 Val Gln Tyr Ile Asp Leu Arg Asp Arg Phe Asp Gly Asp Ile His  
 245 250 255

Thr Leu Glu Leu Leu Leu Arg Leu Val Glu Val Met His Pro Ser  
 260 265 270  
 Phe Gly Phe Ser Trp Val Leu Gln Asp Leu Lys Gly Thr Leu Ala  
 275 280 285  
 Gln Glu Leu Asp Phe Glu Asn Glu Gly Arg Asn Ala Glu Arg Cys  
 290 295 300  
 Ala Arg Glu Leu Ala His Phe Pro Tyr Val Val Pro Arg Val  
 305 310 315  
 His Trp Asp Lys Ser Ser Lys Arg Val Leu Thr Ala Asp Phe Cys  
 320 325 330  
 Ala Gly Cys Lys Val Asn Asp Val Glu Ala Ile Arg Ser Gln Gly  
 335 340 345  
 Leu Ala Val His Asp Ile Ala Glu Lys Leu Ile Lys Ala Phe Ala  
 350 355 360  
 Glu Gln Ile Phe Tyr Thr Gly Phe Ile His Ser Asp Pro His Pro  
 365 370 375  
 Gly Asn Val Leu Val Arg Lys Gly Pro Asp Gly Lys Ala Glu Leu  
 380 385 390  
 Val Leu Leu Asp His Gly Leu Tyr Gln Phe Leu Glu Glu Lys Asp  
 395 400 405  
 Arg Ala Ala Leu Cys Gln Leu Trp Arg Ala Ile Ile Leu Arg Asp  
 410 415 420  
 Asp Ala Ala Met Arg Ala His Ala Ala Leu Gly Val Gln Asp  
 425 430 435  
 Tyr Leu Leu Phe Ala Glu Met Leu Met Gln Arg Pro Val Arg Leu  
 440 445 450  
 Gly Gln Leu Trp Gly Ser His Leu Leu Ser Arg Glu Glu Ala Ala  
 455 460 465  
 Tyr Met Val Asp Met Ala Arg Glu Arg Phe Glu Ala Val Met Ala  
 470 475 480  
 Val Leu Arg Glu Leu Pro Arg Pro Met Leu Leu Val Leu Arg Asn  
 485 490 495  
 Ile Asn Thr Val Arg Ala Ile Asn Val Ala Leu Gly Ala Pro Val  
 500 505 510  
 Asp Arg Tyr Phe Leu Met Ala Lys Arg Ala Val Arg Gly Trp Ser  
 515 520 525  
 Arg Leu Ala Gly Ala Thr Tyr Arg Gly Val Tyr Gly Thr Ser Leu  
 530 535 540  
 Leu Arg His Ala Lys Val Val Trp Glu Met Leu Lys Phe Glu Val  
 545 550 555  
 Ala Leu Arg Leu Glu Thr Leu Ala Met Arg Leu Thr Ala Leu Leu  
 560 565 570  
 Ala Arg Ala Leu Val His Leu Ser Leu Val Pro Pro Ala Glu Glu  
 575 580 585  
 Leu Tyr Gln Tyr Leu Glu Thr  
 590

&lt;210&gt; 20

&lt;211&gt; 841

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2762348CD1

&lt;400&gt; 20

Met Ala Ser Val Phe Arg Ser Glu Glu Met Cys Leu Ser Gln Leu  
 1 5 10 15  
 Phe Leu Gln Val Glu Ala Ala Tyr Cys Cys Val Ala Glu Leu Gly  
 20 25 30  
 Glu Leu Gly Leu Val Gln Phe Lys Asp Leu Asn Met Asn Val Asn  
 35 40 45  
 Ser Phe Gln Arg Lys Phe Val Asn Glu Val Arg Arg Cys Glu Ser  
 50 55 60  
 Leu Glu Arg Ile Leu Arg Phe Leu Glu Asp Glu Met Gln Asn Glu  
 65 70 75  
 Ile Val Val Gln Leu Leu Glu Lys Ser Pro Leu Thr Pro Leu Pro  
 80 85 90

Arg	Glu	Met	Ile	Thr	Leu	Glu	Thr	Val	Leu	Glu	Lys	Leu	Glu	Gly
				95					100					105
Glu	Leu	Gln	Glu	Ala	Asn	Gln	Asn	Gln	Gln	Ala	Leu	Lys	Gln	Ser
				110					115					120
Phe	Leu	Glu	Leu	Thr	Glu	Leu	Lys	Tyr	Leu	Leu	Lys	Lys	Thr	Gln
				125					130					135
Asp	Phe	Phe	Glu	Thr	Glu	Thr	Asn	Leu	Ala	Asp	Asp	Phe	Phe	Thr
				140					145					150
Glu	Asp	Thr	Ser	Gly	Leu	Leu	Glu	Leu	Lys	Ala	Val	Pro	Ala	Tyr
				155					160					165
Met	Thr	Gly	Lys	Leu	Gly	Phe	Ile	Ala	Gly	Cys	Asp	Pro	Thr	Gly
				170					175					180
Lys	Arg	Met	Ala	Ser	Phe	Glu	Arg	Leu	Leu	Trp	Arg	Val	Cys	Arg
				185					190					195
Gly	Asn	Val	Tyr	Leu	Lys	Phe	Ser	Glu	Met	Asp	Ala	Pro	Leu	Glu
				200					205					210
Asp	Pro	Val	Thr	Lys	Glu	Glu	Ile	Gln	Lys	His	Ile	Phe	Ile	Ile
				215					220					225
Phe	Tyr	Gln	Gly	Glu	Gln	Leu	Arg	Gln	Lys	Ile	Lys	Lys	Ile	Cys
				230					235					240
Asp	Gly	Phe	Arg	Ala	Thr	Val	Tyr	Pro	Cys	Pro	Glu	Pro	Ala	Val
				245					250					255
Glu	Arg	Arg	Glu	Met	Leu	Glu	Ser	Val	Asn	Val	Arg	Leu	Glu	Asp
				260					265					270
Leu	Ile	Thr	Val	Ile	Thr	Gln	Thr	Glu	Ser	His	Arg	Gln	Arg	Leu
				275					280					285
Leu	Gln	Glu	Ala	Ala	Ala	Asn	Trp	His	Ser	Trp	Leu	Ile	Lys	Val
				290					295					300
Gln	Lys	Met	Lys	Ala	Val	Tyr	His	Ile	Leu	Asn	Met	Cys	Asn	Ile
				305					310					315
Asp	Val	Thr	Gln	Gln	Cys	Val	Ile	Ala	Glu	Ile	Trp	Phe	Pro	Val
				320					325					330
Ala	Asp	Ala	Thr	Arg	Ile	Lys	Arg	Ala	Leu	Glu	Gln	Gly	Met	Glu
				335					340					345
Leu	Ser	Gly	Ser	Ser	Met	Ala	Pro	Ile	Met	Thr	Thr	Val	Gln	Ser
				350					355					360
Lys	Thr	Ala	Pro	Pro	Thr	Phe	Asn	Arg	Thr	Asn	Lys	Phe	Thr	Ala
				365					370					375
Gly	Phe	Gln	Asn	Ile	Val	Asp	Ala	Tyr	Gly	Val	Gly	Ser	Tyr	Arg
				380					385					390
Glu	Ile	Asn	Pro	Ala	Pro	Tyr	Thr	Ile	Ile	Thr	Phe	Pro	Phe	Leu
				395					400					405
Phe	Ala	Val	Met	Phe	Gly	Asp	Cys	Gly	His	Gly	Thr	Val	Met	Leu
				410					415					420
Leu	Ala	Ala	Leu	Trp	Met	Ile	Leu	Asn	Glu	Arg	Arg	Leu	Leu	Ser
				425					430					435
Gln	Lys	Thr	Asp	Asn	Glu	Ile	Trp	Asn	Thr	Phe	Phe	His	Gly	Arg
				440					445					450
Tyr	Leu	Ile	Leu	Leu	Met	Gly	Ile	Phe	Ser	Ile	Tyr	Thr	Gly	Leu
				455					460					465
Ile	Tyr	Asn	Asp	Cys	Phe	Ser	Lys	Ser	Leu	Asn	Ile	Phe	Gly	Ser
				470					475					480
Ser	Trp	Ser	Val	Gln	Pro	Met	Phe	Arg	Asn	Gly	Thr	Trp	Asn	Thr
				485					490					495
His	Val	Met	Glu	Glu	Ser	Leu	Tyr	Leu	Gln	Leu	Asp	Pro	Ala	Ile
				500					505					510
Pro	Gly	Val	Tyr	Phe	Gly	Asn	Pro	Tyr	Pro	Phe	Gly	Ile	Asp	Pro
				515					520					525
Ile	Trp	Asn	Leu	Ala	Ser	Asn	Lys	Leu	Thr	Phe	Leu	Asn	Ser	Tyr
				530					535					540
Lys	Met	Lys	Met	Ser	Val	Ile	Leu	Gly	Ile	Val	Gln	Met	Val	Phe
				545					550					555
Gly	Val	Ile	Leu	Ser	Leu	Phe	Asn	His	Ile	Tyr	Phe	Arg	Arg	Thr
				560					565					570
Leu	Asn	Ile	Ile	Leu	Gln	Phe	Ile	Pro	Glu	Met	Ile	Phe	Ile	Leu
				575					580					585
Cys	Leu	Phe	Gly	Tyr	Leu	Val	Phe	Met	Ile	Ile	Phe	Lys	Trp	Cys
				590					595					600

<400>	21														
Met	Ser	Glu	Cys	Pro	Leu	Ile	Leu	Tyr	Ile	His	Lys	His	Ile	Asp	
1				5					10					15	
Thr	Tyr	Ser	Gln	Ser	Tyr	Leu	Phe	Asn	Asp	Leu	Phe	Tyr	Pro	Val	
				20					25					30	
Tyr	Ser	Gly	Gly	Arg	Met	Val	Thr	Tyr	Glu	His	Leu	Arg	Glu	Val	
				35					40					45	
Val	Phe	Gly	Lys	Ser	Glu	Asp	Glu	His	Tyr	Pro	Leu	Trp	Lys	Ser	
				50					55					60	
Val	Ile	Gly	Gly	Met	Met	Ala	Gly	Val	Ile	Gly	Gln	Phe	Leu	Ala	
				65					70					75	
Asn	Pro	Thr	Asp	Leu	Val	Lys	Val	Gln	Met	Gln	Met	Glu	Gly	Lys	
				80					85					90	
Arg	Lys	Leu	Glu	Gly	Lys	Pro	Leu	Arg	Phe	Arg	Gly	Val	His	His	
				95					100					105	
Ala	Phe	Ala	Lys	Ile	Leu	Ala	Glu	Gly	Gly	Ile	Arg	Gly	Leu	Trp	
				110					115					120	
Ala	Gly	Trp	Val	Pro	Asn	Ile	Gln	Arg	Ala	Ala	Leu	Val	Asn	Met	
				125					130					135	
Gly	Asp	Leu	Thr	Thr	Tyr	Asp	Thr	Val	Lys	His	Tyr	Leu	Val	Leu	
				140					145					150	
Asn	Thr	Pro	Leu	Glu	Asp	Asn	Ile	Met	Thr	His	Gly	Leu	Ser	Ser	
				155					160					165	
Leu	Cys	Ser	Gly	Leu	Val	Ala	Ser	Ile	Leu	Gly	Thr	Pro	Ala	Asp	
				170					175					180	



Val Ile Lys Ser Arg Ile Met Asn Gln Pro Arg Asp Lys Gln Gly  
 185 190 195  
 Arg Gly Leu Leu Tyr Lys Ser Ser Thr Asp Cys Leu Ile Gln Ala  
 200 205 210  
 Val Gln Gly Glu Gly Phe Met Ser Leu Tyr Lys Gly Phe Leu Pro  
 215 220 225  
 Ser Trp Leu Arg Met Thr Pro Trp Ser Met Val Phe Trp Leu Thr  
 230 235 240  
 Tyr Glu Lys Ile Arg Glu Met Ser Gly Val Ser Pro Phe  
 245 250

&lt;210&gt; 22

&lt;211&gt; 229

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5108194CD1

&lt;400&gt; 22

Met Gly Asn Gly Val Lys Glu Gly Pro Val Arg Leu His Glu Asp  
 1 5 10 15  
 Ala Glu Ala Val Leu Ser Ser Ser Val Ser Ser Lys Arg Asp His  
 20 25 30  
 Arg Gln Val Leu Ser Ser Leu Leu Ser Gly Ala Leu Ala Gly Ala  
 35 40 45  
 Leu Ala Lys Thr Ala Val Ala Pro Leu Asp Arg Thr Lys Ile Ile  
 50 55 60  
 Phe Gln Val Ser Ser Lys Arg Phe Ser Ala Lys Glu Ala Phe Arg  
 65 70 75  
 Val Leu Tyr Tyr Thr Tyr Leu Asn Glu Gly Phe Leu Ser Leu Trp  
 80 85 90  
 Arg Gly Asn Ser Ala Thr Met Val Arg Val Val Pro Tyr Ala Ala  
 95 100 105  
 Ile Gln Phe Ser Ala His Glu Glu Tyr Lys Arg Ile Leu Gly Ser  
 110 115 120  
 Tyr Tyr Gly Phe Arg Gly Glu Ala Leu Pro Pro Trp Pro Arg Leu  
 125 130 135  
 Phe Ala Gly Ala Leu Ala Gly Thr Thr Ala Ala Ser Leu Thr Tyr  
 140 145 150  
 Pro Leu Asp Leu Val Arg Ala Arg Met Ala Val Thr Pro Lys Glu  
 155 160 165  
 Met Tyr Ser Asn Ile Phe His Val Phe Ile Arg Ile Ser Arg Glu  
 170 175 180  
 Glu Gly Leu Lys Thr Leu Tyr His Gly Phe Met Pro Thr Val Leu  
 185 190 195  
 Gly Val Ile Pro Tyr Ala Gly Leu Ser Phe Phe Thr Tyr Glu Thr  
 200 205 210  
 Leu Lys Ser Leu His Arg Glu Tyr Ser Gly Arg Lys Leu Ile Pro  
 215 220 225  
 Phe Ser Glu Gly

&lt;210&gt; 23

&lt;211&gt; 170

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5503122CD1

&lt;400&gt; 23

Met Tyr Asp Asn Leu Tyr Leu His Gly Ile Glu Asp Ser Glu Ala  
 1 5 10 15  
 Gly Ser Ala Asp Ser Tyr Thr Ser Arg Pro Ser Asp Ser Asp Val  
 20 25 30  
 Ser Leu Glu Glu Asp Arg Glu Ala Ile Arg Gln Glu Arg Glu Gln  
 35 40 45

Gln Ala Ala Ile Gln Leu Glu Arg Ala Lys Ser Lys Pro Val Ala  
 50 55 60  
 Phe Ala Val Lys Thr Asn Val Ser Tyr Cys Gly Ala Leu Asp Glu  
 65 70 75  
 Asp Val Pro Val Pro Ser Thr Ala Ile Ser Phe Asp Ala Lys Asp  
 80 85 90  
 Phe Leu His Ile Lys Glu Lys Tyr Asn Asn Asp Trp Trp Ile Gly  
 95 100 105  
 Arg Leu Val Lys Glu Gly Cys Glu Ile Gly Phe Ile Pro Ser Pro  
 110 115 120  
 Leu Arg Leu Glu Asn Ile Arg Ile Gln Gln Glu Gln Lys Arg Gly  
 125 130 135  
 Arg Phe His Gly Gly Lys Ser Ser Gly Asn Ser Ser Ser Ser Leu  
 140 145 150  
 Gly Glu Met Val Ser Gly Thr Phe Arg Ala Thr Pro Thr Ser Thr  
 155 160 165  
 Gly Glu Gly Cys Ser  
 170

&lt;210&gt; 24

&lt;211&gt; 655

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5517972CD1

&lt;400&gt; 24

Met Ser Ser Ser Asn Val Glu Val Phe Ile Pro Val Ser Gln Gly  
 1 5 10 15  
 Asn Thr Asn Gly Phe Pro Ala Thr Ala Ser Asn Asp Leu Lys Ala  
 20 25 30  
 Phe Thr Glu Gly Ala Val Leu Ser Phe His Asn Ile Cys Tyr Arg  
 35 40 45  
 Val Lys Leu Lys Ser Gly Phe Leu Pro Cys Arg Lys Pro Val Glu  
 50 55 60  
 Lys Glu Ile Leu Ser Asn Ile Asn Gly Ile Met Lys Pro Gly Leu  
 65 70 75  
 Asn Ala Ile Leu Gly Pro Thr Gly Gly Gly Lys Ser Ser Leu Leu  
 80 85 90  
 Asp Val Leu Ala Ala Arg Lys Asp Pro Ser Gly Leu Ser Gly Asp  
 95 100 105  
 Val Leu Ile Asn Gly Ala Pro Arg Pro Ala Asn Phe Lys Cys Asn  
 110 115 120  
 Ser Gly Tyr Val Val Gln Asp Asp Val Val Met Gly Thr Leu Thr  
 125 130 135  
 Val Arg Glu Asn Leu Gln Phe Ser Ala Ala Leu Arg Leu Ala Thr  
 140 145 150  
 Thr Met Thr Asn His Glu Lys Asn Glu Arg Ile Asn Arg Val Ile  
 155 160 165  
 Gln Glu Leu Gly Leu Asp Lys Val Ala Asp Ser Lys Val Gly Thr  
 170 175 180  
 Gln Phe Ile Arg Gly Val Ser Gly Gly Glu Arg Lys Arg Thr Ser  
 185 190 195  
 Ile Gly Met Glu Leu Ile Thr Asp Pro Ser Ile Leu Phe Leu Asp  
 200 205 210  
 Glu Pro Thr Thr Gly Leu Asp Ser Ser Thr Ala Asn Ala Val Leu  
 215 220 225  
 Leu Leu Leu Lys Arg Met Ser Lys Gln Gly Arg Thr Ile Ile Phe  
 230 235 240  
 Ser Ile His Gln Pro Arg Tyr Ser Ile Phe Lys Leu Phe Asp Ser  
 245 250 255  
 Leu Thr Leu Leu Ala Ser Gly Arg Leu Met Phe His Gly Pro Ala  
 260 265 270  
 Gln Glu Ala Leu Gly Tyr Phe Glu Ser Ala Gly Tyr His Cys Glu  
 275 280 285  
 Ala Tyr Asn Asn Pro Ala Asp Phe Phe Leu Asp Ile Ile Asn Gly  
 290 295 300

```

Asp Ser Thr Ala Val Ala Leu Asn Arg Glu Glu Asp Phe Lys Ala
305 310 315
Thr Glu Ile Ile Glu Pro Ser Lys Gln Asp Lys Pro Leu Ile Glu
320 325 330
Lys Leu Ala Glu Ile Tyr Val Asn Ser Ser Phe Tyr Lys Glu Thr
335 340 345
Lys Ala Glu Leu His Gln Leu Ser Gly Gly Glu Lys Lys Lys Lys
350 355 360
Ile Thr Val Phe Lys Glu Ile Ser Tyr Thr Thr Ser Phe Cys His
365 370 375
Gln Leu Arg Trp Val Ser Lys Arg Ser Phe Lys Asn Leu Leu Gly
380 385 390
Asn Pro Gln Ala Ser Ile Ala Gln Ile Ile Val Thr Val Val Leu
395 400 405
Gly Leu Val Ile Gly Ala Ile Tyr Phe Gly Leu Lys Asn Asp Ser
410 415 420
Thr Gly Ile Gln Asn Arg Ala Gly Val Leu Phe Phe Leu Thr Thr
425 430 435
Asn Gln Cys Phe Ser Ser Val Ser Ala Val Glu Leu Phe Val Val
440 445 450
Glu Lys Lys Leu Phe Ile His Glu Tyr Ile Ser Gly Tyr Tyr Arg
455 460 465
Val Ser Ser Tyr Phe Leu Gly Lys Leu Leu Ser Asp Leu Leu Pro
470 475 480
Met Arg Met Leu Pro Ser Ile Ile Phe Thr Cys Ile Val Tyr Phe
485 490 495
Met Leu Gly Leu Lys Pro Lys Ala Asp Ala Phe Phe Val Met Met
500 505 510
Phe Thr Leu Met Met Val Ala Tyr Ser Ala Ser Ser Met Ala Leu
515 520 525
Ala Ile Ala Ala Gly Gln Ser Val Val Ser Val Ala Thr Leu Leu
530 535 540
Met Thr Ile Cys Phe Val Phe Met Met Ile Phe Ser Gly Leu Leu
545 550 555
Val Asn Leu Thr Thr Ile Ala Ser Trp Leu Ser Trp Leu Gln Tyr
560 565 570
Phe Ser Ile Pro Arg Tyr Gly Phe Thr Ala Leu Gln His Asn Glu
575 580 585
Phe Leu Gly Gln Asn Phe Cys Pro Gly Leu Asn Ala Thr Gly Asn
590 595 600
Asn Pro Cys Asn Tyr Ala Thr Cys Thr Gly Glu Glu Tyr Leu Val
605 610 615
Lys Gln Gly Ile Asp Leu Ser Pro Trp Gly Leu Trp Lys Asn His
620 625 630
Val Ala Leu Ala Cys Met Ile Val Ile Phe Leu Thr Ile Ala Tyr
635 640 645
Leu Lys Leu Leu Phe Leu Lys Lys Tyr Ser
650 655

```

&lt;210&gt; 25

&lt;211&gt; 184

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5593114CD1

&lt;400&gt; 25

```

Met Trp Val Phe Gly Tyr Gly Ser Leu Ile Trp Lys Val Asp Phe
1 5 10 15
Pro Tyr Gln Asp Lys Leu Val Gly Tyr Ile Thr Asn Tyr Ser Arg
20 25 30
Arg Phe Trp Gln Gly Ser Thr Asp His Arg Gly Val Pro Gly Lys
35 40 45
Pro Gly Arg Val Val Thr Leu Val Glu Asp Pro Ala Gly Cys Val
50 55 60
Trp Gly Val Ala Tyr Arg Leu Pro Val Gly Lys Glu Glu Glu Val
65 70 75

```

Lys Ala Tyr Leu Asp Phe Arg Glu Lys Gly Gly Tyr Arg Thr Thr  
 80 85 90  
 Thr Val Ile Phe Tyr Pro Lys Asp Pro Thr Thr Lys Pro Phe Ser  
 95 100 105  
 Val Leu Leu Tyr Ile Gly Thr Cys Asp Asn Pro Asp Tyr Leu Gly  
 110 115 120  
 Pro Ala Pro Leu Glu Asp Ile Ala Glu Gln Ile Phe Asn Ala Ala  
 125 130 135  
 Gly Pro Ser Gly Arg Asn Thr Glu Tyr Leu Phe Glu Leu Ala Asn  
 140 145 150  
 Ser Ile Arg Asn Leu Val Pro Glu Glu Ala Asp Glu His Leu Phe  
 155 160 165  
 Ala Leu Glu Lys Leu Val Lys Glu Arg Leu Glu Gly Lys Gln Asn  
 170 175 180  
 Leu Asn Cys Ile

<210> 26  
 <211> 154  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 044775CD1

<400> 26  
 Met Gly Ala Phe Glu Cys Val Arg Lys Val Tyr Gln Thr Asp Gly  
 1 5 10 15  
 Leu Lys Gly Phe Tyr Arg Gly Met Ser Ala Ser Tyr Ala Gly Ile  
 20 25 30  
 Ser Glu Thr Val Ile His Phe Val Ile Tyr Glu Ser Ile Lys Gln  
 35 40 45  
 Lys Leu Leu Glu Tyr Lys Thr Ala Ser Thr Met Glu Asn Asp Glu  
 50 55 60  
 Glu Ser Val Lys Glu Ala Ser Asp Phe Val Gly Met Met Leu Ala  
 65 70 75  
 Ala Ala Thr Ser Lys Thr Cys Ala Thr Thr Ile Ala Tyr Pro His  
 80 85 90  
 Glu Val Val Arg Thr Arg Leu Arg Glu Glu Gly Thr Lys Tyr Arg  
 95 100 105  
 Ser Phe Phe Gln Thr Leu Ser Leu Leu Val Gln Glu Glu Gly Tyr  
 110 115 120  
 Gly Ser Leu Tyr Arg Gly Leu Thr Thr His Leu Val Arg Gln Ile  
 125 130 135  
 Pro Asn Thr Ala Ile Met Met Ala Thr Tyr Glu Leu Val Val Tyr  
 140 145 150  
 Leu Leu Asn Gly

<210> 27  
 <211> 438  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 116588CD1

<400> 27  
 Met Leu Leu Val Thr Pro Arg Pro Glu Arg Gly Gly Arg Gly Thr  
 1 5 10 15  
 Glu Leu Gly Glu Phe Cys Gly Thr Pro Leu Leu Phe Ser Ser Tyr  
 20 25 30  
 Phe Cys Tyr Asp Asn Pro Ala Ala Leu Gln Thr Gln Val Lys Arg  
 35 40 45  
 Asp Met Gln Val Asn Thr Thr Lys Phe Met Leu Leu Tyr Ala Trp  
 50 55 60  
 Tyr Ser Trp Pro Asn Val Val Leu Cys Phe Phe Gly Gly Phe Leu  
 65 70 75

Ile	Asp	Arg	Val	Phe	Gly	Ile	Arg	Trp	Gly	Thr	Ile	Ile	Phe	Ser	
				80					85					90	
Cys	Phe	Val	Cys	Ile	Gly	Gln	Val	Val	Phe	Ala	Leu	Gly	Gly	Ile	
				95					100					105	
Phe	Asn	Ala	Phe	Trp	Leu	Met	Glu	Phe	Gly	Arg	Phe	Val	Phe	Gly	
				110					115					120	
Ile	Gly	Gly	Glu	Ser	Leu	Ala	Val	Ala	Gln	Asn	Thr	Tyr	Ala	Val	
				125					130					135	
Ser	Trp	Phe	Lys	Gly	Lys	Glu	Leu	Asn	Leu	Val	Phe	Gly	Leu	Gln	
				140					145					150	
Leu	Ser	Met	Ala	Arg	Ile	Gly	Ser	Thr	Val	Asn	Met	Asn	Leu	Met	
				155					160					165	
Gly	Trp	Leu	Tyr	Ser	Lys	Ile	Glu	Ala	Leu	Leu	Gly	Ser	Ala	Gly	
				170					175					180	
His	Thr	Thr	Leu	Gly	Ile	Thr	Leu	Met	Ile	Gly	Gly	Val	Thr	Cys	
				185					190					195	
Ile	Leu	Ser	Leu	Ile	Cys	Ala	Leu	Ala	Leu	Ala	Tyr	Leu	Asp	Gln	
				200					205					210	
Arg	Ala	Glu	Arg	Ile	Leu	His	Lys	Glu	Gln	Gly	Lys	Thr	Gly	Glu	
				215					220					225	
Val	Ile	Lys	Leu	Thr	Asp	Val	Lys	Asp	Phe	Ser	Leu	Pro	Leu	Trp	
				230					235					240	
Leu	Ile	Phe	Ile	Ile	Cys	Val	Cys	Tyr	Tyr	Val	Ala	Val	Phe	Pro	
				245					250					255	
Phe	Ile	Gly	Leu	Gly	Lys	Val	Phe	Phe	Thr	Glu	Lys	Phe	Gly	Phe	
				260					265					270	
Ser	Ser	Gln	Ala	Ala	Ser	Ala	Ile	Asn	Ser	Val	Val	Tyr	Val	Ile	
				275					280					285	
Ser	Ala	Pro	Met	Ser	Pro	Val	Phe	Gly	Leu	Leu	Val	Asp	Lys	Thr	
				290					295					300	
Gly	Lys	Asn	Ile	Ile	Trp	Val	Leu	Cys	Ala	Val	Ala	Ala	Thr	Leu	
				305					310					315	
Val	Ser	His	Met	Met	Leu	Ala	Phe	Thr	Met	Trp	Asn	Pro	Trp	Ile	
				320					325					330	
Ala	Met	Cys	Leu	Leu	Gly	Leu	Ser	Tyr	Ser	Leu	Leu	Ala	Cys	Ala	
				335					340					345	
Leu	Trp	Pro	Met	Val	Ala	Phe	Val	Val	Pro	Glu	His	Gln	Leu	Gly	
				350					355					360	
Thr	Ala	Tyr	Gly	Phe	Met	Gln	Ser	Ile	Gln	Asn	Leu	Gly	Leu	Ala	
				365					370					375	
Ile	Ile	Ser	Ile	Ile	Ala	Gly	Met	Ile	Leu	Asp	Ser	Arg	Gly	Tyr	
				380					385					390	
Leu	Phe	Leu	Glu	Val	Phe	Phe	Ile	Ala	Cys	Val	Ser	Leu	Ser	Leu	
				395					400					405	
Leu	Ser	Val	Val	Leu	Leu	Tyr	Leu	Val	Asn	Arg	Ala	Gln	Gly	Gly	
				410					415					420	
Asn	Leu	Asn	Tyr	Ser	Ala	Arg	Gln	Arg	Glu	Glu	Ile	Lys	Phe	Ser	
				425					430					435	
His	Thr	Glu													

&lt;210&gt; 28

&lt;211&gt; 237

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 875369CD1

&lt;400&gt; 28

Met	Ala	His	Val	Gly	Ser	Arg	Lys	Arg	Ser	Arg	Ser	Arg	Ser	Arg	
				5					10					15	
Ser	Arg	Gly	Arg	Gly	Ser	Glu	Lys	Arg	Lys	Lys	Lys	Ser	Arg	Lys	
				20					25					30	
Asp	Thr	Ser	Arg	Asn	Cys	Ser	Ala	Ser	Thr	Ser	Gln	Gly	Arg	Lys	
				35					40					45	
Ala	Ser	Thr	Ala	Pro	Gly	Ala	Glu	Ala	Ser	Pro	Ser	Pro	Cys	Ile	
				50					55					60	

Thr Glu Arg Ser Lys Gln Lys Ala Arg Arg Arg Thr Arg Ser Ser  
 65 70 75  
 Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser  
 80 85 90  
 Ser Ser Ser Ser Ser Ser Ser Ser Asp Gly Arg Lys Lys Arg Gly  
 95 100 105  
 Lys Tyr Lys Asp Lys Arg Arg Lys Lys Lys Lys Arg Lys Lys  
 110 115 120  
 Leu Lys Lys Lys Gly Lys Glu Lys Ala Glu Ala Gln Gln Val Glu  
 125 130 135  
 Ala Leu Pro Gly Pro Ser Leu Asp Gln Trp His Arg Ser Ala Gly  
 140 145 150  
 Glu Glu Glu Asp Gly Pro Val Leu Thr Asp Glu Gln Lys Ser Arg  
 155 160 165  
 Ile Gln Ala Met Lys Pro Met Thr Lys Glu Glu Trp Asp Ala Arg  
 170 175 180  
 Gln Ser Ile Ile Arg Lys Val Val Asp Pro Glu Thr Gly Arg Thr  
 185 190 195  
 Arg Leu Ile Lys Gly Asp Gly Glu Val Leu Glu Glu Ile Val Thr  
 200 205 210  
 Lys Glu Arg His Arg Glu Ile Asn Lys Gln Ala Thr Arg Gly Asp  
 215 220 225  
 Cys Leu Ala Phe Gln Met Arg Ala Gly Leu Leu Pro  
 230 235

&lt;210&gt; 29

&lt;211&gt; 219

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1325518CD1

&lt;400&gt; 29

Met Lys Leu Leu Leu Trp Ala Cys Ile Val Cys Val Ala Phe Ala  
 1 5 10 15  
 Arg Lys Arg Arg Phe Pro Phe Ile Gly Glu Asp Asp Asn Asp Asp  
 20 25 30  
 Gly His Pro Leu His Pro Ser Leu Asn Ile Pro Tyr Gly Ile Arg  
 35 40 45  
 Asn Leu Pro Pro Pro Leu Tyr Tyr Arg Pro Val Asn Thr Val Pro  
 50 55 60  
 Ser Tyr Pro Gly Asn Thr Tyr Thr Asp Thr Gly Leu Pro Ser Tyr  
 65 70 75  
 Pro Trp Ile Leu Thr Ser Pro Gly Phe Pro Tyr Val Tyr His Ile  
 80 85 90  
 Arg Gly Phe Pro Leu Ala Thr Gln Leu Asn Val Pro Pro Leu Pro  
 95 100 105  
 Pro Arg Gly Phe Pro Phe Val Pro Pro Ser Arg Phe Phe Ser Ala  
 110 115 120  
 Ala Ala Ala Pro Ala Ala Pro Pro Ile Ala Ala Glu Pro Ala Ala  
 125 130 135  
 Ala Ala Pro Leu Thr Ala Thr Pro Val Ala Ala Glu Pro Ala Ala  
 140 145 150  
 Gly Ala Pro Val Ala Ala Glu Pro Ala Ala Glu Ala Pro Val Gly  
 155 160 165  
 Ala Glu Pro Ala Ala Glu Ala Pro Val Ala Ala Glu Pro Ala Ala  
 170 175 180  
 Glu Ala Pro Val Gly Val Glu Pro Ala Ala Glu Glu Pro Ser Pro  
 185 190 195  
 Ala Glu Pro Ala Thr Ala Lys Pro Ala Ala Pro Glu Pro His Pro  
 200 205 210  
 Ser Pro Ser Leu Glu Gln Ala Asn Gln  
 215

&lt;210&gt; 30

&lt;211&gt; 707

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2060987CD1

&lt;400&gt; 30

```

Met Ala Ala Ala Ala Thr Ala Ala Glu Gly Val Pro Ser Arg Gly
1      5      10
Pro Pro Gly Glu Val Ile His Leu Asn Val Gly Gly Lys Arg Phe
20     25
Ser Thr Ser Arg Gln Thr Leu Thr Trp Ile Pro Asp Ser Phe Phe
35     40
Ser Ser Leu Leu Ser Gly Arg Ile Ser Thr Leu Lys Asp Glu Thr
50     55
Gly Ala Ile Phe Ile Asp Arg Asp Pro Thr Val Phe Ala Pro Ile
65     70
Leu Asn Phe Leu Arg Thr Lys Glu Leu Asp Pro Arg Gly Val His
80     85
Gly Ser Ser Leu Leu His Glu Ala Gln Phe Tyr Gly Leu Thr Pro
95     100
Leu Val Arg Arg Leu Gln Leu Arg Glu Glu Leu Asp Arg Ser Ser
110    115
Cys Gly Asn Val Leu Phe Asn Gly Tyr Leu Pro Pro Pro Val Phe
125    130
Pro Val Lys Arg Arg Asn Arg His Ser Leu Val Gly Pro Gln Gln
140    145
Leu Gly Gly Arg Pro Ala Pro Val Arg Arg Ser Asn Thr Met Pro
155    160
Pro Asn Leu Gly Asn Ala Gly Leu Leu Gly Arg Met Leu Asp Glu
170    175
Lys Thr Pro Pro Ser Pro Ser Gly Gln Pro Glu Glu Pro Gly Met
185    190
Val Arg Leu Val Cys Gly His His Asn Trp Ile Ala Val Ala Tyr
200    205
Thr Gln Phe Leu Val Cys Tyr Arg Leu Lys Glu Ala Ser Gly Trp
215    220
Gln Leu Val Phe Ser Ser Pro Arg Leu Asp Trp Pro Ile Glu Arg
230    235
Leu Ala Leu Thr Ala Arg Val His Gly Gly Ala Leu Gly Glu His
245    250
Asp Lys Met Val Ala Ala Ala Thr Gly Ser Glu Ile Leu Leu Trp
260    265
Ala Leu Gln Ala Glu Gly Gly Gly Ser Glu Ile Gly Val Phe His
275    280
Leu Gly Val Pro Val Glu Ala Leu Phe Phe Val Gly Asn Gln Leu
290    295
Ile Ala Thr Ser His Thr Gly Arg Ile Gly Val Trp Asn Ala Val
305    310
Thr Lys His Trp Gln Val Gln Glu Val Gln Pro Ile Thr Ser Tyr
320    325
Asp Ala Ala Gly Ser Phe Leu Leu Leu Gly Cys Asn Asn Gly Ser
335    340
Ile Tyr Tyr Val Asp Val Gln Lys Phe Pro Leu Arg Met Lys Asp
350    355
Asn Asp Leu Leu Val Ser Glu Leu Tyr Arg Asp Pro Ala Glu Asp
365    370
Gly Val Thr Ala Leu Ser Val Tyr Leu Thr Pro Lys Thr Ser Asp
380    385
Ser Gly Asn Trp Ile Glu Ile Ala Tyr Gly Thr Ser Ser Gly Gly
395    400
Val Arg Val Ile Val Gln His Pro Glu Thr Val Gly Ser Gly Pro
410    415
Gln Leu Phe Gln Thr Phe Thr Val His Arg Ser Pro Val Thr Lys
425    430
Ile Met Leu Ser Glu Lys His Leu Ile Ser Val Cys Ala Asp Asn
440    445
Asn His Val Arg Thr Trp Ser Val Thr Arg Phe Arg Gly Met Ile
455    460
Ser Thr Gln Pro Gly Ser Thr Pro Leu Ala Ser Phe Lys Ile Leu

```

Ala Leu Glu Ser	470	Ala Asp Gly His Gly	475	Gly Cys Ser Ala Gly	480
Asp Ile Gly Pro	485	Tyr Gly Glu Arg Asp	490	Asp Gln Gln Val Phe	495
Gln Lys Val Val	500	Pro Ser Ala Ser Gln	505	Leu Phe Val Arg Leu	510
Ser Thr Gly Gln	515	Arg Val Cys Ser Val	520	Arg Ser Val Asp Gly	525
Pro Thr Thr Ala	530	Phe Thr Val Leu Glu	535	Cys Glu Gly Ser Arg	540
Leu Gly Ser Arg	545	Pro Arg Arg Tyr Leu	550	Leu Thr Gly Gln Ala	555
Gly Ser Leu Ala	560	Met Trp Asp Leu Thr	565	Thr Ala Met Asp Gly	570
Gly Gln Ala Pro	575	Ala Gly Gly Leu Thr	580	Glu Gln Glu Leu Met	585
Gln Leu Glu His	590	Cys Glu Leu Ala Pro	595	Pro Ala Pro Ser Ala	600
Ser Trp Gly Cys	605	Leu Pro Ser Pro Ser	610	Pro Arg Ile Ser Leu	615
Ser Leu His Ser	620	Ala Ser Ser Asn Thr	625	Ser Leu Ser Gly His	630
Gly Ser Pro Ser	635	Pro Pro Gln Ala Glu	640	Ala Arg Arg Arg Gly	645
Gly Ser Phe Val	650	Glu Arg Cys Gln Glu	655	Leu Val Arg Ser Gly	660
Asp Leu Arg Arg	665	Pro Pro Thr Pro Ala	670	Pro Trp Pro Ser Ser	675
Leu Gly Thr Pro	680	Leu Thr Pro Pro Lys	685	Met Lys Leu Asn Glu	690
Ser Phe	695		700		705

<210> 31  
 <211> 279  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2172064CD1

<400> 31

Met Cys Gly Arg Phe	1	Leu Arg Arg Leu Leu	10	Ala Glu Glu Ser Arg	15
Arg Ser Thr Pro Val	5	Gly Arg Leu Leu Leu	20	Pro Val Leu Leu Gly	25
Phe Arg Leu Val Leu	25	Leu Ala Ala Ser Gly	30	Pro Gly Val Tyr Gly	35
Asp Glu Gln Ser Glu	35	Phe Val Cys His Thr	40	Gln Gln Pro Gly Cys	45
Lys Ala Ala Cys Phe	50	Asp Ala Phe His Pro	55	Leu Ser Pro Leu Arg	60
Ser Trp Val Phe Gln	65	Val Ile Leu Val Ala	70	Val Pro Ser Ala Leu	75
Tyr Met Gly Phe Thr	80	Leu Tyr His Val Ile	85	Trp His Trp Glu Leu	90
Ser Gly Lys Gly Lys	95	Glu Glu Glu Thr Leu	100	Ile Gln Gly Arg Glu	105
Gly Asn Thr Asp Val	110	Pro Gly Ala Gly Ser	115	Leu Arg Leu Leu Trp	120
Ala Tyr Val Ala Gln	125	Leu Gly Ala Arg Leu	130	Val Leu Glu Gly Ala	135
Ala Leu Gly Leu Gln	140	Tyr His Leu Tyr Gly	145	Phe Gln Met Pro Ser	150
Ser Phe Ala Cys Arg	155	Arg Glu Pro Cys Leu	160	Gly Ser Ile Thr Cys	165
Asn Leu Ser Arg Pro	170	Ser Glu Lys Thr Ile	175	Phe Leu Lys Thr Met	180



	185		190		195
Phe Gly Val Ser	Gly Phe Cys Leu Leu	Phe Thr Phe Leu Glu Leu			
	200	205			210
Val Leu Leu Gly	Leu Gly Arg Trp Trp	Arg Thr Trp Lys His			Lys
	215	220			225
Ser Ser Ser Ser	Lys Tyr Phe Leu Thr	Ser Glu Ser Thr Arg Arg			
	230	235			240
His Lys Lys Ala	Thr Asp Ser Leu Pro	Val Val Glu Thr Lys Glu			
	245	250			255
Gln Phe Gln Glu	Ala Val Pro Gly Arg	Ser Leu Ala Gln Glu Lys			
	260	265			270
Gln Arg Pro Val	Gly Pro Arg Asp Ala				
	275				

<210> 32  
 <211> 154  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2219267CD1

<400> 32	
Met Val Thr Gly Leu Ala Ser Leu Leu Leu Leu Ala Gly Ala Gln	
1 5 10 15	
Tyr Leu Pro Gly Trp Thr Val Leu Phe Leu Ser Val Leu Gly Leu	
20 25 30	
Leu Ala Ser Arg Ala Val Ser Ala Leu Ser Ser Leu Phe Ala Ala	
35 40 45	
Glu Val Phe Pro Thr Val Ile Arg Gly Ala Gly Leu Gly Leu Val	
50 55 60	
Leu Gly Ala Gly Phe Leu Gly Gln Ala Ala Gly Pro Leu Asp Thr	
65 70 75	
Leu His Gly Arg Gln Gly Phe Phe Leu Gln Gln Val Val Phe Ala	
80 85 90	
Ser Leu Ala Val Leu Ala Leu Leu Cys Val Leu Leu Leu Pro Glu	
95 100 105	
Ser Arg Ser Arg Gly Leu Pro Gln Ser Leu Gln Asp Ala Asp Arg	
110 115 120	
Leu Arg Arg Ser Pro Leu Leu Arg Gly Arg Pro Arg Gln Asp His	
125 130 135	
Leu Pro Leu Leu Pro Pro Ser Asn Ser Tyr Trp Ala Gly His Thr	
140 145 150	
Pro Glu Gln His	

<210> 33  
 <211> 289  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2308629CD1

<400> 33	
Met Val Ala Gly Ala Val Ala Gly Ile Leu Glu His Cys Val Met	
1 5 10 15	
Tyr Pro Ile Asp Cys Val Lys Thr Arg Met Gln Ser Leu Gln Pro	
20 25 30	
Asp Pro Ala Ala Arg Tyr Arg Asn Val Leu Glu Ala Leu Trp Arg	
35 40 45	
Ile Ile Arg Thr Glu Gly Leu Trp Arg Pro Met Arg Gly Leu Asn	
50 55 60	
Val Thr Ala Thr Gly Ala Gly Pro Ala His Ala Leu Tyr Phe Ala	
65 70 75	
Cys Tyr Glu Lys Leu Lys Lys Thr Leu Ser Asp Val Ile His Pro	
80 85 90	
Gly Gly Asn Ser His Ile Ala Asn Gly Ala Ala Gly Cys Val Ala	

	95		100		105
Thr	Leu	Leu	His	Asp	Ala
	110		115		120
Gln	Arg	Met	Gln	Met	Tyr
	125		130		135
Cys	Val	Arg	Ala	Val	Trp
	140		145		150
Arg	Ser	Tyr	Thr	Thr	Gln
	155		160		165
Ile	His	Phe	Met	Thr	Tyr
	170		175		180
Gln	Arg	Arg	Tyr	Asn	Pro
	185		190		195
Ala	Gly	Ala	Val	Ala	Ala
	200		205		210
Lys	Thr	Leu	Leu	Asn	Thr
	215		220		225
Ile	Thr	Gly	His	Ile	Thr
	230		235		240
Tyr	Gln	Val	Gly	Gly	Val
	245		250		255
Arg	Val	Ile	Tyr	Gln	Ile
	260		265		270
Tyr	Glu	Phe	Phe	Lys	Tyr
	275		280		285
Arg	Ala	Gly	Lys		

&lt;210&gt; 34

&lt;211&gt; 300

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2660038CD1

&lt;400&gt; 34

Met	Asp	Phe	Leu	Met	Ser	Gly	Leu	Ala	Ala	Cys	Gly	Ala	Cys	Val
1				5					10					15
Phe	Thr	Asn	Pro	Leu	Glu	Val	Val	Lys	Thr	Arg	Met	Gln	Leu	Gln
				20					25					30
Gly	Glu	Leu	Gln	Ala	Pro	Gly	Thr	Tyr	Gln	Arg	His	Tyr	Arg	Asn
				35					40					45
Val	Phe	His	Ala	Phe	Ile	Thr	Ile	Gly	Lys	Val	Asp	Gly	Leu	Ala
				50					55					60
Ala	Leu	Gln	Lys	Gly	Leu	Ala	Pro	Ala	Leu	Leu	Tyr	Gln	Phe	Leu
				65					70					75
Met	Asn	Gly	Ile	Arg	Leu	Gly	Thr	Tyr	Gly	Leu	Ala	Glu	Ala	Gly
				80					85					90
Gly	Tyr	Leu	His	Thr	Ala	Glu	Ala	Thr	His	Ser	Pro	Ala	Arg	Ser
				95					100					105
Ala	Ala	Ala	Gly	Ala	Met	Ala	Gly	Val	Met	Gly	Ala	Tyr	Leu	Gly
				110					115					120
Ser	Pro	Ile	Tyr	Met	Val	Lys	Thr	His	Leu	Gln	Ala	Gln	Ala	Ala
				125					130					135
Ser	Glu	Ile	Ala	Val	Gly	His	Gln	Tyr	Lys	His	Gln	Gly	Met	Phe
				140					145					150
Gln	Ala	Leu	Thr	Glu	Ile	Gly	Gln	Lys	His	Gly	Leu	Val	Gly	Leu
				155					160					165
Trp	Arg	Gly	Ala	Leu	Gly	Gly	Leu	Pro	Arg	Val	Ile	Val	Gly	Ser
				170					175					180
Ser	Thr	Gln	Leu	Cys	Thr	Phe	Ser	Ser	Thr	Lys	Asp	Leu	Leu	Ser
				185					190					195
Gln	Trp	Glu	Ile	Phe	Pro	Pro	Gln	Ser	Trp	Lys	Leu	Ala	Leu	Val
				200					205					210
Ala	Ala	Met	Met	Ser	Gly	Ile	Ala	Val	Val	Leu	Ala	Met	Ala	Pro
				215					220					225
Phe	Asp	Val	Ala	Cys	Thr	Arg	Leu	Tyr	Asn	Gln	Pro	Thr	Asp	Ala

Gln Gly Lys Gly	230	Leu Met Tyr Arg Gly	235	Ile Leu Asp Ala Leu	240
	245		250		255
Gln Thr Ala Arg	260	Thr Glu Gly Ile Phe	265	Gly Met Tyr Lys Gly	270
	275		280		285
Gly Ala Ser Tyr	290	Phe Arg Leu Gly Pro	295	His Thr Ile Leu Ser	300
Phe Phe Trp Asp		Gln Leu Arg Ser Leu		Tyr Tyr Thr Asp Thr	

<210> 35  
 <211> 382  
 <212> PRT  
 <213> Homo sapiens  
  
 <220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2670745CD1  
  
 <400> 35  
 Met Leu Arg Trp Thr Val His Leu Glu Gly Gly Pro Arg Arg Val  
 1 5 10 15  
 Asn His Ala Ala Val Ala Val Gly His Arg Val Tyr Ser Phe Gly  
 20 25 30  
 Gly Tyr Cys Ser Gly Glu Asp Tyr Glu Thr Leu Arg Gln Ile Asp  
 35 40 45  
 Val His Ile Phe Asn Ala Val Ser Leu Arg Trp Thr Lys Leu Pro  
 50 55 60  
 Pro Val Lys Ser Ala Ile Arg Gly Gln Ala Pro Val Val Pro Tyr  
 65 70 75  
 Met Arg Tyr Gly His Ser Thr Val Leu Ile Asp Asp Thr Val Leu  
 80 85 90  
 Leu Trp Gly Gly Arg Asn Asp Thr Glu Gly Ala Cys Asn Val Leu  
 95 100 105  
 Tyr Ala Phe Asp Val Asn Thr His Lys Trp Phe Thr Pro Arg Val  
 110 115 120  
 Ser Gly Thr Val Pro Gly Ala Arg Asp Gly His Ser Ala Cys Val  
 125 130 135  
 Leu Gly Lys Ile Met Tyr Ile Phe Gly Gly Tyr Glu Gln Gln Ala  
 140 145 150  
 Asp Cys Phe Ser Asn Asp Ile His Lys Leu Asp Thr Ser Thr Met  
 155 160 165  
 Thr Trp Thr Leu Ile Cys Thr Lys Gly Ser Pro Ala Arg Trp Arg  
 170 175 180  
 Asp Phe His Ser Ala Thr Met Leu Gly Ser His Met Tyr Val Phe  
 185 190 195  
 Gly Gly Arg Ala Asp Arg Phe Gly Pro Phe His Ser Asn Asn Glu  
 200 205 210  
 Ile Tyr Cys Asn Arg Ile Arg Val Phe Asp Thr Arg Thr Glu Ala  
 215 220 225  
 Trp Leu Asp Cys Pro Pro Thr Pro Val Leu Pro Glu Gly Arg Arg  
 230 235 240  
 Ser His Ser Ala Phe Gly Tyr Asn Gly Glu Leu Tyr Ile Phe Gly  
 245 250 255  
 Gly Tyr Asn Ala Arg Leu Asn Arg His Phe His Asp Leu Trp Lys  
 260 265 270  
 Phe Asn Pro Val Ser Phe Thr Trp Lys Lys Ile Glu Pro Lys Gly  
 275 280 285  
 Lys Gly Pro Cys Pro Arg Arg Arg Gln Cys Cys Cys Ile Val Gly  
 290 295 300  
 Asp Lys Ile Val Leu Phe Gly Gly Thr Ser Pro Ser Pro Glu Glu  
 305 310 315  
 Gly Leu Gly Asp Glu Phe Asp Leu Ile Asp His Ser Asp Leu His  
 320 325 330  
 Ile Leu Asp Phe Ser Pro Ser Leu Lys Thr Leu Cys Lys Leu Ala  
 335 340 345  
 Val Ile Gln Tyr Asn Leu Asp Gln Ser Cys Leu Pro His Asp Ile  
 350 355 360

Arg Trp Glu Leu Asn Ala Met Thr Thr Asn Ser Asn Ile Ser Arg  
 365 370 375  
 Pro Ile Val Ser Ser His Gly  
 380

<210> 36  
 <211> 287  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2676443CD1

<400> 36  
 Met Ala Ala Glu Ala Arg Val Ser Arg Trp Tyr Phe Gly Gly Leu  
 1 5 10 15  
 Ala Ser Cys Gly Ala Ala Cys Cys Thr His Pro Leu Asp Leu Leu  
 20 25 30  
 Lys Val His Leu Gln Thr Gln Gln Glu Val Lys Leu Arg Met Thr  
 35 40 45  
 Gly Met Ala Leu Arg Val Val Arg Thr Asp Gly Ile Leu Ala Leu  
 50 55 60  
 Tyr Ser Gly Leu Ser Ala Ser Leu Cys Arg Gln Met Thr Tyr Ser  
 65 70 75  
 Leu Thr Arg Phe Ala Ile Tyr Glu Thr Val Arg Asp Arg Val Ala  
 80 85 90  
 Lys Gly Ser Gln Gly Pro Leu Pro Phe His Glu Lys Val Leu Leu  
 95 100 105  
 Gly Ser Val Ser Gly Leu Ala Gly Gly Phe Val Gly Thr Pro Ala  
 110 115 120  
 Asp Leu Val Asn Val Arg Met Gln Asn Asp Val Lys Leu Pro Gln  
 125 130 135  
 Gly Gln Arg Arg Asn Tyr Ala His Ala Leu Asp Gly Leu Tyr Arg  
 140 145 150  
 Val Ala Arg Glu Glu Gly Leu Arg Arg Leu Phe Ser Gly Ala Thr  
 155 160 165  
 Met Ala Ser Ser Arg Gly Ala Leu Val Thr Val Gly Gln Leu Ser  
 170 175 180  
 Cys Tyr Asp Gln Ala Lys Gln Leu Val Leu Ser Thr Gly Tyr Leu  
 185 190 195  
 Ser Asp Asn Ile Phe Thr His Phe Val Ala Ser Phe Ile Ala Gly  
 200 205 210  
 Gly Cys Ala Thr Phe Leu Cys Gln Pro Leu Asp Val Leu Lys Thr  
 215 220 225  
 Arg Leu Met Asn Ser Lys Gly Glu Tyr Gln Gly Val Phe His Cys  
 230 235 240  
 Ala Val Glu Thr Ala Lys Leu Gly Pro Leu Ala Phe Tyr Lys Gly  
 245 250 255  
 Leu Val Pro Ala Gly Ile Arg Leu Ile Pro His Thr Val Leu Thr  
 260 265 270  
 Phe Val Phe Leu Glu Gln Leu Arg Lys Asn Phe Gly Ile Lys Val  
 275 280 285  
 Pro Ser

<210> 37  
 <211> 497  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3295764CD1

<400> 37  
 Met Asp Val Pro Gly Pro Val Ser Arg Arg Ala Ala Ala Ala Ala  
 1 5 10 15  
 Ala Thr Val Leu Leu Arg Thr Ala Arg Val Arg Arg Glu Cys Trp  
 20 25 30

Phe	Leu	Pro	Thr	Ala	Leu	Leu	Cys	Ala	Tyr	Gly	Phe	Phe	Ala	Ser
				35					40					45
Leu	Arg	Pro	Ser	Glu	Pro	Phe	Leu	Thr	Pro	Tyr	Leu	Leu	Gly	Pro
				50					55					60
Asp	Lys	Asn	Leu	Thr	Glu	Arg	Glu	Val	Phe	Asn	Glu	Ile	Tyr	Pro
				65					70					75
Val	Trp	Thr	Tyr	Ser	Tyr	Leu	Val	Leu	Leu	Phe	Pro	Val	Phe	Leu
				80					85					90
Ala	Thr	Asp	Tyr	Leu	Arg	Tyr	Lys	Pro	Val	Val	Leu	Leu	Gln	Gly
				95					100					105
Leu	Ser	Leu	Ile	Val	Thr	Trp	Phe	Met	Leu	Leu	Tyr	Ala	Gln	Gly
				110					115					120
Leu	Leu	Ala	Ile	Gln	Phe	Leu	Glu	Phe	Phe	Tyr	Gly	Ile	Ala	Thr
				125					130					135
Ala	Thr	Glu	Ile	Ala	Tyr	Tyr	Ser	Tyr	Ile	Tyr	Ser	Val	Val	Asp
				140					145					150
Leu	Gly	Met	Tyr	Gln	Lys	Val	Thr	Ser	Tyr	Cys	Arg	Ser	Ala	Thr
				155					160					165
Leu	Val	Gly	Phe	Thr	Val	Gly	Ser	Val	Leu	Gly	Gln	Ile	Leu	Val
				170					175					180
Ser	Val	Ala	Gly	Trp	Ser	Leu	Phe	Ser	Leu	Asn	Val	Ile	Ser	Leu
				185					190					195
Thr	Cys	Val	Ser	Val	Ala	Phe	Ala	Val	Ala	Trp	Phe	Leu	Pro	Met
				200					205					210
Pro	Gln	Lys	Ser	Leu	Phe	Phe	His	His	Ile	Pro	Ser	Thr	Cys	Gln
				215					220					225
Arg	Val	Asn	Gly	Ile	Lys	Val	Gln	Asn	Gly	Gly	Ile	Val	Thr	Asp
				230					235					240
Thr	Pro	Ala	Ser	Asn	His	Leu	Pro	Gly	Trp	Glu	Asp	Ile	Glu	Ser
				245					250					255
Lys	Ile	Pro	Leu	Asn	Met	Glu	Glu	Pro	Pro	Val	Glu	Glu	Pro	Glu
				260					265					270
Pro	Lys	Pro	Asp	Arg	Leu	Leu	Val	Leu	Lys	Val	Leu	Trp	Asn	Asp
				275					280					285
Phe	Leu	Met	Cys	Tyr	Ser	Ser	Arg	Pro	Leu	Leu	Cys	Trp	Ser	Val
				290					295					300
Trp	Trp	Ala	Leu	Ser	Thr	Cys	Gly	Tyr	Phe	Gln	Val	Val	Asn	Tyr
				305					310					315
Thr	Gln	Gly	Leu	Trp	Glu	Lys	Val	Met	Pro	Ser	Arg	Tyr	Ala	Ala
				320					325					330
Ile	Tyr	Asn	Gly	Gly	Val	Glu	Ala	Val	Ser	Thr	Leu	Leu	Gly	Ala
				335					340					345
Val	Ala	Val	Phe	Ala	Val	Gly	Tyr	Ile	Lys	Ile	Ser	Trp	Ser	Thr
				350					355					360
Trp	Gly	Glu	Met	Thr	Leu	Ser	Leu	Phe	Ser	Leu	Leu	Ile	Ala	Ala
				365					370					375
Ala	Val	Tyr	Ile	Met	Asp	Thr	Val	Gly	Asn	Ile	Trp	Val	Cys	Tyr
				380					385					390
Ala	Ser	Tyr	Val	Val	Phe	Arg	Ile	Ile	Tyr	Met	Leu	Leu	Ile	Thr
				395					400					405
Ile	Ala	Thr	Phe	Gln	Ile	Ala	Ala	Asn	Leu	Ser	Met	Glu	Arg	Tyr
				410					415					420
Ala	Leu	Val	Phe	Gly	Val	Asn	Thr	Phe	Ile	Ala	Leu	Ala	Leu	Gln
				425					430					435
Thr	Leu	Leu	Thr	Leu	Ile	Val	Val	Asp	Ala	Ser	Gly	Leu	Gly	Leu
				440					445					450
Glu	Ile	Thr	Thr	Gln	Phe	Leu	Ile	Tyr	Ala	Ser	Tyr	Phe	Ala	Leu
				455					460					465
Ile	Ala	Val	Val	Phe	Leu	Ala	Ser	Gly	Ala	Val	Ser	Val	Met	Lys
				470					475					480
Lys	Cys	Arg	Lys	Leu	Glu	Asp	Pro	Gln	Ser	Ser	Ser	Gln	Val	Thr
				485					490					495
Thr	Ser													

<210> 38  
 <211> 228  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3438320CD1

<400> 38  
 Met Pro Arg Arg Gly Leu Val Ala Gly Pro Asp Leu Glu Tyr Phe  
 1 5 10 15  
 Gln Arg Arg Tyr Phe Thr Pro Ala Glu Val Ala Gln His Asn Arg  
 20 25 30  
 Pro Glu Asp Leu Trp Val Ser Tyr Leu Gly Arg Val Tyr Asp Leu  
 35 40 45  
 Thr Ser Leu Ala Gln Glu Tyr Lys Gly Asn Leu Leu Leu Lys Pro  
 50 55 60  
 Ile Val Glu Val Ala Gly Gln Asp Ile Ser His Trp Phe Asp Pro  
 65 70 75  
 Lys Thr Arg Asp Ile Arg Lys His Ile Asp Pro Leu Thr Gly Cys  
 80 85 90  
 Leu Arg Tyr Cys Thr Pro Arg Gly Arg Phe Val His Val Pro Pro  
 95 100 105  
 Gln Leu Pro Cys Ser Asp Trp Ala Asn Asp Phe Gly Lys Pro Trp  
 110 115 120  
 Trp Gln Gly Ser Tyr Glu Val Gly Arg Leu Ser Ala Lys Thr  
 125 130 135  
 Arg Ser Ile Arg Ile Ile Asn Thr Leu Thr Ser Gln Glu His Thr  
 140 145 150  
 Leu Glu Val Gly Val Leu Glu Ser Ile Trp Glu Ile Leu His Arg  
 155 160 165  
 Tyr Leu Pro Tyr Asn Ser His Ala Ala Ser Tyr Thr Trp Lys Tyr  
 170 175 180  
 Glu Gly Lys Asn Leu Asn Met Asp Phe Thr Leu Glu Glu Asn Gly  
 185 190 195  
 Ile Arg Asp Glu Glu Glu Phe Asp Tyr Leu Ser Met Asp Gly  
 200 205 210  
 Thr Leu His Thr Pro Ala Ile Leu Leu Tyr Phe Asn Asp Asp Leu  
 215 220 225  
 Thr Glu Leu

<210> 39  
 <211> 273  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3986488CD1

<400> 39  
 Met Ala Ala Thr Ile Met Ile Leu Tyr Val Ser Lys Leu Asn Lys  
 1 5 10 15  
 Ile Ile His Phe Pro Asp Phe Asp Lys Lys Ile Pro Val Lys Leu  
 20 25 30  
 Phe Pro Leu Pro Leu Leu Tyr Val Gly Asn His Ile Ser Gly Leu  
 35 40 45  
 Ser Ser Thr Ser Lys Leu Ser Leu Pro Met Phe Thr Val Leu Arg  
 50 55 60  
 Lys Phe Thr Ile Pro Leu Thr Leu Leu Leu Glu Thr Ile Ile Leu  
 65 70 75  
 Gly Lys Gln Tyr Ser Leu Asn Ile Ile Leu Ser Val Phe Ala Ile  
 80 85 90  
 Ile Leu Gly Ala Phe Ile Ala Ala Gly Ser Asp Leu Ala Phe Asn  
 95 100 105  
 Leu Glu Gly Tyr Ile Phe Val Phe Leu Asn Asp Ile Phe Thr Ala  
 110 115 120  
 Ala Asn Gly Val Tyr Thr Lys Gln Lys Met Asp Pro Lys Glu Leu  
 125 130 135  
 Gly Lys Tyr Gly Val Leu Phe Tyr Asn Ala Cys Phe Met Ile Ile  
 140 145 150  
 Pro Thr Leu Ile Ile Ser Val Ser Thr Gly Asp Leu Gln Gln Ala

Thr	Glu	Phe	Asn	155	Gln	Trp	Lys	Asn	Val	160	Val	Phe	Ile	Leu	Gln	165
Leu	Leu	Ser	Cys	170	Phe	Leu	Gly	Phe	Leu	175	Leu	Met	Tyr	Ser	Thr	180
Leu	Cys	Ser	Tyr	185	Tyr	Asn	Ser	Ala	Leu	190	Thr	Thr	Ala	Val	Val	195
Ala	Ile	Lys	Asn	200	Val	Ser	Val	Ala	Tyr	205	Ile	Gly	Ile	Leu	Ile	210
Gly	Asp	Tyr	Ile	215	Phe	Ser	Leu	Leu	Asn	220	Phe	Val	Gly	Leu	Asn	225
Cys	Met	Ala	Gly	230	Gly	Leu	Arg	Tyr	Ser	235	Phe	Leu	Thr	Leu	Ser	240
Gln	Leu	Lys	Pro	245	Lys	Pro	Val	Gly	Glu	250	Glu	Asn	Ile	Cys	Leu	255
Leu	Lys	Ser		260						265						270

<210> 40  
 <211> 206  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 4378816CD1

Met	Gly	Ala	Glu	Trp	Glu	Leu	Gly	Ala	Glu	Ala	Gly	Gly	Ser	Leu	
1				5					10					15	
Leu	Leu	Cys	Ala	Ala	Leu	Leu	Ala	Ala	Gly	Cys	Ala	Leu	Gly	Leu	
				20					25					30	
Arg	Leu	Gly	Arg	Gly	Gln	Gly	Ala	Ala	Asp	Arg	Gly	Ala	Leu	Ile	
				35					40					45	
Trp	Leu	Cys	Tyr	Asp	Ala	Leu	Val	His	Phe	Ala	Leu	Glu	Gly	Pro	
				50					55					60	
Phe	Val	Tyr	Leu	Ser	Leu	Val	Gly	Asn	Val	Ala	Asn	Ser	Asp	Gly	
				65					70					75	
Leu	Ile	Ala	Ser	Leu	Trp	Lys	Glu	Tyr	Gly	Lys	Ala	Asp	Ala	Arg	
				80					85					90	
Trp	Val	Tyr	Phe	Asp	Pro	Thr	Ile	Val	Ser	Val	Glu	Ile	Leu	Thr	
				95					100					105	
Val	Ala	Leu	Asp	Gly	Ser	Leu	Ala	Leu	Phe	Leu	Ile	Tyr	Ala	Ile	
				110					115					120	
Val	Lys	Glu	Lys	Tyr	Tyr	Arg	His	Phe	Leu	Gln	Ile	Thr	Leu	Cys	
				125					130					135	
Val	Cys	Glu	Leu	Tyr	Gly	Cys	Trp	Met	Thr	Phe	Leu	Pro	Glu	Trp	
				140					145					150	
Leu	Thr	Arg	Ser	Pro	Asn	Leu	Asn	Thr	Ser	Asn	Trp	Leu	Tyr	Cys	
				155					160					165	
Trp	Leu	Tyr	Leu	Phe	Phe	Phe	Asn	Gly	Val	Trp	Val	Leu	Ile	Pro	
				170					175					180	
Gly	Leu	Leu	Leu	Trp	Gln	Ser	Trp	Leu	Glu	Leu	Lys	Lys	Met	His	
				185					190					195	
Gln	Lys	Glu	Thr	Ser	Ser	Val	Lys	Lys	Phe	Gln					
				200					205						

<210> 41  
 <211> 235  
 <212> PRT  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 4797137CD1

Met	Gln	Gln	Arg	Gly	Ala	Ala	Gly	Ser	Arg	Gly	Cys	Ala	Leu	Phe	
1				5					10					15	
Pro	Leu	Leu	Gly	Val	Leu	Phe	Phe	Gln	Gly	Val	Tyr	Ile	Val	Phe	

	20		25		30
Ser	Leu	Glu	Ile	Arg	Ala
	35		40		45
Glu	Lys	Ile	Lys	Leu	Lys
	50		55		60
Thr	Asp	Lys	Leu	Thr	Ile
	65		70		75
Ser	His	Thr	Val	Ser	Ile
	80		85		90
Thr	Thr	Ala	Gly	Thr	Phe
	95		100		105
Val	Tyr	Lys	Gly	Asp	Ala
	110		115		120
Lys	Asp	Asn	Gly	Thr	Phe
	125		130		135
Val	His	His	Asn	Ile	Pro
	140		145		150
Gly	Phe	Gly	Thr	Met	Leu
	155		160		165
Val	Phe	Val	Pro	Ser	Ala
	170		175		180
Met	Gly	Arg	Lys	Ala	Ala
	185		190		195
Tyr	Lys	Lys	Ser	Ser	Ile
	200		205		210
Glu	Glu	Glu	Ala	Cys	Met
	215		220		225
Cys	Leu	Asp	Ser	Asp	Tyr
	230		235		

&lt;210&gt; 42

&lt;211&gt; 147

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5470806CD1

&lt;400&gt; 42

Met	Ala	Ser	Leu	Arg	Leu	Phe	Leu	Leu	Cys	Leu	Ala	Val	Leu	Ile
	1		5		10		15		20		25		30	
Phe	Ala	Ser	Glu	Ala	Gly	Pro	Gly	Gly	Ala	Gly	Glu	Ser	Lys	Cys
	35		40		45		50		55		60		65	
Pro	Leu	Met	Val	Lys	Val	Leu	Asp	Ala	Val	Arg	Gly	Ser	Pro	Ala
	70		75		80		85		90		95		100	
Val	Asp	Val	Ala	Val	Lys	Val	Phe	Lys	Lys	Thr	Ala	Asp	Gly	Ser
	105		110		115		120		125		130		135	
Trp	Glu	Pro	Phe	Ala	Ser	Gly	Lys	Thr	Ala	Glu	Ser	Gly	Glu	Leu
	140		145		150		155		160		165		170	
His	Gly	Leu	Thr	Thr	Asp	Glu	Lys	Phe	Thr	Glu	Gly	Val	Tyr	Arg
	175		180		185		190		195		200		205	
Val	Glu	Leu	Asp	Thr	Lys	Ser	Tyr	Trp	Lys	Ala	Leu	Gly	Ile	Ser
	210		215		220		225		230		235		240	
Pro	Phe	His	Glu	Tyr	Ala	Glu	Val	Val	Phe	Thr	Ala	Asn	Asp	Ser
	245		250		255		260		265		270		275	
Gly	His	Arg	His	Tyr	Thr	Ile	Ala	Ala	Leu	Leu	Ser	Pro	Tyr	Ser
	280		285		290		295		300		305		310	
Tyr	Ser	Thr	Thr	Ala	Val	Val	Ser	Asn	Pro	Gln	Asn			
	315		320		325		330		335		340		345	

&lt;210&gt; 43

&lt;211&gt; 147

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 5473242CD1



<400> 43  
 Met Val His Leu Thr Asp Ala Glu Lys Ala Thr Val Asn Gly Leu  
 1 5 10 15  
 Trp Gly Lys Val Asn Pro Val Glu Ile Gly Ala Glu Ser Leu Ala  
 20 25 30  
 Ser Leu Leu Ile Val Tyr Pro Trp Thr Gln Arg Tyr Phe Ser Lys  
 35 40 45  
 Phe Gly Asp Leu Ser Ser Val Ser Ala Ile Met Gly Asn Pro Gln  
 50 55 60  
 Val Lys Ala His Gly Glu Lys Val Ile Asn Ala Phe Asp Asp Gly  
 65 70 75  
 Leu Lys His Leu Asp Asn Leu Lys Gly Thr Phe Ala Ser Leu Ser  
 80 85 90  
 Glu Leu His Cys Asp Lys Leu His Val Asp Pro Glu Asn Phe Arg  
 95 100 105  
 Leu Leu Gly Asn Met Ile Val Ile Met Met Gly His His Leu Gly  
 110 115 120  
 Lys Glu Phe Thr Pro Ser Ala Gln Ala Ala Phe Gln Lys Val Val  
 125 130 135  
 Ala Gly Val Ala Ser Ala Leu Ala His Lys Tyr His  
 140 145

<210> 44

<211> 2701

<212> DNA

<213> Homo sapiens

<220>

<221> misc\_feature

<223> Incyte ID No: 264114CB1

<400> 44

gcggcgggcgc cagcttctctc ggccggaggag gaggcgagac cccaggcgag gccgcggcgcg 60  
 gaggggccacg ccccccacgc cgcgccggag gggcccagtg tggacggggc caccggctgg 120  
 agcggatccc acacctccgg accgagggac gcggttactc cacaggatcc gctgaacata 180  
 ggatgttggc acaaaatcta cctcgtgtat ttttctcttt cactcatgag ctgcacaatt 240  
 gcagatttga gcacaatgtc tgcagactgt gttgaaaaac tctgaagaac ctaattaaca 300  
 caggatgacc taggagtgt tctaagtctg tgtaacaaga tattactcat tagtgaatgt 360  
 gtcagtcttg gtactgaatg ctgcagataa cagcaagtag gttctccttt atttctgaag 420  
 tattcacttg accttccatc agtaagacgg acttttctaa tctgttcttg gagatattaa 480  
 tggaaatacag tcatgtccac tcaagacgag aggcagatca atactgaata tgctgtgtca 540  
 ttgttgggaac agttgaaact gttttatgaa cagcagttgt ttactgacat agtggttaatt 600  
 gttgaggggca ctgaattccc ttgtcataag atgggttctg caacatgtag ctcttatttc 660  
 agggccatgt ttatgagtgg actaagtga agcaaaacaaa cccatgtaca cctgaggaat 720  
 gtcgatgctg ccaccttaca gataataata acttatgcat acacgggtaa cttggcaatg 780  
 aatgacagca ctgtagaaca gctttatgaa acagcttgct tcctacaggt agaagatgtg 840  
 ttacaacggt gtcgagaata ttaattaaa aaaataaatg cagagaattg tgtacgattg 900  
 ttgagttttg ctgatctctt cagttgtgag gaattaaaac agagtgtctaa aagaatggtg 960  
 gagcacaagt tcaactgctg gtatcatcag gatgtagaaa aggaagaaac cgttcgagaa 1020  
 ctgatagata ttctcagtag tgacaattta aatgtagaaa aggaagaaac cgttcgagaa 1080  
 gctgctatgc tgtggctaga gtataacaca gaatcacgat cccagttatt gtcttctgtt 1140  
 cttagccaaa tcagaattga tgcactttca gaagtaacac agagagcttg gtttcaagg 1200  
 ctgccaccca atgataagtc agtggtggtt caagggtctg ataagtccat gcccaagtt 1260  
 ttcaaaccac gacttgggat gactaaagag gaaatgatga ttttcattga agcatcttca 1320  
 gaaaatcctt gtagtcttta ctcttctgtc tgttacagcc cccaagcaga aaaagtttac 1380  
 aagttatgta gccaccagc tgatttgcac aagggttgga ccgttgtaac tcctgataat 1440  
 gatattctaca tagcaggggg tcaagttcct ctgaaaaaca caaaaacaaa tcacagtaaa 1500  
 acaagcaaac ttcagactgc cttcagaact gtgaattgct tttattggtt tgatgcacag 1560  
 caaaatacct ggtttccaaa gaccccaatg ctttttgtcc gcataaagcc atctttggtt 1620  
 tgctgtgaag gctatatcta tgcaattgga ggagatagcg taggtggaga acttaacgg 1680  
 aggaccgtag aaagatacga cactgagaaa gatgagtga cgatggtaag ccctttacct 1740  
 tgtgcttggc aatggagtgc agcagttgtg gttcatgact gcatttatgt gatgacactg 1800  
 aacctcatgt actgttattt tccaaggtct gactcatggg tagaaatggc catgagacag 1860  
 actagtaggt cctttgcttc agctgcagct tttggtgata aaattttcta tattggagg 1920  
 ttgcatattg ctaccaattc cggcataaga ctcccctctg gcactgtaga tgggtcttca 1980  
 gtaactgtgg aaatttatga tgtgaataaa aatgagtga aaatggcagc caacatccct 2040  
 gctaagaggt actctgaccc ctgtgttaga gctgtgtgta tctcaaattc tctatgtgtg 2100  
 tttatgcgag aaaccactt aaatgagcga gctaaatcgc tcacctacca atatgacctg 2160  
 gaacttgacc ggtggtctct gcggcagcat atatctgaac gtgtactgtg ggacttgggg 2220

```

agagattttc gatgcactgt ggggaaactc tatccatcct gccttgaaga gtctccatgg 2280
aaaccaccaa cttatctttt ttcaacggat gggacagaag agtttgaact ggatggagaa 2340
atggttgcac taccacctgt atagtgggga agttcagggg gtgcacgcct gagttatgtg 2400
ctttgtcatt ttctttgcta aacaaaagag gctatgaaa aactaaatat gagtacataa 2460
aattctatct ttgataaatt ttatttttat gcctactta atatttgcac cagtataata 2520
tatatcagtg agtcctacag aaagatatgc ttccataata tgaaatagat tattcaataa 2580
ttgagaaact ttatgtgtaa tcatgagagt ataagaatct ggattatcta acattgttag 2640
ccctgtgtat gtacagttca aaaagtccat ttataaaagt agtttcctgt tcctagttag 2700
t

```

&lt;210&gt; 45

&lt;211&gt; 736

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 1455669CB1

&lt;400&gt; 45

```

gagacttagc gacagacaga cgctgggacc cagcagcaga gaaggcgccg atggccgcgc 60
ctgctgagcc ctgctcgagg caggggggtct ggaaccagac agagcctgaa cctgccgccca 120
ccagcctgct gagcctgtgc ttccctgagaa cagcaggggt ctgggtaccc cccatgtacc 180
tctgggtcct tgggtcccatc tacctcctct tcatccacca ccctggccgg ggctacctcc 240
ggatgtcccc actcttcaaa gccaagatgg tagctgccat ccctgggagc ctggaaccag 300
gcaatgttcg ggggaggcag gggacaggct ggaacctggt gaagtcttaa agtagactcc 360
tcctatcggg gtgtagaagg gaatctgtta atcaaacaga gcaatattag aaaggctaca 420
gaggtcaact cagtggaaaca cggttctccc aaacagattt tgtaattccg aaaatccacg 480
catgcgcaaa catacgcata cactcccatg ttctctggaca gtttatagct accataacct 540
ggcattttcc aaaacatacc atgtagactc ttggatacac aaggtaattt tagagccaca 600
ttaggatgaa ccttttaaaa agttatgcat ttatttttat gttccccac tggtgtatt 660
ataggacaat ttttatatgt gatatgtatt taccttagtg tgttaaataa aactggcat 720
tccaagtgtg aaaaaa

```

736

&lt;210&gt; 46

&lt;211&gt; 1826

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2084989CB1

&lt;400&gt; 46

```

ccttagggcg cagggacagc cgagcgttac ctgggtcccg gcagcggagt tctttaccca 60
ccccagttct ggttctgacg ccctagctca ttccgcaaat ttagggcttg ggtctggctt 120
gttccccctc ggctcgaacc acctcttctc tgagccgagc cagctaccgg ggctcctgga 180
attgccaccc ctccctgggc acccttgagg cctccgtgga gggacgtcac ggggcagagc 240
gggacgtgag cctgagtttg ctgcaggcgt gctctgtgtg gtggctgggt tctgccaatc 300
cccgtgcca cgggtgggc gcggccggga agctcctgcc cctccctgct ggctggcgtc 360
acgctgacg tcccgcgtga tggctgggag ggcccgccgg cgacagcgga ggcagagagg 420
aaggcggttc tgagagcttc agagagcgat ggaaagcaaa atgggtgaat tgcctttaga 480
catcaacatc caggaacctc gctgggacca aagtactttc ctgggcagag cccggcactt 540
tttactgtt actgatcctc gaaatctgct gctgtccggg gcacagctgg aagcttctcg 600
gaacatcgtg cagaactaca gggccggcgt ggtgacccca gggatcaccg aggaccagct 660
gtggaggggc aagtatgtgt atgactccgc cttccatccg gacacagggg agaaggtggg 720
cctgattggc cgcattgtcag cccaggtgcc catgaacatg accatcactg gctgcatgct 780
cacattctac aggaagaccc caaccgtggt gttctggcag tgggtgaatc agtccctcaa 840
tgccattgtt aactactcca accgcagtgg tgacactccc atcactgtga ggcagctggg 900
gacagcctat gtgagtgcca ccaactggag tgtggccacg gccctgggac tcaaatccct 960
caccaagcac ctgccccctt tggctggcag atttgtccc ttgacagcag tggcagctgc 1020
tgatgaggca ggtcagaggc ttggctactc ggtgactgca gccaaagcagg gaatcttcca 1140
ggtgggtgatt tcaagaatct gcattggcgt tcttgccatg gccatccac cactgatcat 1200
ggacactctg gagaagaaag acttcctgaa ggtaggcgac tgtacctctc ttgtcctgga 1260
atggggcagtg gctgggagaa gtgaccaggc cccaactctc tctccagcct cgctgtattc 1320
tctaagactt gccagccctt ctctgacct ctgcaccgcc tctccacct tcgttcattc 1380
agcaagaatg aactgggctg ggggtgaagg actctgcagg ggcaggagga gaggacaaa 1440

```

```

gaaggaaacc aacttcatca gtgttactcc agtggcttct gacacacaga aggggactgt 1500
catagtcatg cttgatctca tgctcattct tttacccctt agtgcctcca tactgagagg 1560
tacacacggg tgaacacgca cacacagaca tgaacaggac acgaaagcaa agcacaggaa 1620
caagctctgg ctcattcaca gaatcattta ttcacaaatg tattgagtgc catgcaccag 1680
gcatgtttta gggctgagga gatggcactg aacacaatgg ttatggcccc tgtcctcatg 1740
aagtttatag tctgatgcag aaaccaataa acaaggaggc acccacataa atacattctt 1800
agaaagtgtg aaaataaaaa aaaaaa

```

1826

&lt;210&gt; 47

&lt;211&gt; 1325

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2501034CB1

&lt;400&gt; 47

```

ccacgggtcc ggttctggac tgcagttgag tggaaatggg caacggcggg cggagcgggc 60
tgcagcaggg gaaggggaac gtggatgggg tggcagcgac tcctactgct gcctcggcct 120
cctgccagta caggtgcatc gaatgcaacc aggaggccaa agagtgtgtac cgagactata 180
accacgggtg gctgaagata accatctgta aatcctgcc aagaaacctgta gacaaatata 240
tcgagtatga tcctgttata atcttgatta atgctatatt gtgcaaagct caggcctaca 300
gacatattct tttcaatact caaataaata tccatggaaa actctgcata ttttgtttgc 360
tttgtgaagc atacctgagg tgggtggcagc ttcaagattc caaccagaat actgcccctg 420
atgacttgat cagatatgct aaggaatggg atttctatag aatgtttgcg attgctgctt 480
tagaacaacac tgcctatttt attggcattt ttaccttct gtgggtagaa cggccccatga 540
cggcaaaaaa aaagcccaac ttcattttgc tgctgaaagc attattatta tctagctacg 600
gaaaactctt gctgattcca gctgtcattt gggaaacatga ctacacatct gtgtgcctca 660
aactcattaa agtatttggc cttacatcaa attttcaggc aattagagtg accctaaaca 720
tcaaccgtaa gctctccttc ttggccgtgt tgagtggcct actgctggaa agcatcatgg 780
tctacttctt ccagagtatg gaatgggatg ttggaagtga ttatgccatc tttaaatctc 840
aggacttctg aagagtttta ttcttcttca ctatctgtgg catgaccagc tgtatctgaa 900
agagaaaaga catgaaatat aaaccaacct cctcatttct gttgagttaa atgaagcaaa 960
gattggaaac actttctgaa aaagaaagca atgataatag cgggtggatac ccacccccac 1020
aatgacaccc aagagacaag ccattttacat acagatattc acagtccac atagaaacac 1080
ccacatggag acaaggaatg ttgctgcaga gactgaatga catgcaacag gtgaagggtt 1140
atacgttata cacaaggcca ggtaagcgct cataattcac acataataaa acatctaggt 1200
ttcattcctt tgacatgttt atatcttttt aatttaaatg ttgttactgg cttaaaatat 1260
tttgtgttct tacaatagaa acgcttttaa taaagtcttt cagaataaac caaaaaaaa 1320
aaaaa

```

1325

&lt;210&gt; 48

&lt;211&gt; 1832

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2745212CB1

&lt;400&gt; 48

```

tgggctgtcg ttggctggag cagcggctgc gcggtgcggt gtgctgtgag gtctgcgggc 60
gctggcaaat ccggcccagg atgtagagct ggcagtgccg gacggcgcgct ctgacgcgga 120
gttgggtggg gtagagagta gggggcggtg gtcgggggtg gtgggagaag gaggaggcgg 180
cgaatcactt ataaatggcg ccgaagcagg acccgaagcc taaattccag gagggtgagc 240
gagtgtgtgt ctttcatggg cctcttcttt atgaagcaaa gtgtgtaaa gttgccataa 300
aggacaaaca agtgaataac ttcatacatt acagtgggtg gaataaaaaat tgggatgaat 360
gggttccgga gagcagagta ctcaaatacg tggacaccaa tttgcagaaa cagcgagaac 420
ttcaaaaagc caatcaggag cagtatgcag aggggaagat gagaggggct gccccaggaa 480
agaagacatc tggctctgca cagaaaaatg ttgaagtga aacgaaaaag aacaaacaga 540
aaacacctgg aatggagat ggtggcagta ccagtggagc cctcagcct cctcggaaga 600
aaaggggccc ggtagatcct actgttgaaa atgaggaaac attcatgaac agagtgaag 660
ttaaagttaa gattcctgaa gagctaaaac cgtggcttgt tgatgactgg gacttaatta 720
ccaggcaaaa acagctcttt tatcttctgt ccaagaagaa tgtggattcc attcttgagg 780
attatgcaaa ttacaagaaa tctcgtggaa acacagataa taaggagtat gcggttaatg 840
aagttgtggc agggataaaa gaatacttca acgtaatgtt gggtagccag ctactctata 900
aatttgagag accacagtat gctgaaattc ttgcagatca tcccagtgca cccatgtccc 960

```

```

agggtgatgg agcgccacat ctcttgagat tatttgtacg aattggagca atgttggcctt 1020
atacacctct ggatgagaag agccttgctt tattactcaa ttatcttcac gatttcctaa 1080
agtacctggc aaagaattct gcaactttgt tcagtgccag cgattatgaa gtggctcctc 1140
ctgagtacca tcggaaagct gtgtgagagg cactctcact cacttatgtt tggatctccg 1200
taaacacatt tttgttctta gtctatctct tgtacaaacg atgtgctttg aagatgttag 1260
tgtataacaa ttgatgtttg ttttctgttt gattttaaac agagaaaaaa taaaaggggg 1320
taatagctcc tttttcttc tttctttttt tttttcattt caaaattgct gccagtgttt 1380
tcaatgatgg acaacagagg gatatgctgt agagtgtttt attgcctagt tgacaaagct 1440
gcttttgaat gctgggtggt ctattccttt gacactacgc acttttataa tacatgttaa 1500
tgctatatga caaaatgctc tgattcctag tgccaaaggt tcaattcagt gtatataact 1560
gaacacactc atccatttgt gcttttgttt ttttttatgg tgcttaaagt aaagagccca 1620
tcctttgcaa gtcattccatg ttgttactta ggcattttat cttggctcaa attgttgaag 1680
aatgggtggc tgtttcatgg tttttgtatt tgtgtctaag gcacgtttta acatgataga 1740
cgcaatgcat tgtgtagcta gttttctgga aaagtcaatc ttttaggaat tgtttttcag 1800
atcttcaata aattttttct ttaattttca aa
1832

```

&lt;210&gt; 49

&lt;211&gt; 1211

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 4833111CB1

&lt;400&gt; 49

```

gacagacccg cctcaaacat ggcgggcgccc agcgcgcgag gacgtgatcc gcttctgctc 60
cggtctggat tgtagccttg acgaggtctg agcgaccatg gaccggccgg gggtcgtggc 120
agcgctgggtg gctgggtggg tagcaggtgt ttctgttgac ttgatattat ttctctgga 180
taccattaaa accaggctgc agagtcccca aggatttagt aaggctgggtg gttttcatgg 240
aatatatgct ggcgttccct ctgctgctat tggatccctt cctaattgctg ctgcattttt 300
tatcacctat gaatatgtga agtgggtttt gcatgctgat tcatcttcgt atttgacacc 360
tatgaaacat atgttggctg cctctgctgg agaagtgtt gcctgcctga ttcgagttcc 420
atctgaagtg gttaagcaga gggcacagg atctgcttct acaagaacat ttcagatttt 480
ctctaacatc ttatatgaag aggggtatcca aggggtgtat cgaggctata aaagcacagt 540
tttaagagag attccttttt ctttgggtcca gtttccctta tgggagtcct taaaagccct 600
ctggctcctgg aggcaggatc atgtgggtgga ttcttggcag tcagcagctc gtggagcttt 660
tgcaaggtgga ttggccgctg cagtcaccac cctctagag gtggcaaga caagaattac 720
gctggcaaaag gctggctcca gcactgctga tgggaatgtg ctctctgtcc tgcatgggg 780
ctggcggtca caggggctgg caggattatt tgcaggtgtc ttccctcgaa tggcagccat 840
cagtcctggga ggtttcatct ttctgggggc ttatgaccga acgcacagct tgctgttgg 900
agtgggcaga aagagtcctt gaagcagaga caagcctcac ctccacttct gtcaagagag 960
gggcctgcag tgcaaaccct ctccgctga gcagctgtct gaactatagg cccagtgct 1020
gaagaccagt tgtgctaaga taccggcatg gagattgtgc catccgtgg ataggctggc 1080
tggtatgaag tcattggcct gtatgccaga gagctaagag aagaaaacgg ggtctgtggc 1140
ggtactctga acaatttctt cagaacctct taataaataa gtttggtaat gctgagaaaa 1200
aaaaaaaaa a
1211

```

&lt;210&gt; 50

&lt;211&gt; 1046

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 876677CB1

&lt;400&gt; 50

```

cccacgcgtc cgggaatgtc tttcactctc tcatactttc tcctctcccc tctcccaagc 60
acatctgagt tgcgcctgt tcttcacact tagctccaaa cccatgaaaa attgccaagt 120
ataaaaagctt ctcaagaatg agatggatc taggggtgtc tcacctgaga agcaagataa 180
agagaatttc gtgggtgtca acaataaacg gcttgggtga tgtggctgga tctgttttc 240
cctctctttc ctgttgggtga tcattacctt ccccatctcc atatggatgt gcttgaagat 300
cattaaggag tatgaacgtg ctgttgtatt ccgtctggga cgcattccaag ctgacaaagc 360
caagggggcca ggtttgatcc tggtcctgcc atgcatagat gtgtttgtca aagttgacct 420
ccgaacagtt acttgaaca ttctccaca agagatctc accagagact ccgtaactac 480
tcaggtagat ggagttgtct attacagaat ctatagtgtc gtctcagcag tggctaagt 540
caacgatgtc catcaagcaa catttctgct gggtcaaac actctagaa atgtcttag 600

```

```

gacacagacc ttgtcccaga tcttagctgg acgagaagag atcgcccata gcatccagac 660
tttactttgat gatgccaccg aactgtgggg gatccgggtg gcccagtggtg aaatcaaaga 720
tggttcggatt cccgtgcagt tgcagagatc catggcagcc gaggtctgagg ccaccgggga 780
agcgagagcc aaggtccttg cagctgaagg agaaatgaat gcttccaaat ccctgaagtc 840
agcctccatg gtgctggctg agtctcccat agtctctcag ctgctgctacc tgcagacctt 900
gagcacggta gccaccgaga agaattctac gattgtgttt cctctgcccc tgaatatact 960
agagggcatt ggtggcgctc gctatgataa ccacaagaag cttccaaata aagcctgagg 1020
tcctcttgcg gtagtcagct attgca

```

1046

&lt;210&gt; 51

&lt;211&gt; 1660

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2326143CB1

&lt;400&gt; 51

```

gctccccctgc ccaccccgcc cccgtggccg agcccgggag tgcagtgagg gtcggccggc 60
cggcgccgggc agccgtgacc cccgcggggg aactgcagc cggagcccgg gaggggcccgc 120
gccgccaccg tctgaactag gatgtcccga catgaagggtg tcagctgtga tgcagtgttta 180
aaaggaaaatt ttcgaggtcg cagatataag tggttaattt gctacgatta cgatctttgt 240
gcatcttggtt atgaaagtgg tgcaacaaca acaaggcata caactgacca cccaatgca 300
tgcatatttaa caagggtaga ttttgattta tactatgggtg gggaagcttt ctctgtagag 360
cagccacagt cttttacttg tccctattgt ggaaaaatgg gctatacggg gacatctctt 420
caagaacatg ttacttctga acatgcagaa acatcaacag aagtgttttg tccaatatgt 480
gcagcgttac ctggaggcga tcctaatacat gtcacggatg actttgcagc tcatcttaca 540
cttgaacaca gagcccctag agattttgat gaatcgagtg gtgttcgaca tgtacgtaga 600
atgtttcacc ctggccgggg attaggaggt cctcgtgctc gtagatcaaa catgcacttt 660
actagcagtt ctactgttgg actttcttct tctcagagtt catattctcc aagcaatagg 720
gaagccatgg atcctatagc tgagctttta tctcagttat caggagtggg acgttctgca 780
ggaggacagc ttaattcctc tggcccttcc gcttctcagt tacaacaact gcagatgcag 840
ctgcagctag aacggcagca tgcccaggga gcacggcaac aactggagac cgcacgcaac 900
gcaacccggc gtactaacac aagcagtgtc accactacaa tcacacaatc cacagcaaca 960
accaacatag ctaatacaga aagcagtcag cagactctac agaattccca gtttctttta 1020
acaaggttga atgatcctaa aatgtctgaa acggagcgcc cttctgtcca ctttagtgcg tgaagagagc 1140
gcagaccgca gctgtttgtt ccaagagctc atggcagatt ttggtgctat gggctgtgta 1200
tcatctcag atgaggatga tcggggggag agctctgctc taaaagagag taataaagga 1260
gatattatgc ctttagatgt tgcttttagaa aacctaaatt taaaagagag gtcctctgtg 1320
aatgagcctc caccacctcc tctttgatga catcccaatt cgtgttgggt ttgtttgggt attgtaattt 1380
ctgtatttgc caatgaaagt ggacaacaac tatcttgggt attcaaaaag aagagagaaa atatatatga 1440
caggtctgtc actcttgtaa cattgtgtac attcaaaaag aagagagaaa atatatatga 1440
taatcatttc cacttaacta atttttactt ctagcaggta aatgtaggta gcagtgcagg 1500
ggtgatctct gcttctgta ccttgacatg caaaaggctc tcctaatact ccacattcaa 1560
actgaagagg aaaattgaaa tctctaatag agctgctgtg tgtatttatg aatattaatg 1620
aataaaaact gcttggatgg tttaccttaa aaaaaaaaaa

```

1660

&lt;210&gt; 52

&lt;211&gt; 1110

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2786302CB1

&lt;400&gt; 52

```

cctttattag agaaagagtg acctatctac cccttacttg gggaagactc tctgcttcaa 60
cagtttcttc tgtcttctgc tcttgacat ggcatgaaaa cggaagtgtg agtatttcaa 120
ggataaagaa taactctagg acaagagttg ctttggggct aaacctatct cctgtattca 180
tttagatttg gctttaaaag ccatactat tttattagct cttctatgct ttttaaaagt 240
ttttaaaaaa attaatgtgg gctttttaat tctttctcag tgggagggtg agtttgaata 300
aacctttctt ccacatgaga agtattttac aagtgtcttg tcaaatttaa aagaaaatga 360
tcaaattctt aagaaaatga tcaaattctt tgtgacaaaa aaatggacaa atattcacca 420
ttgagtgtga atgccatggt gatgctaatt ctgattattt ttcttttctt ttaggccata 480
cttcagagtg ttattttcct ttttgcata aggtgtctgg aaatgaagta tggaaatgaa 540
ataatgaata aagaccagct tttcagaatc tctccacgga gtagagaaac tcatcccaat 600

```

```

ccggaagagc ccgaagaaga agatgaagat gttcaagctg aaagagtcca agcagcaaat 660
gcactcactg ctccaaactt ggaggaggaa ccagtcataa ctgcaagctg tttacacaag 720
gaatattatg agacaaagaa aagttgcttt tcaacaagaa agaagaaaat agccatcaga 780
aatgtttcct tttgtgttaa aaaaggtgaa gttttgggat tactaggaca caatggagct 840
ggtaaaagta cttccattaa aatgataact ggggtgcacaa agccaactgc aggagtgggtg 900
gtgttacaag gcagcagagc atcagtaagg caacagcatg acaacagcct caagtctctg 960
gggtactgcc ctgaggagaa ctgactgtgg cccaagctta ccatgaaaga gcacttggag 1020
ttgtatgcag ctgtggaaag actggggcaa aaaagatgct gctctcagta tttcacgatt 1080
ggtgggaggt cttaagctcc aggaacaact
1110

```

<210> 53  
 <211> 1120  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3735780CB1

```

<400> 53
gaaatccagt tatcaaaatt gactcaagaa gagagaacct aacagaacaa taacaatgga 60
agaaattggg aacattatca caaagctatc atcctgccaa actccaggct cagatgtcac 120
aggttaaaaa aagtccttca tgaaaaagaa agatcttaag cagcatgatg gattcagaag 180
ctcatgaaaa gaggccacca atactaacat cttcaaaaca agatatatca cctcatatta 240
caaagtgttg tgagatgaag cattacttgt gtggctgctg tgcagccttc aacaatgtcg 300
caatcacatt tcccattcag aaggtcctct ttcgacaaca gctgtatggc atcaaaaccc 360
gggatgcaat acttcagttg agaagggatg gatttcgaaa tttgtatcgt ggaatccttc 420
ccccattgat gcagaagaca actacgcttg cacttatgtt tggctctgat gaggatttat 480
ctgccttctt ccacaagcat gtcagtgtct cagagtttgc aaccagtggc gtggcggcag 540
tgcttgcagg gacaacagaa gcaattttca ctccactgga aagagttcag acattgcttc 600
aagaccacaa acatcatgac aaattttacca acacttacca ggctttcaag gcactgaaat 660
gtcatggaat tggagagtat tatcgaggct tgggtgccat tcttttcggg aatggactca 720
gcaatgtctt gtttttcggc cttcgaggct ccattaagga gcatctgcct accgcaacga 780
ctcacagtgc tcatctggtc aatgatttta tctgtggagg tctattgggt gccatgttgg 840
gattcttgtt ttttccaatt aatgttgtta aaactcgcat acagtctcag attgggtggg 900
aatttcagtc tttccccaag gttttccaaa aaatctggct ggaacgggac agaaaactga 960
taaattcttt cagaggtgcc catctgaatt accatcggtc cctcatctct tggggcataa 1020
tcaatgcaac ttatgagttc ttgttaagg ttatatgaaa aaaccatcag ttaagtgcga 1080
tttatcaact gaatagacct tctaagaaga aaaaaaaaaa
1120

```

<210> 54  
 <211> 886  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 039026CB1

```

<400> 54
ggccgcggct cctgtccaga ccctgaccct ccctcccaag gctcaaccgt cccccaacaa 60
ccgccagcct tgtactgatg tcggatgcga gagcctgtgc ttaagtaaga atcaggcctt 120
attggagaca ttcaagcaaa gggtggacaa ctacttttcc agaacagaaa ggaaactcat 180
gcatcagaaa aggtgactaa taaaggtacc agaagaatat ggctgcacaa ataccagaat 240
ctgatcagat aaaacagttt aaggaatttc tggggaccta caataaactt acagagacct 300
gctttttgga ctgtgtttaa gacttcacaa caagagaagt aaaacctgaa gagaccacct 360
gttcagaaca ttgtttacag aaatatttaa aaatgacaca aagaatatcc atgagatttc 420
aggaatatca tattcagcag aatgaagccc tggcagccaa agcaggactc cttggccaac 480
cacgatagag aagtcctgat ggtatgaactt ttgatgaaa attgccaaca gctgctttat 540
tggaatgag gactcatctg atagaatccc ctgaaagcag tagccaccat gttcaaccat 600
ctgtcatgac tgtttggcaa atggaaaccc ctggagaaac aaaattgcta ttaccagga 660
ataatcacia tagaaggtct tattgttcag tgaaataata agatgcaaca tttgttgagg 720
ccttatgatt cagcagcttg gtcacttgat tagaaaaata aaccattgtt tcttcaattg 780
tgactgttaa ttttaaagca acttatgtgt tcgatcatgt atgagataga aaaattttta 840
ttactcaaag taaaataaat ggaaatatca ctgaaaaaaa aaaaaa
886

```

<210> 55  
 <211> 2336

<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 260607CB1

<400> 55

taatacgcctc	actatagggg	atctggccct	cgagcagtaa	ttcggcacga	ggaccatctc	60
tttaggatat	atcttttaaat	tctttgaaac	acataaccaa	aatgggttga	ttcactgact	120
gactttgaaag	ctgcatctgc	cagttacacc	ccaaatggct	ttaatccccct	ctcgggtctg	180
gttgccctttt	gcagtttggg	ttgtggactc	agctcctgtg	aggggtctgg	ttaggagaga	240
gccattttta	aggacaggga	gttttatagc	ccttttctac	tttcctcccc	tcctcccagt	300
ccttatcaat	cttttttccct	ttttcctgac	ccccctcttc	tggaggcagt	tgggagctat	360
ccttgtttat	gcctcactat	tggcagaaaa	gaccccatct	aaaacccaga	gaacactgga	420
gggggatgct	ctagttgggt	ctgtgtccat	tttcctctgt	gccaaagaca	gacagacaga	480
ggctgagaga	ggctgttccct	gaatcaaagc	aatagccagc	tttcgacaca	tacctggctg	540
tctgaggagg	aaggccctct	ggaaaactggg	agctaagggc	gaggcccttc	ccttcagagg	600
ctcctggggg	attaggggtg	gggtgttgcc	aagccaaggg	gtagggagcc	gagaaattgg	660
tctgtcggct	cctgggtgca	ctttggggaa	ggagaggaa	tttggggctc	caggtagctc	720
cctgtttggt	gactgctctg	tcccctgccc	ctactgcaga	gatagcactg	ccgagttccc	780
ttcaggccctg	gcagacgggc	agtgaggagg	ggcctcagtt	agctctcaag	ggtgccttcc	840
cctcctccca	acccagacat	accctctgcc	aaactgggaa	ccagcagtg	tagtaactac	900
ctcacagagc	cccagagggc	ctgcttgagc	cttcttgctc	cacaggagaa	gctgggtgct	960
ctaggcaacc	ccttctctcc	acctctctac	aggggtgggg	gttctccttt	ctttccctctg	1020
aagtgtttat	ggggagatcc	tagtggcttt	gccattcaaa	ccactcgact	gtttgcctgt	1080
ttcttgaaaa	ccagtagaag	ggaaacagca	cagcctgtca	cagtaattgc	aggaagattg	1140
aagaaaaatc	ctcatcaatg	ccaggggaca	taaaagccat	ttcccttcca	aatactcgac	1200
aatttagatg	cagaacattt	ctctgtattc	agacttagag	taacaccagc	tgaaaactgc	1260
agtttcttct	ctttggatac	ataaggcttc	tctatcgggg	tacgggacag	ggaggaggcc	1320
tcatgtctga	agggggattt	aggggagaga	gccgcagccc	tgaccctcgg	tcctgtgcac	1380
cgctttgggg	cacagtctga	tggcgctttt	gctggcgctt	tagtatgggt	gactccggat	1440
ggacaaaaga	aaaaaaattt	tttttcttga	atgaaaatagc	aggaagctcc	tcgggagcat	1500
gtgttttgat	taaccgtagt	gatggatgct	acgagtataa	atggattaac	tacctcaatc	1560
cttacagtaa	gattggaact	aagggcaggg	actcatgcat	aagggtatga	atcccagcca	1620
ggacaagtga	gttgaggctt	gtgccacaaa	aggtttgtcc	ttggggaaca	ggcaggcctg	1680
ccaggatccc	ccccatatcg	attgggctgg	gagggtgggc	cgtgaggctc	ccactttctg	1740
ctttccttgc	ccatgtgtca	cccccttggc	ctccagcttg	tcctctcttc	actttctata	1800
gctttgttgg	accagatggt	gaggaaagga	atggcctctt	cccttctaga	gggggctggc	1860
tggagtgaga	cctggggctt	ggcctggaac	ccaccacaca	gccccaaagt	caggaagcct	1920
ggggaaacca	gagctgagac	ctcttcaaca	gggtttcttt	gagatcctac	acctccattg	1980
ggcccttttt	cagtcttcaa	tggggggcca	gttggtctta	gaaggagaag	aggtgaagca	2040
ggatcctttg	ccctggggga	gtctgagggc	gcggtccttg	gactcattca	ggcgtctttt	2100
gtagttgggg	gagttccact	gggcgatccc	agccccctcc	cacccaccct	ctaattggacc	2160
tcctcataga	agccccatct	cacttttggt	ttatctacct	cttagcaaaa	caatagataa	2220
attaggtagt	ggcagctcca	cttgcttagg	ttaggggggg	aaaaagattt	ctttttccaa	2280
aggaaaaaaa	tattaccttg	agaatacttt	ccaaaaaata	aaatttaaaa	aaaaaa	2336

<210> 56  
<211> 2200  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 1429651CB1

<400> 56

gagaacttta	cgctggatc	tcatctaact	gacacagaaa	ccctgtaagg	atccagaggt	60
ctcggttcagg	accatggaga	gcggcaccag	cagccctcag	cctccacagt	tagatccctt	120
ggatgcgttt	ccccagaagg	gcttgaggcc	tggggacatc	gcgggtgtag	ttctgtactt	180
cctctttgtc	ctggctgttg	gactatggct	cacagtgaag	acaaaaagag	acacagtga	240
aggctacttc	ctggctggag	gggacatggt	gtgggtggca	gtgggtgcat	cctgtttg	300
cagcaatggt	ggaagtggac	atttcattgg	cctggcaggg	tcagggtgctg	ctacgggcat	360
ttctgtatca	gcttatgaac	ttaatggctt	gttttctgtg	ctgatgttgg	cctggatctt	420
cctaccatc	tacattgctg	gtcaggtcac	cacgatggca	gaatacctac	ggaagcgctt	480
cggtggcatc	agaatcccca	tcactctggc	tgtactctac	ctatttatct	acatcttcac	540
caagatctcg	gtagacatgt	atgcaggctg	catcttcac	cagcagtcct	tgcacctgga	600

tctgtacctg	gccatagttg	ggctactggc	catcactgct	gtatacacgg	ttgctgggtg	660
cctggctgct	gtgatctaca	cggatgccct	gcagacgctg	atcatgctta	taggagcgct	720
caccttgatg	ggctacagtt	tgcgcgctg	tggtgggatg	gaaggactga	aggagaagta	780
cttcttgccc	ctggctagca	accggagtga	gaacagcagc	tgcgggctgc	cccgggaaga	840
tgccttccat	attttccgag	atccgctgac	atctgatctc	ccgtggccgg	gggtccctatt	900
tggaatgtcc	atccccatccc	tctggtagctg	gtgcacggat	caggtgattg	tccagcggac	960
tctggctgccc	aagaacctgt	cccattgccaa	aggaggtgct	ctgatggctg	catacctgaa	1020
ggtgctgccc	ctcttcataa	tggtgttccc	tggtgatggct	agccgcatcc	tcttcccaga	1080
tcaagtggccc	tgtgcagatc	cagagatctg	ccagaagatc	tgacagcaacc	cctcaggctg	1140
ttcggacatc	gcgtatccca	aactcgtgct	ggaactcctg	cccacagggc	tccgtgggct	1200
gatgatggct	gtgatgggtg	cggctctcat	gtcctccctc	acctccatct	ttaacagtgc	1260
cagcaccatc	ttcaccatgg	acctctggaa	tcacctccgg	cctcgggcat	ctgagaagga	1320
gctcatgatt	gtgggcaggg	tggttggctg	gctgctggct	ctgggtctcca	tcctctggat	1380
cctgtgggtc	caggccagcc	agggcggcca	gctcttcate	tatatccagt	ccatcagctc	1440
ctacctgcag	ccgctgtgg	cgggtggtctt	catcatggga	tggttctgga	agaggaccaa	1500
tgaaaagggt	gccttctggg	gcctgatctc	gggctgctc	ctgggcttgg	ttaggctggt	1560
cctggacttt	atttacgtgc	agcctcgatg	cgaccagcca	gatgagcgcc	cggtcctggt	1620
gaagagcatt	cactacctct	acttctccat	gatcctgtcc	acggtcaccc	tcactactgt	1680
ctccaccgtg	agctggttca	cagagccacc	ctccaaggag	atggctcagcc	acctgacctg	1740
gtttactcgt	cacgaccccg	tggtccagaa	ggaacaagca	ccaccagcag	ctcccttgtc	1800
tcttaccctc	tctcagaacg	ggatgccaga	ggccagcagc	agcagcagcg	tccagttcga	1860
gatggttcaa	gaaaacacgt	ctaaaaccca	cagctgtgac	atgaccccaa	agcagttcaa	1920
agtgggtgaag	gccatcctgt	ggctctgtgg	aatacaggag	aagggcaagg	aagagctccc	1980
ggccagagca	gaagccatca	tagtttccct	ggaagaaaac	cccttgggtg	agaccctcct	2040
ggacgtcaac	ctcattttct	gcgtgagctg	cgccatcttt	atctggggct	atcttctgta	2100
gtgtggggtg	aaccaggggg	tccaaactct	gtttctcttc	agtgtcccat	ttttttaatg	2160
aaagaaaaaa	taataaagct	tttgtttacc	aaaaaaaaaa			2200

&lt;210&gt; 57

&lt;211&gt; 2823

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2069971CB1

&lt;400&gt; 57

gaaagacata	cacacttcat	gtaatgctac	ctgcaagtct	ccctagaaaa	gcagtttttg	60
taggtgaaaa	caatgaagcc	aggtaatat	gcaaggaggc	tgtaatttta	gcagacctac	120
caacaacact	gatgtaggaa	gctcattatt	ttaattttctg	gagcctttta	atcttttctt	180
tagaaaagtgt	ataaataatt	gcagtgtgct	tttgtctcca	aaactgggca	gtgagttaa	240
caacaacgac	aacaacagcc	gcagctcatc	ctggccgtca	tggaagttct	tgaaagaacg	300
tatcttgtga	atgataaagc	tgccaagatg	tatgttttca	cactagaaaag	aaggagctgc	360
aaatgaacac	ttcatagcaa	tgtggaactc	caacagaaac	cgggtgaataa	agatcagttg	420
cccagagaga	gaccagagga	gctggagtc	ggaggcatgt	accactgcca	cagtggctcc	480
aagcccacag	aaaagggggc	gaatgagtac	gcctatgcca	agtggaaact	ctgttctgct	540
tcagcaatat	gcttcatttt	catgattgca	gaggtcgtgg	gtgggcacat	tgctgggagt	600
cttgctgttg	tcacagatgc	tgcccacctc	ttaattgacc	tgaccagttt	cctgctcagt	660
ctcttctccc	tgtggttgct	atcgaagcct	ccctctaagc	ggctgacatt	tggtatggc	720
cgagcagaga	tccttgggtg	cctgctctcc	atcctgtgca	tctgggtggg	gactggcggt	780
ctagtgtacc	tggtcatgtg	gcgcctgtg	tatcctgatt	accagatcca	ggcgactgtg	840
atgatcatcg	tttccagctg	cgcagtgccg	gccaacattg	tactaactgt	ggttttgcac	900
cagagatgcc	ttggccacaa	tcacaaggaa	gtacaagcca	atgccagcgt	cagagctgtg	960
tttgtgcatg	cccttgagga	tctatttcag	agtatcagtg	tgctaattag	tgactttatt	1020
atctacttta	agccagagta	taaaatagcc	gacccaatct	gcacattcat	cttttccatc	1080
ctggtcttgg	ccagcaccat	cactatctta	aaggacttct	ccatcttact	catggaaggt	1140
gtggccaaaga	gcctgaatta	cagtgggtgtg	aaagagctta	tttttagcagt	cgacgggggtg	1200
ctgtctgtgc	acagcctgca	catctggtct	ctaacaatga	atcaagtaat	tctctcagct	1260
catgttgcta	cagcagccag	ccgggacagc	caagtgggtc	ggagagaaat	tgctaaagcc	1320
cttagcaaaa	gctttacgat	gcactcactc	accattcaga	tggaatctcc	agttgaccag	1380
gaccccgact	gccttttctg	tgaagacccc	tgtgactagc	tcagtcacac	cgtagtttct	1440
ccaaatttga	caggccacct	tcaaacatgc	ttctatgcag	ttctctcatc	atagaaaaata	1500
aggaacaaaa	ggaagaaatt	catgtcatgg	tgcaatgcac	atcttatctc	tttatttagt	1560
tccattcacc	atgaaggaag	aggcactgag	atccatcaat	caattggatt	atatactgat	1620
cagtagctgt	gttcaattgc	aggaatgtgt	atatagatta	ttcctgagtg	gagccgaagt	1680
aacagctgtt	tgtaactatc	ggcaatacca	aattcatctc	ccttccaata	atgcactctg	1740
agaacacata	ggtaaaattg	aactcaggaa	agtccttacta	gaaatcagtg	gaagggacaa	1800



```

atagtcacaa aattttacca aaacattaga aacaaaaaat aaggagagcc aagtcaggaa 1860
taaaagtgtgac tctgtatgct aacgccacat tagaacttgg ttctctcacc aagctgtaat 1920
gtgattttttt tttctactct gaattggaaa tatgtatgaa tatacagaga agtgcttaca 1980
actaatttttt atttacttgt cacatttttg caataaatcc ctcttatttc taaattctaa 2040
cttggtttatt tcaaaacttt atataatcac tgttcaaaag gaaatatttt cacctaccag 2100
agtgtcttaaa cactggcacc agccaaagaa tgtggttga gagaccaga agtcttcaag 2160
aacagccgac aaaaacattc gagttgaccc caccaagtgt ttgccacaga taatttagat 2220
atttacctgc aggaaggaat aaagcagatg caaccaattc attcagtcca cgagcatgat 2280
gtgagcactg ctttgtgcta gacattgggc ttagcattga aactataaag aggaatcaga 2340
cgcagcaagt gcttctgtgt tctggtagca actcaacact atctgtggag agtaaactga 2400
agatgtgagc gccaacattc tggaaatcct atgtcactgg gtttgggttg gaaacctgga 2460
cttctgcatt tttaaaagtt acccagagat gcttctaaag atgagccata gtctagaaga 2520
ttgtcaacca caggagttca ttgagtggga cagctagata catacattgg cagctacaat 2580
agtatcatga attgcaatga tgtagtgggg tataaaagga aagcagtgga tattgccgga 2640
tgggcatggc cagtgtgtt tcacgtcatt gaggtgacag ctctgtctga ctttgaatta 2700
catatggagg ctctccagga agacgaagaa gagaaggaca ttctaggcaa aaagaagact 2760
aggcacaagg cacacttatg tttgtctgtt agcttttagt tgaanaagca agatacaggg 2820
tcg

```

&lt;210&gt; 58

&lt;211&gt; 1491

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2329339CB1

&lt;400&gt; 58

```

cgcctccctc cagctgcgag tgcggcctcg gctggcgggc gcaccaggcc acagttgtaa 60
gggatcttgt ggctgtcagg atggcagagg agcaggaggt caccagctc tgcaagttgc 120
ctgcacagcc ctacacacca cactgcgtga acaacaccta ccgcagcgca cagcactccc 180
aggctctgct ccgagggctg ctggctctcc gggacagcgg aatcctcttc gatgttggtc 240
tgggtgggga gggcagacac atcgaggccc atcgcatcct gctggctgctg tcctgctgatt 300
acttcagagg aatgtttgct gggggattga aggagatgga acaggaagag gtccctgatcc 360
acgggtgtgtc ctacaatgct atgtgccaaa tctacattt catatacacc tccgagctgg 420
agctcagcct gagcaatgta caagagacac tgggtggctgc ctgccagctt cagatcccag 480
aaattatcca tttctgctgt gatttctctca tgtcctgggt ggacgaagag aacattctcg 540
atgtctaccg gctggcagag ctgtttgact tgagccgcct gactgagcaa ctggacacct 600
atatectcaa aaactttgtg gccttctctc ggactgacaa gtaccgccag ctccattgg 660
agaaggtcta ctccctcctc agcagcaatc gcctggaggt ctccctgcgag accgaggtat 720
atgagggggc ccttctctac cattatagcc tggagcaggt gcaggctgac cagatctcgc 780
tgcacgagcc cccaaagctc cttgagacag tgcgggttcc gctgatggaa gctgaggtcc 840
tgcagcggct gcatgacaag ctggacccca gccctttgag ggacacagtg gccagcggc 900
tcatgtacca ccggaacgag agcctacagc ccagcctgca gagcccgcaa acggagctgc 960
ggtcggactt ccagtgcgtt gtgggcttcg ggggcattca ctccacgccg tccactgtcc 1020
tcagcgacca gcccaagtat ctaaaacctc tactgggaga gtggaagcac ttcactgcct 1080
ccctggcccc ccgcatgtcc aaccagggca tgcgggtgct caacaacttc gtatacttga 1140
ttggaggggga caacaatgtc caaggatttc gagcagagtc ccgatgctgg aggtatgacc 1200
cacggcaciaa ccgctgggtc cagatccagt ccctgcagca ggagcacgcc gacctgtccg 1260
tgtgtgttgt aggcaggtac atctacgctg tggcggggccg tgactaccac aatgacctga 1320
atgctgtgga gcgctacgac cctgccacca actcctgggc atacgtggcc ccaactcaaga 1380
gggaggtgta tgcacacgca ggcgcgacgc tggaggggaa gatgtatatc acctgcccga 1440
gcaagcttat tccctttagt gagggttaat tttagcttgc actggccgtc g 1491

```

&lt;210&gt; 59

&lt;211&gt; 986

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2540219CB1

&lt;400&gt; 59

```

ggagcgccaa ctcggtttga agtccaggcc agtggctcct gcggacagcc agcataccag 60
gggccaagtgc actgcattac aaccattgtg aggaatgagg gcctgtgccc tgtggctggc 120
gggcggcatg gcaggagcaa tttcttgggg gacagcgact cctatggatg tcgtgaaaaa 180

```

tcgactccaa	gctgatgggg	tttattttaa	caaataataa	gggtgcctgg	actgtatctc	240
ccagagttac	cagaaggaa	gtcttaaagt	gtttttcaga	ggcatcactg	tgaacgcggg	300
gcggggcttc	cccatgagtg	cggccatgtt	ccttgggtac	gagctgtcgc	tgcaggctat	360
ccgcggggac	cacgcagtg	cgagcccata	agcgccagga	ggatgaacaca	ggatgactac	420
agtgttcccc	tgggcctcat	ctctgcatgt	gaagccctga	gagctgcaga	tgtttgccct	480
ttggacctcc	aagtggacat	caattagcaa	gcgtgggcta	ggatgggtga	gacactgacg	540
tggcccttct	gatgcctggg	atgcctcatg	agtcactgat	tcaaagccctc	caaggttctg	600
atccccaatg	cccactctgc	taggctggca	tcaaagagct	ttccaagaaa	tgtttggtcc	660
agctgagaag	tcctgaccat	gagcaccagg	gagccagaaa	ccaccagag	aaacgttgct	720
tcactcctct	gtctgaggat	ggggaggggc	cagtgcctc	tgggctcagc	cactccctcc	780
agtctcaagt	aacacgtccc	cgtgcctcca	gtctcctctc	agcaccgacc	aggtttttcc	840
ccgctcctgc	acccgtggat	cctgaggaca	gcggtagcgc	cttccctacc	gcacgctgag	900
tccagtgcgt	gtcctctcact	gtgcacttat	tagtgtctgt	tgagtgatta	aatcacatcc	960
tcagggtctgc	agcaataaaa	tgaaag				986

&lt;210&gt; 60

&lt;211&gt; 4023

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2722462CB1

&lt;400&gt; 60

gtccggggcg	gcgcgtatgg	cggcactatg	ccatccacga	ctcccaggcc	cccagtctca	60
gctctggggg	tgagagttcc	ccctccagcc	ccgcacacaa	ctgggagatg	aattaccaag	120
aggcagcaat	ctacctccag	gaaggcgaga	acaacgacaa	gttcttcacc	caccccaagg	180
atgccaaagg	gctggcggcc	tacctctttg	cacacaatca	cctctttctac	ctgatggagc	240
tggccacggc	cctgctgctg	ctgctgctct	cctgtgcgga	ggcccccgcc	gtccccgcac	300
tccggcttgg	catctatgtc	cacgccaccc	tggagctgtt	tgccctgatg	gtggtagtgt	360
ttgaactctg	catgaagtta	cgctggctgg	gcctccacac	cttcatccgg	cacaagcgga	420
ccatggtcaa	gacctcggtg	ctggtggtgc	agtttgcga	ggccatcggt	gtgttggtac	480
ggcagatgtc	ccatgtgcgg	gtgacccgag	cactgcgctg	cattttcctg	gtggactgtc	540
gggtattgcgg	tggcgctccg	cgcaacctgc	ggcagatctt	ccagtccctg	ccgcccctca	600
tggacatcct	cctgctgctg	ctgttcttca	tgatcatctt	tgccatccctc	ggtttctact	660
tgttctcccc	taacccttca	gacctctact	tcagcacctc	ggagaacagc	atcgctagtc	720
tgtttgcctc	tctgaccaca	gccaatttcc	cagatgtgat	gatgccctcc	tactcccgga	780
acccctgggtc	ctgctgtctc	ttcatcgtgt	acctctccat	cgagctgtat	ttcatcatga	840
acctgcttct	ggctgtggtg	ttcgacacct	tcaatgacat	tgagaaacgc	aagttcaagt	900
ctttgctact	gcacaagcga	accgctatcc	agcatgccta	ccgcctgctc	atcagccaga	960
ggaggcctgc	cggcatctcc	tacaggcagt	ttgaaggcct	catgcgcttc	tacaagcccc	1020
ggatgagtgc	cagggagcgc	tatcttacct	tcaaggccct	gaatcagaac	aacacacccc	1080
tgctcagcct	aaaggacttt	tacgatctct	acgaagtgtc	tgctttgaag	tggaaggcca	1140
agaaaaacag	agagcactgg	tttcatgagc	ttcccaggac	ggcgctcctc	atcttcaaag	1200
gtattaatat	ccttgtgaag	tcgaaggcct	tccagtattt	catgtacttg	gtgggtggcag	1260
tcaacggggg	ctggatcctc	gtggagacat	ttatgtctgaa	aggtgggaac	ttcttctcca	1320
agcacgtgcc	ctggagtac	ctcgtctttc	taactatcta	tggggtggag	ctgttctctga	1380
aggttgccgg	cctgggcccct	gtggagtact	tgtcttccgg	atggaaacttg	tttgacttct	1440
ccgtgacagt	gttcgccttc	ctgggactgc	tggcgctggc	cctcaacatg	gagcccttct	1500
atctcatcgt	ggtcctgcgc	ccctccagc	tgtgaggtt	gtttaagttg	aaggagcgct	1560
accgcaacgt	gctggacacc	atgttcgagc	tgctgccccg	gatggccagc	ctgggcccga	1620
ccctgctcat	cttttactac	tccttcgcca	tcgtgggcat	ggagttcttc	tgcgggatcg	1680
tcttccccaa	ctgctgcaac	acgagtacag	tggcagatgc	ctaccgctgg	cgcaaccaca	1740
ccgtggggcaa	caggaccgtg	gtggaggaag	gctactatta	tctcaataat	tttgacaaca	1800
tcctcaacag	ctttgtgacc	ctgtttgagc	tcacagttgt	caacaactgg	tacatcatca	1860
tgggaaggcgt	cacctctcag	acctcccact	ggagccgcct	ctacttcatg	acctttttaca	1920
ttgtgaccat	ggtggtgatg	acgatcattg	tcgcctttat	cctcgaggcc	ttcgtcttcc	1980
gaatgaacta	cagccgcaag	aacaggactc	cggaggttga	tgggtggcatc	acctttgaga	2040
aggaaatctc	caaagaagag	ctgggtggcg	ctaccgggag	ctaccgggag	gcacgggggg	2100
cctcctcgga	tgtcaccagg	ctgctggaga	ccctctccca	gatggagaga	taccagcaac	2160
attccatggt	gtttctggga	cggcgatcaa	ggaccaagag	cgacctgagc	ctgaagatgt	2220
accaggagga	gatccaggag	tggatgagg	agcatgccag	ggagcaagag	cagcagcgac	2280
aactcagcag	cagtgcagcc	cccgccgccc	agcagccccc	aggcagccgc	cagcgctccc	2340
agaccgttac	ctagcccagc	gcccgaagag	cgtctcttct	atgcaataac	acaatagtat	2400
tactctactg	cgatgtacgg	aactgcgggtg	tgtgtacaca	tactcacgta	tatgcacata	2460
tttatataca	ggaagaaaaa	agacagacaa	gatggggcct	ggtttataac	caccttgccc	2520
tgtcttctct	aactccagaa	gccagtttgg	tgaggggtgg	gggtgcggcc	accaggtctg	2580

agctcttctt	actgtggaag	gctccagaag	gcccttcaca	aggagacccc	tcacctggat	2640
ccagtcgact	gcggggcttg	cccctcatgt	gggctggcct	ccatcggcca	cgctccaaagc	2700
tgtcactgct	actgcttcag	gctcacatcc	ccccgacctg	atggcggtgcc	cgccccctct	2760
ccctgcggcc	catgccacag	gtttctgtgt	tttgcttttag	ggacagaacc	acttaggaaa	2820
gaaagaactc	ccggtctcca	gggtgggtatt	tcagtgtctg	tgataatgtc	acgcaacacc	2880
tcttcgggga	ccagtgcaca	ggatctaattg	gaagcggaat	tggggcaact	gggcccgtgt	2940
ggccagagct	cagttagcca	gtgccggggc	gccacagatt	acactgacca	atctcctccc	3000
ttggctctgc	aagcctccca	cccagccttc	tctggcttaa	cccttggttg	cgaaaaactct	3060
tcacacagtg	cctccttggg	gaccagaac	ccggagggaag	gggcatgagg	caggaagtgg	3120
ggccgatgtc	tgcaaccag	accacttcgt	ggaatgggct	cttgaccaa	tccctttttt	3180
ttgagattta	cccgttcaag	caaaacaacg	ttttgggttaa	ctaaggattg	tgctaaagcc	3240
gataccaggt	ccttcacacg	tgtgcactag	gaacaggagc	gaacagcaca	gagagacgct	3300
ccctgtggga	cgcagcagcc	ggcccagttc	ggcccagttc	ccagccaccc	tccctggctc	3360
tgctcacacc	agagatttcc	atagcaggag	cggttgggtg	agaagtaggt	tcagatgaac	3420
ctcagttaac	gtcccccacc	ctcctcccac	catggtaccc	tgtaggagcc	ctgtatgaca	3480
tctgagctgt	gtggaggtag	gagggttgcc	agctgcagtg	accctgccac	agaggcaggg	3540
tcagtgcaga	ggtcgctttg	gttcgcttc	cctggggccac	agaacggaac	acagcatagg	3600
ttctgcagca	ggagccgcag	tggcaggatg	gaggggtcga	agggcaagga	gtgcaactgt	3660
gggcattcct	ggccagcccc	ggccctctgg	tgctgtcttc	ctgtgacttc	agaaggcagg	3720
tggaacagag	ctccctctgg	cttgtcttc	ttcccagcca	cagaacgggc	aggggtggc	3780
ccgaccccag	gggagcagta	cctgggtccc	caccccctcc	tcccaaccac	ctccaaggcc	3840
aagctgggtc	ccatagccag	cacggcatgg	ttctcccctt	cccccccttc	caggtcaggg	3900
gagttggaca	agtagcaggt	gtttgttttt	aaagcacagc	cctttgggaa	agcaacacat	3960
tattgagact	cactgtgatt	ccccggggag	tcagactggc	tttgtctctt	tctctctgga	4020
ggg						4023

&lt;210&gt; 61

&lt;211&gt; 2345

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2739264CB1

&lt;400&gt; 61

gaaaggaggc	ggaactcggg	gatctgactg	gcgggtttccc	ggccgggactg	agaaggggag	60
cgcgctgcgc	gtcgcaggag	taacctactt	gggtctcctgc	tttcgcgaca	tgcccttcaa	120
ttttggggct	ccctcgggca	cctcgggtac	cgctgcagcc	accgcgggccc	ccgcggggtgg	180
gtttggaggga	tttggggacaa	catctacaac	tgcaagttct	gcattcagct	tttctgcccc	240
aactaacaca	ggcactactg	gactcttttg	tggtactcag	aacaaagggtt	ttggattttg	300
tactggtttt	ggcacaacaa	cgggaactag	tactggttta	ggtactgggtt	tgggaaactgg	360
actgggattt	ggaggattta	atcacacagca	gcagcagcaa	actacattag	gtggctctct	420
cagtcagcct	acacaagctc	ctaccagctc	caaccagctg	ataaatactg	cgagtgtctt	480
ttctgtctcca	acgctgtttg	gagatgagag	agatgctatt	ttggcaaaat	ggaatcaact	540
gcaggccctt	tggggaacag	gaaaagggta	tttcaacaat	aatattccgc	cagtggaatt	600
cacacaagaa	aatccctttt	gccgatttaa	gccagtaggt	tatagttgca	tgcccagtaa	660
taaaagatgaa	gatgggctag	tggttttagt	tttcaacaaa	aaagaaacag	agatttcgaag	720
ccaacaacaa	cagttggtag	aatcattgca	taaagttttg	ggaggaaacc	agacccttac	780
tgtaaatgta	gagggcacta	aaacattgcc	agatgatcag	acagaagttg	ttattttatgt	840
tgttgagcgt	tcgccaatg	gtacttcaa	aagagttcca	gctacaacgc	tatatgccca	900
ttttgaacaa	gccaatataa	aaacacaatt	gcagcaactt	ggtgtaaccc	tttctatgac	960
tagaacagaa	ctttctcctg	cacagatcaa	acagctttta	cagaatcctc	ctgctgggtg	1020
tgatcctatt	atctgggaac	aggccaaggt	agataaccct	gattctgaaa	agttaattcc	1080
tgtaccaatg	tggggtttta	aggaaactct	ccgaagactg	aaggttcaag	atcagatgac	1140
taagcagcat	caaaccagat	tagatatcat	atctgaagat	attagtgagc	tacaaaagaa	1200
tcaaactaca	tctgtagcca	aaattgcaca	atacaagagg	aaactcatgg	atctttccca	1260
tagaacttta	caggtcctaa	tcaaacagga	aattcaaagg	aagagtgggt	atgccattca	1320
ggctgatgaa	gagcagttgc	gagttcagct	ggatagcatt	caggggtgaac	taaatgcacc	1380
tactcagttc	aagggccgac	taaatgaatt	gatgtctcaa	atcaggatgc	agaatcattt	1440
tggagcagtc	agatctgaag	aaaggtatta	catagatgca	gatctgttac	gagaaatcaa	1500
gcagcatttg	aggaaggcct	tagccatttg	attagcatca	accatccaca	ttaaagacga	1560
tctagaagat	ataaagctgg	tcgaacatgg	attgaatgaa	accatccaca	tcagaggtgg	1620
tgtcttttagt	tgacagttca	caaacttgtg	taaaggtttg	tgaaatgcat	cttcttactg	1680
catcagacct	tccttaagaa	tgaaccgcac	cacatggagg	gaaaaagaaa	acaattcttt	1740
cttggatttg	ttttttgaga	agtttactga	caaattactg	ttcatcaa	ctgaaatagt	1800
cacctcacag	ctcttcaaa	aaaacctttg	aaagatttat	atctaaaagc	tgtatttact	1860
ttaaaagaag	tgacataatta	ccaaaattgt	atgtactatt	gtacattttt	acaacagcat	1920

```

tttcttaaac ataatctgtg tttaatgatt attgtccatt gagcctgtac tctgctttcc 1980
ataccaagta aatatgaaat aatctacttt gcacataaca gaacaaacta taattacttg 2040
gctgttgagg atttgtactt gagtataaat gtacaccagt ttttgtatgt gtgaactcat 2100
ctgtgggagg agtaaagaaa atccaaaagc atttaagtgt ttgtttttgt tctataaaga 2160
tatgaaaatg tatttttata ttattttact tatttggaa ttacagagca cacctaagca 2220
attaggatat aacaaaacta ctttaaccatt tttgcaacca ttttgttttt taagcctttt 2280
tatttctaaa aagatgaaaa cttataaata aattcttaat ttgtaattac ttttaaaaaa 2340
aaaaa 2345

```

&lt;210&gt; 62

&lt;211&gt; 2085

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2758310CB1

&lt;400&gt; 62

```

cggagatgtg gcgaccgttt ctggcatcat tctgagactc ggcagttgct tctcactgct 60
gcggccgggc ctgtctgtgg gagctgcac ctcctcatct gcaggcgtg gaaaaccaga 120
cacgatcgga catgcatgtg gttctgcggc caaagcacgc ctttgggttg tgaacttcat 180
gatacctgtg tgcagctctg tcatctccac tctgctctgc tgcacagaag gcagaagccc 240
tgcccggtcc ctgctgtgtt cttcaggaga aacgtcaggg gccttcctcc aaggttcttc 300
agccccacac ccctgtggag gaaggtgtct tccaccgcgg tagtgggggc gccctgtctc 360
ctcggagccc gctatgtcat ggcagaggca cgggagaaga ggaggatgcg gctcgtgggtg 420
gatggcatgg ggcgctttgg caggtctctg aaggtcggcc tgcagatctc cctggactac 480
tggtgggtgca ccaatgttgt ccttcgaggg tggaagagcc caggctactt ggaggtgatg 540
tctgctgtgc accagcgggc ggctgatgcc ctggtggcag gggccatcag caacgggggc 600
ctctacgtga agctgggcca ggggctgtgc tccttcaacc acctgcttcc ccccgagtat 660
accgggacct tgcgctgtct agaggacagg gccctcaagc ggggcttcca ggaggtggat 720
gagttgttcc ttgaggactt ccaggccctc ccccacgagc tcttcaggga gtttgactac 780
cagccaattg ctgccgccag cctggcacag gtgcacagag ccaagctgca cgatggcacc 840
agcgtggctg tgaaggtgca gtacatcgac ctgcgggacc gctttgatgg ggacatccac 900
accctggagc tcctgtgcg gctcgttgag gtcatgcacc ccagctttgg cttcagctgg 960
gtcctccagg acctgaagg gacctggcc caggagctgg acttcgagaa tgagggccgc 1020
aacgcagagc gctgtgctgc ggagctggcg cacttccctt acgtcgtggg gccccgcgtg 1080
cactgggaca agtccagcaa gcgctgtctc actgcccact tctgcgccgg ctgcaaggtc 1140
aacgatgtgg aggccatcag gagccagggg ctggcagtg atgacatagc agaaaagctc 1200
atcaaggcct ttgctgagca agggccggac gggaaagcgg agctggtgct gctggaccac 1320
gggctctacc agttcctgga ggagaaggac cgcgcagccc tctgccagct gtggcggggc 1380
atcatcctgc gggacgacgc cgcatgagg gcgcacgcag ccgactggg ggtgcaagac 1440
tacctcctgt tcgcccagat gctcatgag cgccccgtgc gcctggggca gctgtggggc 1500
tcgcacctac tgagcccgga agaggcggcc tacatggtgg acatggcccg cgagcgttcc 1560
gaggccgtca tggcgtgtgt cagggagctg ccgcccga tgctgtggt gctgcgcaac 1620
atcaaacacc tgccgctat caacgtggcc ctgcgcccc ccgtggaccg ctacttctt 1680
atggctaaaa gggctgtccg gggctggagc cgcctggcgg gcgccacgta tcgggggtgtc 1740
tacggcacca gcctcctgcg ccacgccaag gtctgtctgg agatgtcaa gtttgaagtg 1800
gcgctcagggc tggagacctt ggccatgcgg ctgaccgccc tcctggctcg tgctctgggtc 1860
cacctgagcc tcgtgcccc agcggaggag ctctaccagt acctggagac ctagggtgca 1920
gccgcccagg gccggcgggg cccttttcac cttgggctga cggaggtggc ggggctagag 1980
gtgtagacac cccgagcccc gtgggcactc gcactggggg gctgtgacag cagctggggc 2040
aggaggccgt gtaatgacca cacactcctc tcaagcaaaa aaaaa 2085

```

&lt;210&gt; 63

&lt;211&gt; 3014

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 2762348CB1

&lt;400&gt; 63

```

ggaggcaaaag gccaggctga ggatcagggg gggccgggtg gcagcgggga ggcgctgcat 60
gctggaggct gtgctgagtg cccggtgcag gtgagccggg cctgcggagt tgtgccgagt 120
gcctgctgca gaccgaggct gggccaagat ggcgtctgtg tttcgaagcg aggagatgtg 180

```

```

ttgtgcacaa ctgtttctcc aggtggaagc tgcataattgc tgtgtggctg agctcggaga 240
gctcggattg gttcagttca aagatttaaa tatgaatgtg aacagctttc aaaggaaatt 300
tgtgaatgaa gtcagaaggt gtgaatcact ggagagaatc ctccgttttc tggaaagaca 360
gatgcaaaat gagattgtag ttcagttgct cgagaaaagc ccactgaccc cgctcccacg 420
ggaaatgatt accctggaga ctgttctaga aaaactggaa ggagagttac aggaagccaa 480
ccagaaccag caggccttga aacaaagctt cctagaactg acagaactga aatacctcct 540
gaagaaaacc caagacttct ttgagacgga aaccaattta gctgatgatt tctttactga 600
ggacacttct ggcctcctgg agttgaaagc agtgcttgca tatatgaccg gaaagtgtgg 660
gttcatagcc ggggtgtgat caacagggaa gaggatggct tccttttgagc ggttactgtg 720
gcgagtctgc cgaggaaacg tgtacttga catattcact atggagcggc ctctggagga 780
tcctgtgacg aaagaagaaa ttcagaagca gttcagtgag atattttacc aaggagagca 840
gctcaggcag aaaatcaaga agatctgtga tgggtttcga gccactgtct acccttggcc 900
agagcctgcg gtggagcgca gagagatgtt ggagagcgtc aatgtgaggc tggaaagattt 960
aatcaccgtc ataacacaaa cagagtctca ccgccagcgc ctgctgcagg aagccgtctg 1020
caactggcac tcctggctca tcaaggtgca gaagatgaaa gctgtctacc acatcctgaa 1080
catgtgcaac atcgacgtca cccagcagtg tgtcatcgcc gagatctggg tcccgggtgg 1140
agatgccaca cgtatcaaga gggcactgga gcaaggcatg gaactaagtg gctcctccat 1200
ggccccatc atgaccacag tgcaatctaa aacagccctt cccacattta acaggaccaa 1260
taaattcaca gctggcttcc agaataattgt tgatgcctat tttcccttcc ctgttcgctg 1320
gataaaccga gccccttaca ccacatcac tttcccttcc ggctgcactt tggatgattc tgaatgagag 1440
agactgtggg catggaaccg tgatgctcct gatttggaa accttcttcc acgggcgcta 1500
acgcttgctc tcccagaaga tcttctccat ctacacgggt ttgatctaca atgactgctt 1560
tctgatccta ctatgggca tcttctccat ttgggtcttc tggagtgtc caaccatgt tcagaaacgg 1620
ctccaagtcc ttgaacatct tgggtcttcc tctatatctg cagctggacc cagccatacc 1680
cacatggaat actcatgtaa tggaggaaag cagatttgat ccgatttggg acttggcttc 1740
aggagtgtat tttggaaatc cataccggtt aatgaagatg tcgggtgatcc tgggaattgt 1800
aaacaaactc acatttctga actcgtataa tttcaatcac atatacttca gaagaactct 1860
ccagatgggt tttgggtgca tcctcagcct gatttttatc ctgtgtctgt ttggatacct 1920
caacatcatt ctgcaattta tccttgagat ctttgacgtc cacgtatctc agcacgccc 1980
ggttttcatg atcattttca aatgggtgctg tctgtttaac tacagtgact cttccaacgc 2040
cagcatcctc atccacttca aagaagtcca aagtttcttt gtggttatgg ctttgatttc 2100
acccctctac aaacatcagc ttaagccgtt tattcttaga gccagtcac ggaaatccca 2160
tgtgccgtgg atgcttctga aagaagatgc cactgagaac attgaagggt atagctccag 2220
gctgcaggca tccaggatcc agaggacttc tgcagatacc cacggggctc tggacgacca 2280
ccttcttagc cgttctggcc gagacgtctt tgtccaccaa gccatccaca ccacgagta 2340
tggagaagag ttcaactttg acacagcctc ctctggggcc tcagcctggc 2400
ctgctggaag ctgtctgaag tgctctggac tatgggtgat aacagcggcc ttcagacggc 2460
tcattgcaaa aggcgtgcg ggggttttat tatttttgcc gtatttgctg tcttgacagt 2520
agccatcctt ctgatcatgg agggcctctc tgctttcctg cagccctgc gactgcaact 2580
ggttgagttc cagaacaagt tctatgtcgg ggtatggtac aagttttctc cattctcctt 2640
taaacacatc ctggatggca cagccgagga gtaggctgag ggctgcacct cccacggtgg 2700
tcaccatgcc aatgaaggaa gttcagtcct gtctttgata tcagccctg caaggcgctc 2760
aatgggaagg ttgttcttgg ctacactgaa gcatgaaact gtgtattatt tggacgtcag 2820
cctgtggatt tgatacgact taaccacgtc agaggaaagga ctttggcaag tgatattgtc 2880
ttcatgtggg gtattaattc tcaaataata aagtaattga caaatgaggg gagaatgcta 2940
aacagatgtc ttcttgcaat atttttaata ttgtatttga gaaaaataac atctgagta 3000
ttcaaaaaaa aaaa

```

3014

&lt;210&gt; 64

&lt;211&gt; 1726

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3715961CB1

&lt;400&gt; 64

```

tgaaaatgcc tgctgcgtac caaggtatgt actagggcat ctggggtaag taaaaacaaa 60
cacatagagc ctgcctggag aagctcatgg tctgatggaa agataagcaa gaagagttaa 120
tttctaatac atatgataaa aaggtcagag agcagtttct gaaaaacatg tttttgagtt 180
gagtcctgaa agacaaggag atgttagtaa agcagagaag ggagaattca ttctagaag 240
atcagacagt gtgtgggaag ggcagagtct gaaaagagca tgccccattt ggagaagcat 300
caagaagccc acgtgttaga agcaccggcc ccatgagaca aagacacagc tagagagatt 360
gactaggcca tgtcggaaatg tcctcttatt ttatacatat ataagcatat agatacatat 420
agccaaagtt acctttttaa tgatcttttt taccagtggt attctggagg tcgaatgggtc 480
acatatgaac atctccgaga ggttgtgttt ggcaaaagtg aagatgagca ttatccctt 540

```

```

tggaatcag tcattggagg gatgatggct ggtgttattg gccagttttt agccaatcca 600
actgacctag tgaagggttca gatgcaaatg gaaggaaaaa ggaaactgga aggaatacca 660
ttgcgatttc gtggtgtaca tcatgcattt gcaaaaatct tagctgaagg aggaatacga 720
gggctttggg caggctgggt acccaatata caaagagcag cactggtgaa tatgggagat 780
ttaaccactt atgatacagt gaaacactac ttggtattga atacaccact tgaggacaat 840
atcatgactc acggtttatc aagtttatgt tctggactgg tagcttctat tctgggaaca 900
ccagccgatg tcatcaaaag cagaataatg aatcaaccac gagataaaca aggaagggga 960
cttttgataa aatcatcgac tgactgcttg attcaggctg ttcaagggtga aggattcatg 1020
agtctatata aaggcttttt accatcttgg ctgagaatga ccccttggtc aatggtgttc 1080
tggtctactt atgaaaaaat cagagagatg agtggagtca gtccatttta aaccctataa 1140
gatgcaaccc ttaaagatac agtgttcagt attattgaaa tatgggcacg tgcaacacat 1200
accccttatt atttctacct ctttaggaag acacctattc cacagagact gatttatagg 1260
gggcagcact ttattttttt ctggaacccc aagttctctt tgactcctct ttttgtccaa 1320
aagtgatctg gtcggatctc acaaggccat ccaatgagac cccgcacagc attttctaaa 1380
gaagaatcga agcctgacca ctttcacctt gggcaagaag gtttggcctt tgagttgcta 1440
ttctatgctg aagagcctgc tttagaggag agtaccagga gggagccagc atttcagatc 1500
tgaagtagag gataggaaat tggaagaaca catacatagt gcttaagaaa tacatttaac 1560
ctgttatgtc agtatattatc aatgaagttt gataattcac ttttctgtca ttgttaaacg 1620
gtacatactg taaattaaag ggaggtgaat ggaattaat gaataaacat tttgagtttc 1680
cctagtgttg aagggaaggtg tactttttct tgtcagaaag ataaaaa 1726

```

<210> 65  
 <211> 899  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5108194CB1

```

<400> 65
gcgccggcga cgcgtccggg ccggtgaggg ggcgcggggg gcgcccgggg ggcccaagcg 60
tcagcgcgcc cgcctgtcgg gctgaactga ggaccgagtc tcctgccatt ccgagcagcg 120
ctgggtatgg taatggtgtg aaggaaaggg ccgtgcgatt gcatgaggat gctgaggctg 180
tcctgtcctc gtccgtctca tcaaagcgtg accacaggca agtgctcagc tccctgtctg 240
ctggggccct ggctggtgcc cttgccaaaa ctgccgtagc tccctgggac cgaacaaaa 300
tcattctcca agtgtcttca aaaagatttt ctgccaaagg ggccttccgg gtcctctact 360
acacctacct caacgagggg tttctcagct tgtggcgcgg gaactcggcc accatggtgc 420
gcgtggtgcc ctacgccgcc atccagtcca gcgcacacga ggagtacaag cgcattcctg 480
gcagctacta tggcttccgt ggagaagccc tgcccccttg gcctcgccctc ttcgccggcg 540
cactggctgg aacgacagcc gcttcactga cctacccctt ggacctggtc agagcgcgga 600
tggccgtaac cccgaaggaa atgtacagca acatctttca tgtcttcacg cgcattctga 660
gagaagaggg gctgaagact ctctaccatg gatttatgcc caccgtgctg ggggtcattc 720
cctacgctgg cctgagcttc ttcacctatg agacgttcaa gagcttgacg agagagtaca 780
gcggccgcaa gcttattccc tttagtggag gttaatttta gcttggcact ggccgtcggt 840
ttacaacgctc gtgactggga aaaccctggc gttacccaac ttaatcgctt tgcagcaca 899

```

<210> 66  
 <211> 643  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5503122CB1

```

<400> 66
ctttaagctg tagctgtggt ttctgcagca attttgtttt tgccttgaaa gaggtgctct 60
ggattatcac acctccatgt atgacaattt gtacctgcat ggaattgaag actcggagcg 120
tggttcagcg gattccctaca caagcaggcc gtctgactcc gatgtctctt tggaagagga 180
ccgggaagca attcgacagg agagagaaca gcaagcagct atccagcttg agagagcaaa 240
gtccaaacct gtagcatttg ccgtgaagac aaatgtgagc tactgcggcg ccctggacga 300
ggatgtgcct gttccaagca cagctatctc ctttgatgct aaagactttc tacatattaa 360
agagaaatat aacaatgatt ggtggatagg aaggctggtg aaagagggct gtgaaattgg 420
cttcattcca agtccactca gattggagaa catacggatc cagcaagaac aaaaaagagg 480
acgtttttcac ggagggaaat caagtggaaa ttcttcttca agtcttggag aaatggtatc 540
tgggacattc cgagcaactc ccacatcaac aggtgagggt tgtagttaaa ctctttttca 600
tacactgtat tcctttttaaa aatatttgaa cacacatgca agc 643

```

<210> 67  
<211> 2574  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 5517972CB1

<400> 67  
gcgcctccca cgccggccgc cgccgacgtg atcgctcggg cgccgggggc gtgggtgggg 60  
gaagggggtg tgccgcgcga cggctctgct gctgtgccc ctcaaaagggt tccggggcgcg 120  
caggacggaa gacgcagtgc tcgccactcc cactgagatt gagagacgcg gcaaggaggc 180  
agcctgtgga ggaactgggt aggatattagg aacgcaccgt gcacatgctt ggtgggtcttg 240  
ttaagtggaa actgctgctt tagagtttgt ttggaaggtc cgggtgactc atcccaacat 300  
ttacatcctt aattgttaaa gcgctgcctc cgagcgcacg catcctgaga tcctgagcct 360  
ttggttaaga ccgagctcta ttaagctgaa aagataaaaa ctctccagat gtcttccagt 420  
aatgtcgaag tttttatccc agtgtcacaa ggaaacacca atggcttccc cgcgacagct 480  
tccaatgacc tgaaggcatt tactgaagga gctgtgttaa gttttcataa catctgctat 540  
cgagtataaac tgaagagtgg ctttctacct tgtcgaaaa cagttgagaa agaaatatta 600  
tcgaatatca atgggatcat gaaacctggg tctcaacgcca tcctgggacc cacagtgga 660  
ggcaaatctt cgttattaga tgtcttagct gcaaggaaag atccaagtgg attatctgga 720  
gatgttctga taaatggagc accgcgacct gccaatttca aatgtaattc aggttacgtg 780  
gtacaagatg atgttgtgat gggcactctg acggtgagag aaaacttaca gttctcagca 840  
gctcttcggc ttgcaacaac tatgacgaat catgaaaaaa acgaacggat taacagggtc 900  
attcaagagt taggtctgga taaagtggca gactccaagg ttggaactca gtttatccgt 960  
ggtgtgtctg gaggagaaag aaaaaggact agtataggaa tggagcttat cactgatcct 1020  
tccatcttgt tcttggatga gcctacaact ggtcttagct caagcacagc aaatgctgtc 1080  
cttttgcctc tgaaaaggat gtctaagcag ggacgaacaa tcatcttctc cattcatcag 1140  
cctcgatatt ccatcttcaa gttgtttgat agcctcacct tattggcctc aggaagactt 1200  
atgttccacg ggcctgctca ggaggccttg ggatactttg aatcagctgg ttatcactgt 1260  
gaggcctata ataaccctgc agacttcttc ttggacatca ttaatggaga ttccactgct 1320  
gtggcattaa acagagaaga agactttaaa gccacagaga tcatagagcc ttccaagcag 1380  
gataagccac tcatagaaaa attagcggag atttatgtca actcctcctt ctacaagag 1440  
acaaaagctg aattacatca actttccggg ggtgagaaga agaagaagat cacagtcttc 1500  
aaggagatca gctacaccac ctcttctgt catcaactca gatgggttcc caagcgttca 1560  
ttcaaaaact tgctgggtaa tccccaggcc tctatagctc agatcattgt cacagtcgta 1620  
ctgggactgg ttataggtgc catttacttt gggctaaaa atgattctac tggaatccag 1680  
aacagagctg gggttctctt ctctctgacg accaaccagt gtttcagcag tgtttcagcc 1740  
gtggaactct ttgtggtaga gaagaagctc ttcatacatg aatacatcag cggatactac 1800  
agagtgtcat cttatttcct tggaaaactg ttatctgatt tattaccatc gaggatgtta 1860  
ccaagtatta tatttacctg tatagtgtac ttcatgttag gattgaagcc aaaggcagat 1920  
gccttcttcg ttatgatgtt tacccttatg attggtgctt attcagccag ttccatggca 1980  
ctggccatag cagcaggtca gagtgtggtt tctgtagcaa cacttctcat gaccatctgt 2040  
tttgtgttta tgatgatttt ttcaggtctg ttggtcaatc tcacaacctat tgcattcttg 2100  
ctgtcatggc ttcagtactt cagcattcca cgatattgat ttacggcttt gcagcataat 2160  
gaatttttgg gacaaaactt ctgcccagga ctcaatgcaa caggaaacaa tccttgtaac 2220  
tatgcaacat gtactggcga agaataattg gtaaaagcagg gcatcgatct ctaccctgg 2280  
ggcttgtgga agaatacagt ggccttggct tgtatgattg ttattttcct cacaattgcc 2340  
tacctgaaat tgttatttct taaaaaatat tcttaaatat ccccttaatt cagtatgatt 2400  
tatcctcaca taaaaaagaa gcactttgat tgaagtattc aatcaagttt ttttgtgtgt 2460  
ttctgttccc ttgccatcac actgttgac agcagcaatt gttttaaaga gatacatatt 2520  
tagaaatcac aacaaactga attaaacatg aaagaaccca agaaaaaaa aaaa 2574

<210> 68  
<211> 1571  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 5593114CB1

<400> 68  
ggcgcggcga cagctagggg tcacggccac tggggcagag gagccgcgag aagatgtggg 60  
tttttggtta cgggtccctg atctggaagg ttgatttccc ctatcaggac aagctggctg 120  
gatacatcac caactacagc aggcgcttct ggcagggcag cagggaccac cgcggggtcc 180  
ccggcaagcc tggaaagagt gtgactcttg ttgaagatcc tgcgggatgt gtatgggggtg 240

```

ttgcttacag attgccagta ggaaaggaag aagaagtaaa agcatacctt gacttcagag 300
aaaaaggagg ctacagaacc acaacagtca ttttttatcc aaaagatccc acaacaaaac 360
cattcagtggt attgctatat attggaacat gtgataatcc tgattatctt ggtctctgcac 420
ctctggaaga cattgctgaa caaattttta atgcagctgg tccaagtgga agaaatacag 480
aatatctttt tgaacttgca aattctatta ggaaccttgt gccagaagaa gcagatgagc 540
atcttttcgc tttggaaaaa ttagtaaagg aacgtttaga agggaaacag aacctcaatt 600
gcatataaatt tagtcttcag agaattaact tcagtgcaca atgacaatat gatttgaaa 660
tacgtttact taaagatctt atttttaatt tagtgaggat attattttaa cttttatatt 720
aactggaaat gtccgaaac acatatttaa aatattggga tacagtgaat gaaaaattca 780
aattttaata acataaagat ttcctaactt tatgttattg aacacttact cactagaagt 840
gagttcttta gaaaaataca gtgaaggact cagttcagtc ttgtttttat cagagtgaat 900
atcatcctgt ttcacatccc aatactatct tgaaattcta aacaattaaa ccaaaattcc 960
aataaataata aggttatgcc ttcaatatat tcctatacaa ttctgtaacc atggtttaa 1020
atacacaagc ttaaaataac atgcttagaa atacacaata atatgaacag tatttcagcc 1080
ttaattgtga atttcttctg tattcaagta ttaaataaaa tcttttgagt ttttagccaa 1140
aaattggcat ttttaaaata cgaataattt cttggaatta taatgtactg tacctcttct 1200
tttttaaaata aaggcatttt actatatgga aaataactca cttaaagcata aattacatta 1260
tacaatcat gatcactaat gatgtagtct gtcattcact ttgtattaat cttataccaa 1320
aactgaaaaa gatgggctga tactacaaat taatggcaca tataatgaaa atttagtttt 1380
taaaacagct tttggaattc tttgtctgtc actatctcaa tttgtgtgtg tgtgtgtgta 1440
tatacataaa tatacatata aaattttttt tttctttgca gcctgcgtct ggccatccca 1500
caggctggaa agtgaacct ctggcagaag ccaagaacag gcacctcctg gaattataat 1560
tttgttttgt t

```

&lt;210&gt; 69

&lt;211&gt; 1549

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 044775CB1

&lt;400&gt; 69

```

cgcgcttagg caggcggtgg cgcggtgga gtgcccggg gagggctgtg ccggttgctt 60
tctgcagccg catctcggcc agctctcctc gccgtcccc gggcgctgtg cgtctccagt 120
ccgggaccga agccgcctgc cgtagcgggc ggccagatcc gcgtcccgcc tcagcggccg 180
gaggacatgc gggagagaga atgagccaga gggacacgct ggtgcatctg ttgcccggag 240
gatgtggtgg tacagtggga gctattctga catgtccact ggaagtgtga aaaacacgac 300
tgcagtcac tctgtgacg ctttatattt ctgaagtcca gctgaacacc atggctggag 360
ccagtgtcaa ccgagttagt tctcccggac ctcttcattg cctaaagggt atcttgaaa 420
aagaagggcc tcgttccttg ttttagaggac taggccccaa tttagtgggg gtagccctt 480
ccagagcaat atactttgct gcttattcaa actgcaagga aaagtgtgat gatgtattt 540
atcctgattc taccgaagta catatgattt cagctgcaat ggcaggcatt tactgtacat 600
ttctcccag tggactaact taaacctcgt tttactgcaa tcacagcaac caacccatt 720
ttagaataa agactcgggt acagcttgat gcaaggaacc gcggggaaa gcaaatgggt 780
tggtttataa gtgttcgtaa agtgatcag acagatggac taaaaggatt ttataggggc 840
gcttttgaat catatgctgg tatatcagag actgttatcc attttgttat ttatgaaagt 900
ataaaacaaa aactactgga atataagact gcttctacaa tggaaaatga tgaagagtct 960
gtgaaagaag catcagattt tgtgggaatg atgctagctg ctgccacctc aaaaacttgt 1020
gccacaacta tagcatatcc acatgaagtt gtaagaacaa gactacgtga agagggaaca 1080
aaatacagat ctttttttca gactctatct ttgcttgctc aagaagaagg ttatgggtct 1140
ctttatcgtg gtctgacaac tcacttagtg agacagattc caaacacagc cattatgatg 1200
gccacctatg aattggtggg ttacctactc aatggatagc agcacgagga ctgctgtact 1260
gcaaaaaaag aagacaaaaa gattacagtg gacctggga tacagaagcc agcatggcag 1320
acagaagaaa aatagtttgg gaacatgtaa ctattctaag tggaaagttt gttgtaggaa 1380
ttatagtaat cacaccacat tacttggcct ttcggtaatg tgaaaaaaa aaaaaaacct 1440
cagagcctcc aaggaaatgc ctttagaagc actcctctct caaaattgac attttctcta 1500
ccatgtcccc cagacacagt tgggttttgt tgatttatgg cagtcttct 1549

```

&lt;210&gt; 70

&lt;211&gt; 2237

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature



&lt;223&gt; Incyte ID No: 116588CB1

&lt;400&gt; 70

```

gtcactgggtg ccacggggcc tcagegcact tctgtcttaa gtccttgggc ctccttattt 60
tcccctttgc gatcgattcc agccacacct gtggatgttg ctagttactc cgcgtccgga 120
acgtggaggt cgagggactg agctgggcga gttttgtggc actcctttgc tcttcagcag 180
ctattttttgc tatgataatc ctgctgccct tcagactcaa gttaaaccgag atatgcaagt 240
gaataaccacg aaattcatgc tgctgtatgc ctggtattct tggcccaatg tagttttgtg 300
tttctttggg ggctttttga tagaccgagt atttggaaata cgtatggggca caatcatttt 360
tagctgcttt gtttgcattg gacaggttgt ttttgccctg ggtggaatat ttaatgcttt 420
ttggtctgat gaatttggaa gatttgtatt tgggattggg ggcgagtcct tagcagttgc 480
ccagaataca tatgctgtga gctggtttaa aggcaaaaga ttaaacctgg tgtttggact 540
tcaacttagc atggctagaa ttggaagtac agtaaaccatg aacctcatgg gatggctgta 600
ttctaagatt gaagctttgt taggttctgc tggtcacaca accctcggga tcacacttat 660
gattgggggt gtaacgtgta ttctttcact aatctgtgcc ttggctcttg cctacttggg 720
tcagagagca gagagaatcc ttcataaaga acaaggaaaa acaggtgaag ttattaaatt 780
aactgatgta aaggacttct ccttaccctt gtggcttata tttatcatct gtgtctgcta 840
ttatgttgct gtgttccctt ttattggact tgggaaagtt ttctttacag agaaatttgg 900
atthttcttcc caggcagcaa gtgcaattaa cagtgttgta tatgtcatat cagctcccat 960
gtcccgggtg tttgggctcc tgggtggataa aacagggaag aacatcatct gggttctttg 1020
cgcagtagca gccactcttg tgtccacat gatgtggcc tttacgatgt ggaacccttg 1080
gattgctatg tgtcttctgg gactctccta ctcattgctt gcctgtgcat tgtggccaat 1140
ggtggcattt gtagttcttg aacatcagct gggaactgca tatggcttca tgcagtccat 1200
tcagaatctt gggttggcca tcatttccat cattgtcgtg atgatactgg attctcgggg 1260
gtatttgttt ttggaagtgt tcttcattgc ctgtgttctt ttgtcacttt tatctgtggt 1320
cttactctat ttggtgaatc gtgccagggg tgggaacctt aattattctg caagacaaag 1380
ggaagaaata aaattttccc atactgaatg agaagttaaa atgaatgtgt catgagaatg 1440
ggcttaacac atcgttgggt tgaaaacttc catttttaaa aatttagagt ttagtcatta 1500
gaaaaaataa tggactggaa agttatattt atatccaaat atacctattt caaagtgtat 1560
ttgtgaggcc tgttttagcc tgtgtctttt gtattgtgtg ttgctaaaga attctacttt 1620
tagtaggcta atcaacaatg aaagggttag aaaattgctg tggaaacatcc aggtgaactt 1680
caggaaaagc agtgaaaaat ggaaaacgtt ggagcttctg ttgagataat cttcattagg 1740
tatatatctt agggatacag ccttttcttt atcttatagc aggaaaaaaa aacttttgag 1800
ggaaatagaa gggctcgtt acacaaaata aacaatggca ttgtcatagg ccttctcttt 1860
actagtaggg cataatgcta ggggaatatgt gaagatgttt ttatgaagtc tctttctgat 1920
cacgaacaat agcttgcgt ctactctgta gttatgtgga ttgccgagca atgacctttt 1980
tcaatttctt atttctgtgt tactgaggac cctaatacact tagggatgta attttatagt 2040
ataaactttc tgtacagttt ttcttatagt ctaataagta aaaagtgtcc ttcaaattat 2100
gataattgcc tatgtacatg gataaattaa aacactgcac acggagtaaa aaaaaaaa 2160
aaaaaaaaaa aaaaaatgag cggccgcaag cttattccct ttagtgaggg ttaattttag 2220
cttggcactg gccgtcg 2237

```

&lt;210&gt; 71

&lt;211&gt; 1114

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 875369CB1

&lt;400&gt; 71

```

tgggaacgga gcagccccgg gggccccctt gaggcggcga ggccgcgaag ggcgcggggc 60
tggagggtag gagagcgcgg gaaagcgccc cagacgccac tcgcggcgga cggcggccag 120
ttcccagggg tttggagcgg gcccgcggcg ccattggtca cgtcggctcc cgcaagcgct 180
cgaggagtcg cagccgggtcc cggggacggg ggtcggaaaa gagaaagaag aagagcagga 240
aagacacctc gaggaactgc tcggcctcca catcccaagg tcgcaaggcc agcacggccc 300
ctggggcgga ggcctcacct tctccctgca tcacagagag aagcaagcag aaggcccgga 360
ggagaacaag atccagctcc tctcctctct tctccagtgc gccggaagaa gcgggggaag tacaaggaca 480
cctcgctctc ctccctcttc tccagtgtat gcccgaagaa gaagggaag gagagggcgg 540
agaggaggaa gaagaagaag aagaggaaga agctgaagaa gaagggaag gagagggcgg 600
aagcacagca ggtggaggct ctgccgggcc cctcgctgga ccagtggcac cgtacagctg 660
gggaggaaga ggatggccca gtcctgacgg atgagcagaa gtcccgaatc caggccatga 720
agcccatgac caaggaggag tgggatgccc ggcagagcat catccgcaag gtggtggacc 780
ctgagacggg gcgcaccagg cttatttaagg gagatggcga ggtcctagag gaaatcgtaa 840
ccaaagaacg acacagagag atcaacaagc aagccaccgg aggggactgc ctggccttcc 900
agatgcgagc tgggttgctt ccctgagggc ccccgctggc caaggcctgt ggacgacgct 960
ggcggcccg cctgggcagg tttcagggtg ccagtgggaa gcctgatggg tgctggtggc

```

```

ctttcccccg tggattgggc tctggcccag cccagctctct tctcaggggc aggggggtgga 1020
gggttgggggc accggcctgc ttggcaccgc catctgaaaag agcagcactt ctcagctatt 1080
aaaggccccc tggatagaca aaaaaaaaaa aaaa 1114

```

```

<210> 72
<211> 998
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 1325518CB1

```

```

<400> 72
gactaaaaaa gccatgtatt ctttcgtttc tctctaaaag aagaaaaata taatttaaaa 60
atacattgcg tatttttctaa aacaataaat ttatagtgtt aatattcata gggccaatca 120
aaatgaagct tctccttttg gcctgcattg tatgtgttgc ttttgcaagg aagagacggg 180
tccccttcat tggtaggat gacaatgacg atgggcaccc acttcacca tctctgaata 240
ttccttatgg catacgggaat ttaccacctc ctctttatta tcgcccagtg aatacagtcc 300
ccagttacc tgggaatact tacactgaca caggggtacc ttcgtatccc tggattctaa 360
cttctcctgg attccccctat gtctatcaca tccgtggttt tcccttagct actcagttga 420
atgttcctcc tctccctcct aggggtttcc cgtttgtccc tccctcaagg tttttttcag 480
cagctgcagc acccgctgcc ccacctattg cagctgagcc tgctgcagct gcacctctta 540
cagccacacc tgtagcagct gagcctgctg caggggcccc tggtgcagct gagcctgctg 600
cagaggcacc tgttggagct gagcctgctg cagaggcacc tggtgcagct gagcctgctg 660
cagaggcacc tgttggagtg gagccagctg cagaggaacc ttcaccagct gagcctgcta 720
cagccaagcc tgctgccccca gaacctcacc ctctccctc tcttgaacag gcaaatcagt 780
gaaattctct agaagagtac catgggttca tttctatact gatgcagaaa taagtgaat 840
ctacaaaagt tttctttctt ttccaaagac tatttcattc tggtgtattc agagtattca 900
tctcactaca ttgatttgtt tgtggtagtt tttccttggc ctttaattat attgaaaaaa 960
cattgataat taaataaata aaatagataa ttttagaca 998

```

```

<210> 73
<211> 2348
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 2060987CB1

```

```

<400> 73
gggcccagccg gctcgcgccg gggccatggc agcagcggct actgcagccg aggggggtccc 60
cagtcggggg cctcccgggg aagtcaccca tctgaatgtg ggaggcaaga gattcagtag 120
ctctcgcagc actctcacct ggatcccaga ctccctcttc tccagttctc tgagcgggacg 180
catctcgacg ctgaaagatg agaccggagc aatcttcacg gacaggggacc ctacagttct 240
cgccccatc ctcaacttcc tgcgcaccaa agagtggat cccaggggtg tccacgggtc 300
cagcctcctc catgaagccc agttctatgg gctcactcct ctgggttcgtc gcctgcagct 360
tcgagaggag ttggatcgat ctcttctgtg aaacgtcctc ttcaatgggt acctgccgcc 420
accagtgttc ccagtgaagc ggcggaaccg gcacagccta gtggggcctc agcagctagg 480
aggacggcca gcccctgtcc gacggagcaa cacgatgcc cccaaccttg gcaatgcagg 540
gctgctgggc cgaatgctgg atgagaaaac cctccctca cctcaggac aacctgagga 600
gccggggatg gtgcgcctgg tgtgtggaca ccataattgg atcgtgtgtg cctataccca 660
gtttctagtc tgctacaggt tgaagggaagc ctctggctgg cagctgggtg tttccagccc 720
ccgcctggac tggcccatcg aacgactggc gctcacagcc cgggtgcatg gtggggcttt 780
gggtgaacat gacaagatgg tggcagcagc caccggcagc gagatcctgc tatgggctct 840
gcaggcgga ggcgggtggc ccgagatagg ggtctttcat ctgggggtgc ctgtggaggc 900
cttgttcttc gtcgggaacc agctcattgc tacaagccac acaggggcga tcgggggtgtg 960
gaatgccgtc accaagcact ggcaggtcca ggaggtgcag ccatcacca gttatgacg 1020
ggcaggctcc ttcctcctcc tgggctgcaa caacggctcc atttactacg tggatgtgca 1080
gaagtcccc ttgctcatga aagacaacga cctccttgtc agcagactct atcgggaccc 1140
agcggaggat ggggtcaccg ccctcaggtg ctacctcacc cccaagacca gtgacagtgg 1200
gaactggatc gagatcgct atggcaccag gtgcgggtca tcgtgcagca accgcagccc 1260
cccggagact gtgggctcgg ggcctcagct ctccagacc ttcactgtgc accgcagccc 1320
tgtcaccaag atcatgctgt cggagaagca cctcatctca gtctgtgccc acaacaacca 1380
cgtcgggaca tggctgtgta ctgccttccg cggcatgatt tccaccagc cgggtccac 1440
cccactcgct tctttaaaga tcttggtctc gatgggcag gcggctgcag 1500
tgctggcaat gacattggcc cctacggtga gcgggacgac cagcaagtgt tcatccagaa 1560

```

```

ggtggtgccc agtgccagcc agctcttcgt gcgtctctca tctactgggc agcgggtgtg 1620
ctccgtgcgc tccgtggacg gctcaccac gacggccttc acagtgtctg agtgccaggg 1680
ctcccgccgg ctccgctctc ggccccggcg ctacctgtct actggccagg ccaacggcag 1740
cttgcccatg tgggacctaa ccaccgccat ggacggcctc ggccaggccc ctgcagggtg 1800
cctgacggag caagagctga tggaaacagct ggaacactgt gagctggccc cgccggctcc 1860
ttcagctccc tcatggggct gtctccccag cccctcacc cgcattctccc tcaccagcct 1920
ccactcagcc tccagcaaca cctccttgtc tggccaccgt gggagcccaa gcccccgca 1980
ggctgaggcc cggcgccgtg gtggggggcag ctttgtggaa cgctgccagg aactgggtgcg 2040
gagtgggcca gacctccgac ggccaccac accagccccg tggccctcca gcggtctcgg 2100
cactcccttc acacctcca agatgaagct caatgaaact tccttttgaa caacgcagct 2160
gcatgatgc cttgggatgc cctggctcctg ggggactcag gtgcctccct gattcctgtg 2220
ggaaccccggt gttcaggggc agggcctcct tggaaataat ggttattgtt actaggctcc 2280
caccttccct cttttctgga agccaaagtc accctcccca ataaagtcct cactgccaaa 2340
aaaaaaaaa 2348

```

<210> 74  
 <211> 1139  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2172064CB1

```

<400> 74
ctcgagctgg gatgtgtggc aggttcctgc ggcggtgctt ggcggaggag agccggcgct 60
ccacccccgt ggggcgcctc ttgcttcccg tgctcctggg attccgcctt gtgctgctgg 120
ctgccagtgg gcctggagtc tatggtgatg agcagagtga attcgtgtgt cacaccagc 180
agccgggctg caaggctgcc tgcttcgatg ccttccaccc cctctccccg ctgcgttcc 240
gggtcttcca ggtcatcttg gtggctgtac ccagcgccct ctatatgggt ttcactctgt 300
atcacgtgat ctggcactgg gaattatcag gaaaggggaa ggaggaggag accctgatcc 360
agggacggga gggcaacaca gatgtcccag gggctggaag cctcaggctg ctctgggctt 420
atgtggctca gctgggggct cggcttgctc tggagggggc agccctgggg ttgcagtacc 480
acctgtatgg gttccagatg ccagctcctt ttgcatgtcg ccgagaacct tgccttggtg 540
gtataacctg caatctgtcc cgccctctct agaagaccat ttctctaaag accatgtttg 600
gagtcagcgg tttctgtctc ttgtttactt ttttggagct tgtgttctct ggtttgggga 660
gatggtggag gacctggaag cacaatatct cctcttctaa atacttctta acttcagaga 720
gcaccagaag acacaagaaa gcaaccgata gcctcccagt ggtggaaaacc aaagagcaat 780
ttcaagaagc agttccagga agaagcttag ccaggaaaaa acaaagacca gttggacca 840
gagatgcctg agttggagat gaactttggc caactttcct catcaccata cttaaaatcc 900
tgtccaagga ggagcttatt caccattttc tatacatgtg acatatgtag cagcataacc 960
gacaactggg actgcgctgc cttgactcca cctctacata caatgactca gctaaccaga 1020
ctaataaaag ccatgtttgc accattgtct agggaggcat tgctttgggg aattattccc 1080
agtgtcctcc ttacttatcg caagtaataa aaatccccct ggaaatcctc aaaaaaaaaa 1139

```

<210> 75  
 <211> 863  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2219267CB1

```

<400> 75
tcagagctag cttccgccgc agcctggcac ctcagggtgcc gaccttctac ctgccctact 60
tcttgaggcc cggcctggag gcggcagcct tgggtcttcc gtecttgacg gcagattgct 120
gtggacgccg ccccgctgct ctgctgggca ccatggctac aggcctggca tccctgctgc 180
tcttcgctgg ggcccagtat ctgccaggct ggactgtgct gttcctctct gtctggggc 240
tcttgccctc ccgggctgtg tccgcactca cgagcctctt cgcgggccgag gtcttccca 300
cggtgatcag gggggccggg ctgggcccgt tgctgggggc cgggttctct ggccaggcag 360
ccggcccttc ggacacctg cacggccggc agggcttctt cctgcaacaa gtctcttcg 420
cctcccttgc tgtccttgcc ctgctgtgtg tctgtgctgt gcctgagagc cgaagccggg 480
ggctgccccg gtcactgcag gacggccgac gcctgcgccc cteccactc ctgcggggc 540
gcccccgcca ggaccacct cctctgctgc cgccctccaa ctctactgg gccggccaca 600
cccccgagca gcactagtcc tgctgggtgg ccctgggagc caggatggga ccaaagtcaa 660
ggcctggggc atggctgagt accccagacg tctgggtccg ggcagacaca ttcctctcag 720
aagcccgtgt ctacgtgcag gtggagccgt ggggacagcg tgaagggtgc tccagccagg 780

```

ccccaggcac tgggagggccc tgggtctccc cccagccaca cccagtaggt gtggaggata 840  
aaggcttctg tggaaaaaaa aaa 863

<210> 76  
<211> 1322  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2308629CB1

<400> 76  
ccggggggccc ggcgcgcggg gaggcgcggg cctgcaggcc cccggtagca caagatccgg 60  
actccggccc ggactacgag gcgctgccgg ctggagccac tgtcaccacg cacatgggtg 120  
caggcgccgt ggcagggatc ctggagcact gcgtgatgta ccccatcgac tgcgtcaaga 180  
cccggatgca gactctacag cctgaccag ctgcccgtat tcgcaatgtg ttggaggccc 240  
tctggaggat tataagaacg gagggcctat ggaggcccat gagggggctg aacgtcacag 300  
caacaggcgc agggcctgcc cacgcccttt attttgctg ctacgaaaag ttaaaaaaga 360  
cattgagtga tgtaateccac cctgggggca atagccatat tgccaatggt gcggccgggt 420  
gtgtggcaac attacttcat gatgcagcca tgaacctgc ggaagtgggt aagcagagga 480  
tgcatagtga caactacca taccaccggg tgacagactg tgtacgggca gtgtggcaa 540  
atgaaggggc cggggccttt taccgcagct acaccacca gctgaccatg aacgttcctt 600  
tccaagccat tcacttcatg acctatgaat tcctgcagga gcactttaac cccagagac 660  
ggtacaaccc aagctcccac gtccctctctg gagcttgcgc aggagctgta gctgcccgag 720  
ccacaacccc actggacgtt tgcaaaacac tgcacaacac ccaggagtcc ttggcttga 780  
actcacacat tacaggacat atcacaggca tggctagtgc cttcaggacg gtatatcaag 840  
taggtggggg gaccgcctat ttccgagggg tgcaggccag agtaatttac cagatccct 900  
ccacagccat cgcattggtc gtgtatgagt tcttcaaata cctaactact aaaaggcaag 960  
aagagtggag ggctggcaag tgaagttaga ctgaacgaag ccagggggtc agatgacact 1020  
gctgcattct ggtcacattc tctgtctcct ggaatgctcc cacctcaagt ggagttagaa 1080  
ggaaggttaga ggggctctcc cccaggattt tgggtgtttg actaacacca gtctctgcca 1140  
acctctgttg ccaccacctt tccttcagg ccctaagcac gtgcagcaaa gcacaccaca 1200  
gcacctttga taacctctct ccatctctgg cctgatgacc tgccttagac tgttatagag 1260  
ggataagcag ctcatcccc tggttcttaa taaaaagcct ttaaattaaa aaaaaaaaaa 1320  
aa 1322

<210> 77  
<211> 1869  
<212> DNA  
<213> Homo sapiens

<220>  
<221> misc\_feature  
<223> Incyte ID No: 2660038CB1

<400> 77  
cagggacgct cattcctggc cagagtctgt acttctttct cggatcgaga tctcgttgct 60  
ggctcgagga aatcaccggc tcttctcccg gatccttttc ttctctcact tgcctggctt 120  
cttcttctct cctccctggc ttccatctcc caaccccgcg attccctcct ctactcccg 180  
gctcccgctg cccgccatcc tgagccatcc cacttgcaac cttctgtctt ttgcccctcc 240  
ttgacctcag agggctcctgc cttaagcttc tcaccagaat ctcttagatt tctatctctt 300  
cctgcttgc cagctcttga ctcccaaat ccagctgacg ttgaccact tgacatttga 360  
ccctgacacc cttgactgca aatctaaatt cttatcttct gcaacctgta ctgctgaaac 420  
agggccttccc cctgtcttcc aaccccgctt tctgacaccc atttttactt tcttactctt 480  
gggtcagttc ctgctacagc tatcccacca tggacttctt gatgagtggc ctggcagcct 540  
gcggggcctg tgtattcacc aatcccctgg aggtggtgaa gaccaggatg cagttgcaag 600  
gagaactgca ggcccctggc acataccagc ggcactaccg aatgtcttcc catgccttca 660  
tcaccatcgg caaggtggat ggccttgctg ccctgcagaa aggcctggcc cccgcctctt 720  
tgtaccagtt cctgatgaat ggcattccgac tgggaccta tgggctgggt gaggtgggg 780  
gttacctgca cacagccgaa gcgaccaca gtctgcccg cagcgcagca gctggggcca 840  
tggctggggg catgggagcc tacttgggga gccccatcta catggtgaag acacacctgc 900  
aggcacaggc agcctcagaa attgctgtag ggcaccagta taagcatcag ggcattgttc 960  
aggcgctaac cgagattggc cagaaacatg gtctggtggg gttatggcgt ggggctctgg 1020  
gcggcctgcc ccgagttatc gtcggttctt ccaccagct gtgcaccttc tcatccacca 1080  
aggacctcct gagccagtgg gagatcttcc ctccccagag ctggaagtgt gcgctggtg 1140  
ctgccatgat gagtggcatt gcagttgtct tggccatggc accctttgat gtggcctgca 1200  
caaggctcta caaccagccc acagatgcac agggcaaggg cctcatgtac cgggggatag 1260

```

tggacgctct gctgcagaca gctcggaccg agggcatttt tggcatgtac aagggtatag 1320
gtgcctccta ctccgcctc ggccccaca ccatectctc cctcttcttc tgggaccagc 1380
tgcgctccct ctactacaca gacactaaat aacagccgct tccccagtct ccaccaaag 1440
agcactcctt ggccacttgt gcctccacca ctatgtcctg gtgactactg attaggtgac 1500
ctttcatcca tccatggggg acagccaacc ccactcccca tctgttctca ggggtgaatc 1560
actacaagag atgagtttcc ctcttttctc tgggtgttgc tttaaacctt ccctacccat 1620
tccctgggta actcacaccc ctctctcagg gctgaacgag tcatcccaaa gtgtatttcc 1680
tcccactcac cactgccacc cttgagtcct tcctgtctcc atgcacagtt ttaaactcct 1740
ccctccaaaa ccaaagggaa tcgagagacc caattccag gcgtctggga cccaggtgtc 1800
ctgttagatt caaaggcaca gagattatat tgattataaa gcaagtttat tctgaaaaaa 1860
aaaaaaaaa

```

<210> 78  
 <211> 1881  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2670745CB1

```

<400> 78
gaagaaccga gcttggctgt gtttatctcg ttggggacta aggcgtcggg tggcgcgcaa 60
cgggttctag gctgcaggca gctcaggagc ccgcggcccc gcccgggctc ggccctggcag 120
atagcagagg cagcaggccg tgccgggggg gcatgttgct gtaaccagtg gccagggga 180
tgttacggtg gacagtgcac ctggagggcg ggccccgcag ggtgaaccat gctgcagtgg 240
ctgtcgggca tcgggtatac tccttcgggg gttactgtct tgggtaagac tatgagacac 300
tgcgctcagat agatgtgcac attttcaatg cagtgtcctt gcgttggaac aagctgcccc 360
cgggtgaagtc tgccatccgt gggcaagctc ctgtggtaac ctacatgcgc tatggacact 420
caaccgtcct catcgacgac acagtcctcc tttggggcgg gcggaatgac accgaagggg 480
cctgcaatgt gctctatgcc tttgacgtca atcagcctg tgcctaggc aagatcatgt 600
cagggacagt tcctggggcc cgggatggac actgttttcc caatgacatt cacaagctag 660
acatttttgg gggctacgag cagcaggcgg actgttttcc gtacaaaagg cagccctgca cgtggaggg 720
ataccagcac catgacatgg actcttatct gtacaaaagg acatgtatgt ctttgggggc cgtgccgacc 780
acttccactc agccacaatg ctgggaagtc tttactgcaa ccgcattcga gcttttgaca 840
gctttggggc attccattcc aacaatgaga gactgtcccc cgactccagt gctgcctgag gggcgccgga 900
ccagaactga ggcttggctg aatggggagc tgtacatctt tgggtggtat aatgcaaggc 960
gccaactggc cttccatgac ctctggaagt ttaatcctgt gtcctttacc tggaaaaaga 1020
tgaaccgaa ggggaagggg ccatgtcccc gccggcgcca gtgctgctgt attgttggtg 1080
acaagattgt cctcttttgg ggtaccagtc catctcctga ggaaggcctg ggagatgaat 1140
ttgaccttat agatcattct gacttacaca ttttggaact tagccctagt ctgaagactc 1200
tgtgcaaaact ggccgtgatt gactataacc tagaccagtc ctgtttgcct catgatata 1260
ggtgggagct gaatgccatg accaccaaca gcaatatcag tcgccccatc gtctcctccc 1320
atgggtagga ggaagtttct gccacctccc ctctctgagc tgctgtcatc ttcactgccc 1380
ctgcccactc gtcacccacc tgctccttgg acccctggac ttggtatacc tccatgtgga 1440
gttgttgggc gctctgtctg tgaattcagt ggggagctgt agcgggtggg 1500
gggctagggt cctccccctt tgggcccagg gcccttccc cttggtgtct tgctcccatc 1560
cacctccttt cagctgctcc tgggcctcag ctctgcccag ggccagccag gttctgtctg 1620
gaagggaagg gaatggggag aagggagaag caagcagtgt ctgagcctca ggagcttccc 1680
cctccccctt tgcctatccc ctcccctctg cttgagcctt gagccttgac tgggagctga 1740
aaggagtgtc agctgttggc atgagacctc cttctccccg tcttggggag gtggggacca 1800
gcagataaat cccacccttc cttgagctgt cgctgtactc tgaagttcag ccagctcaga 1860
ttttataaaa attaatataa a

```

<210> 79  
 <211> 2004  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 2676443CB1

```

<400> 79
aatcgagggg ggcgtggcct agcgggtccc ccccgggggc gcgcgcgcgc attggctgtg 60
cgggggtgagg gcgcgcgggg ggcgctttga accgggagcg gggcgcgggg cgcggggcgc 120
tgcgccgggt acacgccggg gtaggggcgg ggtcgggttg tggtcggggc gggattgggc 180

```

tctcctgggc	catggcagcc	gagggcgcgc	tgtcgcgctg	gtacttcggg	gggctggcct	240
cctgcggggc	cgcttgctgc	acgcacccgc	tggacctgct	caaggtgcat	ctgcagacgc	300
agcaggaggt	gaagctgcgc	atgacgggca	tggcgcgtgc	ggtgggtgct	accgacggca	360
tcctggcact	ctacagcggc	ctgagcgcct	cgctgtgcag	acagatgacc	tactccctga	420
ctcgggttcgc	catctacgag	actgtgcggg	accgcgtggc	caagggcagc	caggggcctc	480
tccccctcca	cgagaagggtg	ttgctgggct	ccgtcagcgg	tttagctgga	ggcttcgtgg	540
ggacgccccg	agacttggtc	aacgtcagga	tgcagaacga	cgtgaagctg	ccccagggtc	600
agcggcgcaa	ctacgccccat	gcgctggatg	gcctgtaccg	cgtagctcgt	gaagaggggtc	660
tcaggagact	gttctcgggt	gcaaccatgg	catccagccg	agggggcctta	gtcactgtgg	720
gccagctgtc	ctgtctacgac	caggccaagc	agctggctct	tagcaccggg	tacctctctg	780
acaacatctt	cactcacttt	gtcgcagct	ttattgcagg	tggatgtgcc	acgttcctgt	840
gccagccccct	ggatgtgctg	aagactcgcc	tgtatgaactc	caagggggag	tatcagggcg	900
ttttccactg	cgccgtggag	acagcgaagc	tcgggcctct	ggccttttac	aagggcctcg	960
tcccagctgg	catccgcctc	atccccaca	ccgtgctcac	ttttgtgttt	ctggaacagc	1020
tacgcaaaaa	ctttggcatc	aaagtgccat	cctgaccagc	cgtggggaatg	gctgggctgc	1080
caggccagac	acgctagggt	cttccaaaga	gtcccaagcc	cagcacctgc	tccctggggcc	1140
acgaacctcc	tggccgtggc	caccatcct	ccgcagcagg	ccccctgctgt	ccccccacct	1200
gctggctgag	ctcctcctgg	cctcgtcccc	tctcagctgt	agctgcacca	ccccgcctct	1260
ggctaccagg	ctctcccgcc	tgggcaactgc	gtggccttgc	ccctctcccc	ctggcagctc	1320
ctcaggggaa	caggggctac	cagaggctga	tttctccccct	ctcctggggc	aggggagggg	1380
tattatccct	gcctcctgcc	cccgatgccc	aaagcagcat	cttccagcac	tttccatcga	1440
ggacttgggt	ggcagagtgt	gggtgcagcc	tggctgttgc	tcacccaagt	gctagctctg	1500
cacttcgtgt	ctgctgagag	caaccagacc	ttccatgtcc	tcgggcagct	gcaactcccc	1560
gcgagacccc	gcagctgggt	gggatgaaca	agcaacgcag	accacaagcg	agtgcctggg	1620
agggagtggc	ccaggttggg	tctggagcca	ttgtgggtga	gggtcgaggg	ccaccgaggt	1680
cccgcgcacc	gctgcctgcc	ctgcagtggc	tttaacagtt	agttttgcc	aagcctctcc	1740
actcaccagc	aggcgtgtct	tgtcttcagg	gattgtgcct	gcgtccctcg	ggcacctggg	1800
cccccccgct	tggctccctg	ggggaatggc	ccaggcgggc	cgcgggttct	ccttagggcc	1860
ttctccccga	caaggagtcc	gacggggcgc	atgctgcac	ctctgcctcc	ctggctcgtg	1920
ggcttcaccc	cacctgggaa	gggcagtgtg	ctctgtgggg	gctgcaatca	ataaatgccg	1980
ggagctgcc	aaaaaaaaa	aaaa				2004

&lt;210&gt; 80

&lt;211&gt; 3555

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3295764CB1

&lt;400&gt; 80

gaccgtgggc	gagtcagaa	cgctcctggcc	ttacaggagg	aaggcgtcac	tcgcgggttac	60
aagtgcctga	ccctcactcc	agttggcgga	ggaggagaag	gaagggggcg	ggccgggtcc	120
cctccccctg	cgccccggat	ggatgtgccc	ggcccggtgt	ctcggcgggc	ggcggcgggc	180
gcggccactg	tgtcctctgc	gaccgctcgg	gtccgtcgcg	aatgctggtt	cttgcgcgac	240
gcgctgctct	gcgcctacgg	cttcttcgcc	agcctcaggc	cgtccgagcc	cttcctgacc	300
ccgtacctgc	tggggccgga	caagaacctg	accgagaggg	aggtcttcaa	tgaatttat	360
ccagtatgga	cttactctta	cctgggtgcta	ctgtttcctg	tgttccttgc	cacagactac	420
ctccgttata	aacctgttgt	tctactgcag	gggtcagcc	ttattgttac	atggtttatg	480
ctgctctatg	cccagggaact	gctggccatt	caatttctag	aattttttta	tggcatcgcc	540
acagccactg	aaattgccta	ttactcttat	atctacagtg	tgggtggacct	gggcatgtac	600
cagaaagtca	caagttactg	tcgaagtgcc	actttggtgg	gctttacagt	gggctctgtc	660
ctagggcaaa	tccttgtctc	agtggcaggc	tggctcgtgt	tcagcctgaa	tgtcatctct	720
cttacctgtg	tttcagtggc	tttctcctgc	gcctgggttt	tacctatgcc	acagaagagc	780
ctcttctttc	accacattcc	ttctacctgc	cagagagtga	atggcatcaa	ggtacaaaat	840
ggtggcattg	ttactgacac	cccagcttct	aaccaccttc	ctggctggga	ggacattgag	900
tcaaaaatcc	ctctaaatat	ggaggagcct	cccggtggag	aaccggaacc	caagccagac	960
cgtctccttg	tattgaaagt	actatggaat	gatttctcta	tgtgctactc	ctctcgccct	1020
cttctctgct	ggtctgtgtg	gtgggcccctc	tctacctgtg	gctattttca	agttgtgaac	1080
tacacacagg	gcctgtggga	gaaagtgtatg	ccttctcgct	atgctgctat	ctataatggg	1140
ggcgtggagg	ccgtttcaac	cttactgggt	gctgttgctg	tgtttgcagt	tggttatata	1200
aaaatatcct	ggtcaacttg	gggagaaaatg	acattatctc	tcttttctct	cctgattgct	1260
gctgcagtgt	atatcatgga	cactgtgggt	aacatttggg	tgtgctatgc	atcctatggt	1320
gtcttcagaa	tcattctacat	gttactcatc	acgatagcaa	cttttcaaat	tgtgcaaac	1380
ctcagcatgg	aacgctatgc	cctagtattt	ggtgtaataa	ccttcattgc	cctggcactg	1440
cagacgctgc	tcactctaata	tgtggtagat	gccagtggcc	ttggattaga	aattaccact	1500
cagtttttga	tctatgccag	ttattttgca	ctcatcgctg	tggtttttct	ggccagtggg	1560

```

gcagtcagtg ttatgaagaa atgtagaaag ctggaagatc cacaatcaag ttctcaagta 1620
accacttcat aatatactgc tgaagggtc ctcttctatag caagaactct gcacagcaac 1680
tgcctggatg tatttgattt tttaaagcgt agacatatat ttatgaatgt gcatttcttg 1740
acttcacagc agccacttga ctaatacctt gtgttccggg aataacatga tactattcag 1800
aggagccaga agtaaaagttt atttcatgga ttatttatga gagctaattt aaggatgact 1860
ttttttctga ttcaaaagtg aacttgattt taaaaaccag tcaagagcaa tcaaagcagc 1920
acatggtgtt gtatacttca ttagcaagtg agtttggtgt tttataggtc acatatgtct 1980
gtatctactt agccagatgc ttggcctggg gggaccaggg ctccacagag gccacaaaat 2040
gttgcaagtc atgatggatg gaaatatgtt ctaacagcat ctgcctctat tcaatttaat 2100
tcttatttct gtgttactca tgtacattgg tcttctaca tagttattct atcactggca 2160
atatttgttc tggtttagtg ttctgtattt taaggtgtac gtatcatttc taattttaag 2220
ttattttaaa aaaattcatc atatgaatgt tcttggttcc cattgtgacg attatttatt 2280
tctgtaaaaa ttgtttagaa gtacgttttt gcattattca tatgcttccc agagaagctc 2340
atthagttag aaaataaggc aagttttgaa gcttgctaaa tgaagagact taagaaagct 2400
taagggtacgc ttgcttgtct ttaaactctc aatatgaagg actattaatt ccaagattaa 2460
aagttcatat ataggctaaa gatgtaacta ggccatttgt atttgtattc ccttttattt 2520
ccaaaataaa atgaaaaatc ttttttaat aatttcatcc ctatttatag tttttatatt 2580
aatttgtttt tcttatccaa gtaaagatgt caataggaat tgcattagtc caaggccttt 2640
ttcataaaact gagcctcttt tcaattattt caatgggaca ggaactagga tagatgtgat 2700
tctgcatttt ttttacctta aatctgcctt tgtttctaaa ggtagatcat cttgaatatt 2760
tgcttaaaat tgctagtgat ttcatocca agttacttga aaaaatgttc tatatgcatt 2820
taattctgaa atcagttctac caaggggtcg ctagtatatg tcagacatga aaactatttt 2880
aaagctgact ttggtgcctt atcttgaaaa gaatctagat aggtgctttt aactggggta 2940
ttaacttttt tagaatgaca cagctgaaca gtgttaataa tagtgtgtca agattgcaaa 3000
gtcgacatac tcatttggtt taagcaggaa tcttagaagc aaatggatgg ggataagaat 3060
aggtcatttt ctattcacca tctttacta ttcaagggaaa ggaaaagaac actagctaa 3120
gaagggaaaag ggaagtgata tcataaaagt agcaaccttc attttacatt ctgtctgttt 3180
tctttttttt gctttgtttt gtttgtgtca atttgggaat tgtgtactcc gaaacaagta 3240
gaaaagtgtc gtttgaggga ttttattaaa tcttttttta atggaatgtg gtacaaaatt 3300
ttcatgttac caaagcaata tttccctgga atttaattca aagtttggg catacaacct 3360
gagccttttc ttatataaga caagaatatg ttcacatctt ggtatgtggc catatttata 3420
gaatgctgaa ctcaatgtgc aagttgtact gtatgcagtt ttgtaataaa gtgaaaataa 3480
tttgtgttac tttttattca attctgtata gattataaaa ttatttttat taaataaata 3540
ttttacagta ttttt 3555

```

&lt;210&gt; 81

&lt;211&gt; 1293

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;223&gt; Incyte ID No: 3438320CB1

&lt;400&gt; 81

```

atatgcccca cgcggggactc atactacgtt tcccgtgaac acgtgcagtc caaaccccg 60
ccctgatatt tatctcagtg gacgggtggc ggaagggac aatgggtttcc atgtcagcgg 120
ataaacgctc tcccctcggc tcccggacgc gacggaggtc gtagtagtag tgagtacgtg 180
ctgaggagca aaggagtaac caagagatcc agtgaccgac agagcaagag ccatgcccg 240
ccggggcctg gtggctgggc cagacttgga gtattttcag cgtcgctatt tcacgccggc 300
ggagggtggc caacataaca ggcccgaaga cctctgggta tcttacctgg gacgcgtgta 360
cgacctaacg tcattggcac aggaatacaa ggggaacctg ctgctgaaac ccacgtgga 420
agttgcaggc caggatatca gccactggtt tgatccaaag accagagaca tccgcaagca 480
catagatccg ctgaccggct gcctgaggta ctgcaccccg cggggccgct tbtgtcacgt 540
tccgcctcag ctgccctgtt cggaactggg caacgatttt gggagccct ggtggcaggg 600
gtcgtattat gaggtggggc ggctgtctgc caagaccgg agcatccgca tcattaacac 660
gctcacgtcg caggagcaca cactggagggt gggggttctg gagtccatat gggaaatcct 720
acaccgctat ctcccctata actcacatgc tgccagctac acgtggaaat atgaagggaa 780
gaacctgaac atggatttta ccctggaaga gaatgggac cgggatgagg aggaagaatt 840
tgactatctc agtatggacg gtacacttca cacacctgca atacttctgt acttcaatga 900
tgatctcacg gagttgtagg caaggagatg tacactcgtg tagactcaag acgtatttctg 960
agtttggctt tttctgtgcc ttgaggaaaa gtgggtggggc cgaggggtgc ctggaccagg 1020
atctccactc ctctccagga gctagcctgt cccttctga agtgtaaagg ccctattccc 1080
tgcttcattt acagtttgct ctgagaaaaat tagtgaatta atctttggga atgatacaag 1140
aagatcaagt accttggttt agggagatgt agaagaggat agtcagagtt caggcagaac 1200
tgtttgatag ttaagagaga gtatgtctac aggggtgagg gatggaagga cttttttggc 1260
aatgatggaa atgagatgtc tgcaggagat ggg 1293

```

<210> 82  
 <211> 1489  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 3986488CB1

<400> 82  
 cggggcgctggc gggcgggcgct cttctacggg acctgctcct tcctcatcgt gcttgtcaac 60  
 aaggcgctgc tgaccaccta cgggtttcccg tcaccaattt tccttggaat tggacagatg 120  
 gcagccacca taatgatact atatgtgtcc aagctaaaca aaatcattca cttccctgat 180  
 tttgataaga aaattcctgt aaagctgttt cctctgcctc tcctctacgt tggaaaccac 240  
 ataagtggat tatcaagcac aagtaaatta agcctaccga tgttcaccgt gctcaggaaa 300  
 ttcaccattc cacttacctt acttctggaa accatcatatc ttgggaagca gtattcactc 360  
 aacatcatcc tcagtgtctt tgccattatt ctgggggctt tcatagcagc tgggtctgac 420  
 cttgcttttta acttagaagg ctatatTTTT gtattcctga atgatatctt cacagcagca 480  
 aatggagttt ataccaaaca gaaaatggac ccaaaggagc tagggaaata cggagtactt 540  
 ttctacaatg cctgcttcat gattatccca actcttatta ttagtgtctc cactggagac 600  
 ctgcaacagg ctactgaatt caaccaatgg aagaatgttg tgtttatcct acagtcttctt 660  
 ctttctctgtt ttttgggggt tctgctgatg tactccacgg ttctgtgcag ctattacaat 720  
 tcagccctga cgacagcagt ggttggagcc atcaagaatg tatccgttgc ctacattggg 780  
 atattaatcg gtggagacta catTTTTctt ttgttaaact ttgtagggtt aaatatTTgc 840  
 atggcagggg gcttgagata ttctTTTTta acactgagca gccagttaaa acctaaacct 900  
 gtgggtgaag aaaacatctg tttggatttg aagagctaaa gagtctgcag caggattgga 960  
 gactgacttg tgactgcggg ctggggggggc attcccagta ggaatgtgaa gccagaggtt 1020  
 tcggattcgt gacatccacc ccctgggcaa gtgagagcat ctgcaaaatg caaagagaaac 1080  
 tacctcatat gcaggatgag ccaatggcag tctcaagaaa tgtactcggg cgacacctta 1140  
 cctgtggaata gcaaactctt tcaaaaataag ccactgggac tcggtagggtg gagccccagc 1200  
 tgctcttcta gggacctatg gggccttctg ggcattctctg tgctgtgtgc tggggaggag 1260  
 gttgatgtaa tgggtgactct tttctgatca gcaccttggc cgtgattccc aagggtccag 1320  
 ccaaagcaaa gggccagttg tttcagttta aacagacatg tctttagtct aataaaatta 1380  
 gtaactgcc agtaaagtta tttgttagct ttgatgaaag ctatgttggg atctttccct 1440  
 aatcatcaaa gtaataaaaa aatcatttct atgtaaaaaa aaaaaaaaaa 1489

<210> 83  
 <211> 927  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 4378816CB1

<400> 83  
 ctccttgctt tcctctgccg catggctcctg ggccgttggc gtcggaagcc tgaagcatgg 60  
 gcgctgagtg ggagctgggg gccgaggctg gcggttcgct gctgctgtgc gccgcgctgc 120  
 tggcgccggg ctgcgccctg ggcctgcgcc tggccgcggg gcagggggcg gcggaccgcg 180  
 gggcgctcat ctggtctctg tacgacgcgc tgggtgcaact cgcgctggaa ggcccttttg 240  
 tctacttgct tttagtagga aacgttgcaa attccgatgg cttgattgct tctttatgga 300  
 aagaatatgg caaagctgat gcaagatggg tttattttga tccaaccatt gtgtctgtgg 360  
 aaattctgac cgtcgccctg gatgggtctc tggcattgtt cctcatttat gccatagtca 420  
 aagaaaaata ttaccggcat ttcttgacga tcaccctgtg cgtgtgagag ctgtatggct 480  
 gctggatgac cttctccca gagggtctca ccagaagccc caacctcaac accagcaact 540  
 ggctgtactg ttggctttac ctgtttttt ttaacgggtg gtgggttctg atcccaggac 600  
 tgctactgtg gcagtcattg ctagaactca agaaaatgca tcagaaagaa accagttcag 660  
 tgaagaagtt tcagtgaact ttcaaaacca taacacccat tatctaactt catgaaccag 720  
 aatgaatcaa atctttttgt ttggccaaaa tgtaatacat tccagtctac actttgtttt 780  
 tgtattgttg ctctgaaca acctgtttca aattggtttt aaggcgacca gtttctgttg 840  
 tattgttgtt caattaaatg gtgatatagg gaaaagagaa caaatttgaa tttgtaataa 900  
 taaaatgttt aattatacaa aaaaaaa 927

<210> 84  
 <211> 970  
 <212> DNA  
 <213> Homo sapiens



<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 4797137CB1

<400> 84

ggatgcagca	gagaggagca	gctggaagcc	gtggctgcgc	tctcttccct	ctgctgggcg	60
tctgttctt	ccaggggtgt	tatatcgtct	tttccttggg	gattcgtgca	gatgcccatg	120
tccgaggtta	tggtggagaa	aagatcaagt	tgaaatgcac	tttcaagtca	acttcagatg	180
tcactgacaa	gcttactata	gactggacat	atcgccctcc	cagcagcagc	cacacagtat	240
caatatttca	ttatcagtct	ttccagtacc	caaccacagc	aggcacattt	cgggatcggg	300
tttcctgggt	tggaatgta	tacaaagggg	atgcatttat	aagtataagc	aaccctacca	360
taaaggacaa	tgggacattc	agctgtgctg	tgaagaatcc	cccagatgtg	caccataata	420
ttcccagac	agagctaaca	gtcacagaaa	gggggttttg	caccatgctt	tctctgtggt	480
cccttcttct	catccttgct	tttgtgccct	cagccgtggt	gggtgctctg	ctgctgggtg	540
gaatggggag	gaaggctgct	gggctgaaga	agaggagcag	gtctggctat	aagaagtcac	600
ctattgaggt	ttccgatgac	actgatcagg	aggaggaaga	ggcgtgtatg	gcgaggcttt	660
gtgtccgttg	cgctgagtgc	ctggattcag	actatgaaga	gacatattga	tgaaagtctg	720
tatgacacaa	gaagagtcac	ctaaagacag	gaaacatccc	attccactgg	cagctaaagc	780
ctgtcagaga	aagtggagct	ggcctggacc	atagcgatgg	acaatcctgg	agatcatcag	840
taaagacttt	aggaaccact	tatttataga	ataaatgttc	ttgtgtatt	tataaactgt	900
tcagggaagtc	tcataagaga	ctcatgactt	cccctttcaa	tgaattatgc	tgtaattgaa	960
tgaagaatc						970

<210> 85  
 <211> 594  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5470806CB1

<400> 85

gacaggatgg	cttcccttcg	cctgttcctc	ctctgcctcg	ctgtactgat	atttgcgtct	60
gaagctggcc	ctgggggtgc	tggagaatcc	aagtgtcctc	tgatgggtcaa	agtcctggat	120
gctgtccgag	gcagccctgc	tggtgatgtg	gccgtgaaag	tgttcaaaaa	gactgcagac	180
ggaagctggg	agccgtttgc	ctctgggaag	accgccgagt	ctggagagct	gcacgggctc	240
accacagatg	agaagttcac	ggaaggggtg	tacagggtag	aactggacac	caaatcgtag	300
tggaaggctc	ttggcatttc	cccattccat	gaatacgcag	aggtgggttt	cacagccaat	360
gactctggtc	atcgccacta	caccatcgca	gccctgctca	gcccgtactc	ctacagcacc	420
actgctgtcg	tcagtaaccc	ccagaactga	gggaccacgc	ccagtaggac	caggatcttg	480
ccaaagcagt	agcttcccat	ttgtactgaa	acagtgttct	tgctctataa	accgtgttag	540
caactcggga	agatgccgtg	aaacgatctt	attaaaccac	ctgtgatgcc	aaaa	594

<210> 86  
 <211> 618  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> misc\_feature  
 <223> Incyte ID No: 5473242CB1

<400> 86

gtgttgactc	gcaacctcag	gaacagacac	catggtgcac	ctaactgatg	ctgagaaggc	60
tactgttaat	ggcctgtggg	gaaaggtgaa	ccctgttgaa	attggcgctg	agtcccttgc	120
cagtctgctg	attgtctacc	cttggaccca	gaggtacttt	tctaaatttg	gggacctgtc	180
ctctgtctct	gctatcatgg	gtaaccccca	ggtgaaggcc	catggcgaaa	aggtgataaa	240
cgccttcgat	gatggcctga	aacacttgga	caacctcaag	ggcacctttg	ccagcctcag	300
tgaactccac	tgtgacaagc	tgcatgtgga	tcttgagaac	ttcaggctcc	tgggcaatat	360
gattgtgatt	atgatgggcc	accacctggg	caaggaattc	accccgagtg	cacaggctgc	420
cttccagaag	gtgggtggctg	gagtgccag	tgccctggct	cacaagtacc	actaaacctc	480
tttctctgct	cttgtctttg	tgcaatggct	aattgttccc	aagagagctt	ctgtcagttg	540
ttgtcaaaat	gacaaagacc	tttgaaaatc	tgctctacta	attaaagcat	ttgggttcaag	600
tgttctgttg	agataccc					618